

# MATHEMATICS

## STUDENT TEXT BOOK

### GRADE 7

This book was developed by Addis Ababa city Administration Education Bureau and adapted by Southern Nations, Nationalities and peoples' Regional Education Bureau.

#### **Adapted by**

Natan Labiso (MSc)

Meseret Ayenew (MA)

#### **Editted by:**

Firehiwot Eyuel (Bsc)

#### **Designer:**

Getahun Tadesse



**Southern Nations, Nationalities and peoples' Regional  
Education Bureau**

*This book was developed by Addis Ababa city dministration Education Bureau and adapted with budget and technical support of Southern Nations, Nationalities and peoples' Regional Education Bureau. The printing cost is covered by SNNPREBs.*

*Copy Right at © 2015 E.C Southern Nations, Nationalities and peoples' Regional Education Bureau .*



# Table of Contents

<b>UNIT 1: BASIC CONCEPTS OF SETS .....</b>	<b>1</b>
<b>Introduction.....</b>	<b>1</b>
<b>1.1. Sets and Elements of Sets.....</b>	<b>1</b>
<b>1.2. The Relation among sets .....</b>	<b>8</b>
<b>1.3. Operations on Sets .....</b>	<b>11</b>
<b>Summary on Unit 1.....</b>	<b>15</b>
<b>Review Exercises on Unit 1 .....</b>	<b>15</b>
<b>Unit 2: INTEGERS .....</b>	<b>17</b>
<b>Introduction .....</b>	<b>17</b>
<b>2.1. Revision of whole and natural numbers.....</b>	<b>17</b>
<b>2.2. Introduction to integer.....</b>	<b>20</b>
<b>2.3. Comparing and Ordering Integers.....</b>	<b>26</b>
<b>2.4. Addition and subtraction of Integers.....</b>	<b>33</b>
<b>2.5. Multiplication and Division of Integer numbers.....</b>	<b>42</b>
<b>2.6. Even and Odd integers.....</b>	<b>54</b>
<b>Summary on Unit 2.....</b>	<b>60</b>
<b>Review Exercise for unit 2.....</b>	<b>61</b>
<b>Unit 3: RATIO, PROPORTION AND PERCENTAGE.....</b>	<b>64</b>
<b>Introduction .....</b>	<b>64</b>
<b>3.1. Ratio and proportion.....</b>	<b>65</b>
<b>3.2. Revision on Percentages.....</b>	<b>81</b>
<b>3.3. Application of Ratio, Proportion and Percentage.....</b>	<b>93</b>
<b>Summary for unit 3.....</b>	<b>113</b>
<b>Review exercise for unit 3.....</b>	<b>115</b>
<b>Unit 4: LINEAR EQUATIONS.....</b>	<b>118</b>
<b>Introduction.....</b>	<b>118</b>
<b>4.1. Algebraic terms and expressions.....</b>	<b>119</b>



4.2 Solving linear equations.....	126
4.3. Cartesian coordinate system.....	137
4.4.Applications of linear Equations.....	151
Summary for unit 4.....	156
Review Exercise for unit 4.....	158
<b>Unit 5: PERIMETER AND AREA OF PLANE FIGURES .....</b>	<b>160</b>
Review Exercise for unit 5.....	161
5.1.Revision of triangles.....	161
5.2.Four - sided Figures.....	165
5.3.Areas and perimeters of triangles .....	177
5.4.Perimeters and Areas of four sided figures.....	182
5.5.Circumference and Area of a circle.....	192
5.6.Applications of Perimeter and Area of Plane Figures.....	197
Summary for unit 5.....	201
Review exercise for unit 5.....	202
<b>Unit 6: CONGRUENCY OF PLANE FIGURES .....</b>	<b>204</b>
Review Exercise for unit 6 .....	204
6.1.Congruent of Plane Figures.....	204
6.2.Applications.....	219
Summary for unit 6.....	223
Review exercise for unit 6.....	224
<b>Unit 7: DATA HANDLING .....</b>	<b>226</b>
Review Exercise for unit 7.....	226
7.1.Organization of data using frequency table.....	226
7.2. Construction and Interpretation of line graph and pie charts .....	231
7.3.The Mean, Mode, Median and Range of Data.....	241
7.4.Application of Data Handling.....	248
Summary for unit 7.....	252
Review exercise for unit 7.....	253

# UNIT 1

## BASIC CONCEPTS OF SETS

### *Learning Outcomes:*

**At the end of this unit, learners will be able to:**

- understand the concept of set.
- describe the relation between two sets.
- perform two operations (intersection and union) on sets.

### *Introduction*

The idea of a set is familiar in everyday life. In your surrounding there are different groups of objects or individuals. For example group of all grade 7 students in your school, group of all teachers in your school is a collection of individuals. In this unit you will learn about some sets and properties of sets.

### ***1.1. SETS AND ELEMENTS OF SETS***

In our daily life things are grouped together with a certain property in common such as family members, a collection of clothes, fingers of hands, collection of students in a class, a herd of cattle, a flock of sheep, a swarm of bees, etc. all these groups are well defined.

**Definition:**

***A Set is a well-defined collection of objects or individuals. The objects in the set are called its elements or members of the set.***

By well-defined, we mean that, given a collection and an object or individual, we have to say that the object or the individual is in the collection or not without any ambiguity. Repeating elements in a set do not add new elements to the set.

***For Example,  $\{b, b, b\}$  is the same as  $\{b\}$ .***

**Example 1.1**

***Identify each of the following collections are a set or not***

- A. The collection of whole numbers less than 6.
- B. The group of short students in a certain school.
- C. The collection of vowel letters in the English alphabet.
- D. A group of rich people in Hawassa.

**Solution:**

- A. The collection of whole numbers less than 5 is a well-defined collection. The whole numbers 1, 2, 3, 4, and 5 are in collection and anything other than these is not in the collection.*
- B. The group of short students in a certain school is not well-defined. It is because shortness is relative someone may be short for some group, but not for others.*
- C. The vowels in the English alphabet are a, e, i, o, u are well-defined; anything other than these five letters is not in the collection.*
- D. The group of rich people in Hawassa is not well-defined because richness is relative*

*Note:*

1. Sets are denoted by capital letters like  $A, B, C$ , etc, and elements of a set are denoted by small letters like  $a, b, c, d, x, y, z$ , etc.
2. The symbol  $\{ \dots \}$  is used to group the members of a set called braces or curly brackets with elements separated by commas.
3. Given a set  $A$  and an object  $x$ ,
  - i. If  $x$  is an element of  $A$ , then we denote this relation by " $x \in A$ " and read as " $x$  is an element of set  $A$ " or " $x$  belongs to set  $A$ ".
  - ii. If  $x$  is not an element of  $A$ , then we denote the relation by " $x \notin A$ " and read as " $x$  is not an element of set  $A$ " or " $x$  does not belong to set  $A$ ".

### Example 1.2

If  $A$  is the set of natural numbers less than 5, then  $A$  can be written as  $A = \{1, 2, 3, 4\}$ . In this case,  $1 \in A$ ,  $2 \in A$ ,  $3 \in A$ , and  $4 \in A$ , but  $5 \notin A$  and also  $7 \notin A$  and so on. Every object in a set is unique. The same object cannot be included in the set more than once.

For example, if  $A = \{1, 2, 3, 4, 5\}$  then every element of  $A$  is included not more than once. We do not write

$$A = \{1, 1, 2, 3, 4, 5\} \text{ or}$$

$$A = \{1, 2, 2, 3, 4, 5\} \text{ or}$$

$$A = \{1, 2, 3, 3, 4, 5\} \text{ and so on.}$$

### Example 1.3

**Determine the set  $X$  as the set of all days in a week.**

**Solution:**

*There are seven days in a week: Monday, Tuesday, Wednesday, Thursday, Friday, Saturday and Sunday.*

*Therefore =  $\{\text{Monday, Tuesday, Wednesday, Thursday, Friday, Saturday, Sunday}\}$ .*

**Example 1.4**

Let  $M$  be the set of multiples of 2 between 1 and 9; determine the set  $M$  and members of the set  $M$ .

**Solution:**

$M = \{2, 4, 6, 8\}$  and 2, 4, 6 and 8 are the members of set  $M$ .

**Exercise 1.1**

1. Which of the following collections are sets? Justify your answer.

- A. The beautiful birds in Hawassa.
- B. The natural numbers less than 1000.
- C. The prime factors of 2500.
- D. The months of a year.
- E. Students in grade 6 that are 12 years of age in Ethiopia.

2. Identify each of the following statements are true or false

- A.  $-1 \in \{-2, -1, 0, 1, 2\}$
- B.  $b \in \{a, b, c, d, e, f\}$
- C.  $6 \notin \{\text{factors of } 48\}$
- D.  $3 \notin \{1, 2, \{3\}, 4\}$ .
- E.  $\{3\} \in \{3\}$ .

**1.1.1. WAYS TO DESCRIBE SETS****i. Listing or Roster Method**

Listing or roster notation also called enumeration method is a list of elements separated by commas enclosed in curly braces. The curly braces are used to indicate that the elements written between them belong to that set. The elements in such types of sets can be completely or partially listed and the elements follow certain pattern.

**Example 1.5**

Let  $P$  be the set of all prime numbers less than 9. Write  $P$  using listing method.

**Solution:**

The prime numbers less than 9 are 2, 3, 5 and 7.

Therefore,  $P = \{2, 3, 5, 7\}$  is tabulation or complete listing method.

### **Example 1.6**

If  $A$  is the set of natural numbers less than 10, then write  $B$  using listing or roster method.

### **Solution:**

The natural numbers less than 10 are numbers from 1 to 9. Instead of writing all the 9 numbers, you can list the first three numbers to indicate that the numbers follow a certain pattern followed by three dots to indicate that the numbers are continuing up to the last number, which is 9.

Thus,  $A = \{1, 2, 3 \dots 9\}$  is a partial listing method.

### **ii. Verbal Method**

There are times when it is not practical to list all the elements of a set. In such cases, it is better to describe the set using verbal method.

### **Example 1.7 Describe each of the following sets**

- A. The set of letters in the English alphabet.
- B. The set of Grade 6 students in Ethiopia in 2014 E.C.

### **Solution:**

- A. There are 26 letters in the English alphabet and instead of listing all the English alphabets, you can write the set as  $\{\text{the English alphabets}\}$ .
- B. It is very difficult to list all Grade 6 students in Ethiopia in 2014 E.C. and the better way to write it is as  $\{\text{Grade 6 students in Ethiopia in 2014 E.C.}\}$ .

### **iii. Set builder method**

It is called method of defining property.

It is another way to specify a set that enables us to decide whether or not any given objects belong to the set.

### Example 1.8

A. F is the set of all females who are living in Hawassa can be described as :-

$$F = \{x/x \text{ is a female living in Hawassa}\}$$

which is read as “F is the set of x such that x is a female living in Hawassa”.

B. Let  $A = \{2, 3, 4, 5, 6\}$  can be described as  $A = \{x | x \in \mathbb{N} \text{ and } 2 < x < 7\}$

### Exercise 1.2

1. Describe each of the following sets by the listing method.

- A. The set of prime numbers between 4 and 25.
- B. The set of capital cities of the regional states in Ethiopia.
- C. The set of all subjects that a grade 7 student is learning.
- D. The set of even natural numbers.

2. Describe each of the following sets by a verbal method

- A.  $\{1, 2, 3, \dots, 100\}$
- B.  $\{1, 3, 5, 7, \dots\}$
- C.  $\{\text{January, February, March, } \dots, \text{December}\}$
- D.  $\{2, 3, 5, 7, 11, \dots\}$

## 1.1.2. SOME SPECIAL TYPES OF SETS

### Empty set

**Defnition:-**

*A set which contains no element is called empty set (null or void). Empty set is denoted by  $\emptyset$  or  $\{\}$ .*

### Example 1.9

- A. A set of whole numbers that is odd and even.
- B. A group of birds which have four legs.

- C.  $X$  is the set of whole numbers that are even and prime.  
 D. The set of cats that can fly.

**Solution:**

- A. There is no whole number which is both even and odd. So  $A$  is empty set, that is,  $A = \{ \}$  or  $\emptyset$   
 B. There is no bird which has four legs.  
 C. The only whole number that is both even and prime is 2.  
 Therefore,  $X = \{2\}$  is not empty set.

**Definition:-**

**A set  $G$  is said to be a finite set if  $G$  contains exactly  $n$  elements for some positive integer  $n$  or  $G = \emptyset$ . A set that is not finite is called infinite set.**

**Notation:**

If  $G$  is a finite set, then the number of elements in  $G$  is denoted by  $n(G)$ .

**Example 1.10**

**Identify each of the following sets as finite or infinite.**

- A.  $M$  is the set of all weeks with 9 days.  
 B.  $L = \{1, 2, \dots, 200\}$ .  
 C.  $K$  is the set of all natural numbers that are multiples of 3.

**Solution:**

- A. For every week there is 7 days. Therefore,  $M = \{ \}$  and hence  $M$  is a finite set, with  $n(M) = 0$ .  
 B.  $L$  contains 200 elements. Thus,  $L$  is a finite set and  $n(L) = 200$ .  
 C.  $M = \{4, 8, 12, \dots\}$  and  $M$  does not contain finite elements. Therefore,  $M$  is an infinite set.

D.  $K = \{3, 6, 9 \dots\}$  and  $K$  does not contain finite elements. Therefore,  $K$  is an infinite set.

### Exercise 1.3

Identify each of the following sets as finite set, infinite set or empty set

- A.  $D$  is the set of integers less than 0
- B.  $S$  is the set of natural numbers less than 1.
- C.  $Q$  is the set of grade 7 teachers who are 13 years old.
- D.  $X$  is the set whole numbers between -2 and 10.

## 1.2. THE RELATION AMONG SETS

### i. Subsets



#### Activity 1.1:

What relationships do you observe?

- A.  $M = \{-2, -1, 0, 1, 2\}$  and  $N = \{-2, -1, 0, 1, 2\}$
- B.  $P = \{1, 2, 3\}$  and  $Q = \{1, 2, 3, 4, 5\}$

#### Definition:

Given two Sets  $A$  and  $B$  if every element of set  $A$  is an element of the set  $B$ , then we say that  $A$  is a subset of  $B$ , and we denote this relation by  $A \subseteq B$ . If set  $A$  is not subset of  $B$ , we denote by  $A \not\subseteq B$ .

Note:

For any set,

A. Empty set is a subset of every set or  $\emptyset \subseteq A$

B. Every set is the subset of itself or  $A \subseteq A$

#### Example 1.11

- A. List all subsets of the set  $A = \{a, b, c, d\}$ . How many are there?
- B. List all subsets of the set  $L = \{1, 2, 3\}$ . How many are there?

#### Solution:

- A. The subsets of  $A$  are  $\{a\}$ ,  $\{b\}$ ,  $\{c\}$ ,  $\{d\}$ ,  $\{a, b\}$ ,  $\{a, c\}$ ,  $\{a, d\}$ ,  $\{b, c\}$ ,  $\{b, d\}$ ,  $\{c, d\}$ ,  $\{a, b, c\}$ ,  $\{a, b, d\}$ ,  $\{b, c, d\}$ ,  $\{a, b, c, d\}$  and  $\emptyset$ .

There are 16 subsets of the set  $A = \{a, b, c, d\}$ .

B. The subsets of L are  $\{1\}$ ,  $\{2\}$ ,  $\{3\}$ ,  $\{1, 2\}$ ,  $\{1, 3\}$ ,  $\{2, 3\}$ ,  $\{1, 2, 3\}$ , and  $\emptyset$ .

There are 8 subsets of the set  $N = \{1, 2, 3\}$ .

## ii. Proper subset

### Definition:

**Set A is a proper subset of set B if every element of set A is an element of the set B but there exists at least one element in B which is not an element of the set A. It is denoted by  $A \subset B$ . Which is read as 'A is a proper subset of B'. That is,  $A \subset B$  means  $A \subseteq B$  but  $B \not\subseteq A$ .**

### Example 1.12

Given set  $X = \{a, b, c\}$ , the sets  $\emptyset$ ,  $\{a\}$ ,  $\{b\}$ ,  $\{c\}$ ,  $\{a, b\}$ ,  $\{a, c\}$  and  $\{b, c\}$  are proper subsets of set X but  $\{a, b, c\}$  is not proper subsets of set X. i.e. is  $X \not\subset X$ .

**Fact:** If a set A is finite with n elements, then

**i. The number of subsets of A is  $2^n$ .**

**ii. The number of proper subsets of A is  $2^n - 1$ .**

## i. Power set

### Definition:

**Let A be any set, the power set of A, denoted by  $P(A)$ , is the set of all subsets of A. That is,  $P(A) = \{S/S \subseteq A\}$ .**

### Example 1.13

Let  $L = \{-1, 1\}$ , then the subsets of L are  $\emptyset$ ,  $\{-1\}$ ,  $\{1\}$ , and L. There

fore,  $P(A) = \{\emptyset, \{-1\}, \{1\}, L\}$ .

## ii Equality of Sets

### Definition:

Given two sets A and B, if every element of A is also an element of B and if every element of B is also an element of A, then the sets A and B are said to be equal. We write this as  $A = B$ .

$\therefore A = B$ , if and only if  $A \subseteq B$  and  $B \subseteq A$  i.e. they have the same number of elements and their elements are the same.

If A and B are not equal, we write  $A \neq B$ .

## iii. Equivalence of sets

### Definition:

Two sets A and B are said to be equivalent, written as  $A \leftrightarrow B$  or  $(A \sim B)$ , if there is a one to one correspondence between them.

Observe that

two finite sets A and B are equivalent, if and only if  $n(A) = n(B)$ .

### Example 1.14

1. Let  $A = \{1, 2, 3, 4\}$  and  $B = \{1, 4, 2, 3\}$ . Since two sets contain exactly the same/identical elements,  $A = B$ .
2. Let  $A = \{a, b, c, d\}$  and  $B = \{1, 2, 3, 4\}$  since two sets contain equal number of elements,  $A \leftrightarrow B$ .

## Exercise 1.4

1. State whether each of the following statements is true or false. If it is false justify your answer.

A. Every equal set are equivalent sets.

D.  $\emptyset \subseteq \{\{3\}\}$

B.  $\{a, b, c, d\} \subseteq \{b, a, d, c\}$

E.  $\{0,1\} \not\subseteq \{\{0,1\}, 0,1\}$ .

C.  $\{3\} \subseteq \{\{3\}\}$

2. Let  $Q = \{0, 1, 2\}$  then find all

- i. Subsets of  $Q$
  - ii. Proper subsets of  $Q$
  - iii. Power sets of  $Q$
3. Let  $A = \{0, \{1, 2\}\}$  then find all (i-iii) in question 2 above.
4. Which of the following pairs represent equal sets and which of them represent equivalent sets?
- i.  $\{\emptyset\}$  and  $\{1\}$
  - ii.  $K = \{x \in \mathbb{N} / x \text{ is a factor of } 6\}$  and  $H = \{x \in \mathbb{N} / 0 < x < 7\}$
  - iii.  $A =$  The set of even natural numbers between 1 and 7  
 $B =$  The set of even divisors of 12 less than 12.

### 1.3. OPERATIONS ON SETS

As there are operations on the sets of numbers there are operations on the sets.


Here we consider only two operations among the operations of sets. These are the intersection of sets and the union of sets. We will see this one by one in detail as follows.

#### 1.3.1 THE INTERSECTION OF SETS

##### **Definition:**

*The intersection of two sets  $A$  and  $B$ , denoted by  $A \cap B$  and read as “ $A$  intersection  $B$ ” is the set of all elements common to both set  $A$  and set  $B$ . That is,  $A \cap B = \{x | x \in A \text{ and } x \in B\}$*

Using the Venn diagram, can be represented by the white region:



**Example: :**


- Let  $A = \{1, 3, 5, 6, 7, 8\}$  and  $B = \{2, 3, 6, 8, 9\}$ . Then  $A \cap B = \{3, 6, 8\}$ .
- Let  $S = \{a, b, c, d\}$  and  $T = \{f, b, d, g\}$ . Then  $S \cap T = \{b, d\}$
- Let  $W = \{4, 6, 8\}$  and  $X = \{1, 3, 5, 7\}$ . Here these two sets have no elements in common, so their intersection is empty set, that is  $W \cap X = \emptyset$

**Definition:**

*Two or more sets that have no common element are called disjoint sets.*

*Sets  $A$  and  $B$  are disjoint sets, if  $A \cap B = \emptyset$*

*In Venn diagram, the disjoint sets  $A$  and  $B$  can be represented as:*

**Properties of the intersection of sets**

For any sets  $A$ ,  $B$  and  $C$  and the universal set  $U$

Commutative property:  $A \cap B = B \cap A$

Associative property:  $(A \cap B) \cap C = A \cap (B \cap C)$


Identity property:  $A \cap U = A$

### 1.3.2 THE UNION OF SETS

**Definition:**

*The union of two sets  $A$  and  $B$ , denoted by  $A \cup B$  and read as “ $A$  union  $B$ ” is the set of all elements that are members of set  $A$  and set  $B$  or both of the sets. That is,  $A \cup B = \{x / x \in A \text{ or } x \in B\}$*

We can represent using Venn diagram as follows:



**Example:** Let  $A = \{a, b, c, d, e\}$

$B = \{c, d, e, f, g\}$ , then  $A \cup B = \{a, b, c, d, e, f, g\}$

**Example:**  $A = \{a, b\} \cup \{c, d, e\} = \{a, b, c, d, e\}$

$B = \{1, 2, 3, 4, 5\} \cup \emptyset = \{1, 2, 3, 4, 5\}$

#### Properties of the union of sets

For any sets  $A$ ,  $B$  and  $C$  the universal set  $U$

- i. Commutative property:  $A \cup B = B \cup A$
- ii. Associative property:  $(A \cup B) \cup C = A \cup (B \cup C)$
- iii. Identity property:  $A \cup \emptyset = A$

#### Exercise 1.5

1. Given  $A = \{a, b, \{c\}\}$ ,  $B = \{b, c\}$  and  $C = \{\{c\}, d\}$  find:

A.  $A \cap B$

C.  $B \cap C$

B.  $A \cap C$

D.  $(A \cap B) \cap C$

2. Given  $A = \{1, 2, \{3\}\}$ ,  $B = \{2, 3\}$ , and  $C = \{\{3\}, 4\}$  find

- A.  $A \cup B$                       B.  $B \cup C$                       C.  $A \cup C$   
 D.  $(A \cup B) \cup C$               E.  $A \cup (B \cup C)$

3. Determine whether each of the following statements is true or false:

- A. If  $x \in A$  and  $x \in B$ , then  $x \in (A \cap B)$   
 B. If  $x \notin A$  and  $x \in B$ , then  $x \in (A \cap B)$ .  
 C. If  $x \notin A$  and  $x \notin B$ , then  $x \notin (A \cup B)$ .  
 D. For any set,  $A$ ,  $A \cup \emptyset = A$ .  
 E. If  $A \subseteq B$ , then  $A \cup B = B$   
 F. If  $x \in (A \cap B)$ , then  $x \in A$  and  $x \in B$   
 G. If  $A \subseteq B$  then  $A \cap B = A$   
 H. For any set,  $A$ , then  $A \cap A = A$   
 I. If  $A \cup B = \emptyset$ , then  $A = \emptyset$  and  $B = \emptyset$

4. Let  $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$  be the universal set and


let  $A = \{0, 2, 3, 5, 7\}$                        $B = \{0, 2, 4, 6, 8\}$  and

$C = \{x / x \text{ is a factor of } 6\}$  Find

- i.  $A \cap C$                       ii.  $B \cap C$                       iii.  $A \cap (B \cap C)$

5. Use the Venn diagram given below to find each of the following set relations.

- A.  $A \cap (B \cap C)$   
 B.  $A \cup B$   
 C.  $A \cap B$   
 D.  $A \cup C$   
 E.  $A \cup (B \cap C)$   
 F.  $B \cap C$



## Summary on Unit 1

- ✓ A set is a well-defined collection of objects and the objects of a set are called *elements* of the set. Elements of a set are also called members of the set.
- ✓ Sets can be described by *listing method* or *verbal method*.
- ✓ A set that has no element is called an *empty set* or *vacuous set* and denoted by  $\emptyset$  or  $\{ \}$ .
- ✓ A set A is called *finite* if and only if it is the empty set or has exactly *n* elements, where *n* is a natural number. Set that is not finite is called an *infinite set*.
- ✓ Two sets A and B are said to be *equal* if every element of one is an element of the other.
- ✓ A set A is a subset of B if every element of set A is an element of set B.
- ✓ For any two sets A and B,
  - A. The union of A and B, denoted by  $A \cup B$ , is the set of all elements that are members of either A or B.
  - B. The intersection of A and B, denoted by  $A \cap B$ , is the set of all elements that are members both A and B.

## REVIEW EXERCISES ON UNIT 1

1. Determine whether each of the following is set or not.
  - A. The collection of all tall students in your class.
  - B. The collection of all whole numbers divisible by 2.
  - C. The collection of model farmers in your locality.
  - D. The collection of odd numbers.
  - E. The collection of all intelligent students in south region.

2. Which of the following represent equal sets?
  - A.  $A = \{a, b, c, d, e\}$
  - B.  $B = \{a, e, i, o, u\}$
  - C.  $F =$  The first five letters in the English alphabet.
  - D.  $V =$  The vowel letters in the English alphabet.
3. Rewrite each of the following statements using correct notation of elements and sets.
  - A. A is a set whose elements are 1,2,3,4 and 5.
  - B. 6 is not an element of set A.
  - C. W is the set of all natural numbers between 3 and 21
4. Which of the following pairs of sets are equivalent?
  - A.  $\{1,2,3,4,5\}$  and  $\{m,n,o,p,q\}$
  - B.  $\{1,2,3,4,5\}$  and  $\{m,n,o,p,q,r\}$
  - C.  $\{x | x \text{ is a factor of } 6\}$  and The set of natural number less than 9
5. If  $X = \{a,b,c,d,e,f,g,h,i,j\}$ ,  $Y = \{b,d,f,h\}$  and  $Z = \{a,b,e,f,g,h\}$ , then find each of the following sets:
  - A.  $X \cap Y$
  - B.  $Y \cap X$
  - C.  $Z \cup Y$
  - D.  $Y \cup Z$
  - E.  $(X \cup Y) \cup Z$
  - F.  $(X \cap Y) \cap Z$
6. Identify each of the following as finite or infinite.
  - A=All integers less than 20.
  - B=All students in your class who are older than 20 years of age.
  - C=All natural numbers less than 1.

# UNIT 2

# INTEGERS

## *Learning Outcomes:*

At the end of this unit, learners will be able to:

- ↪ represent integers on a number line
- ↪ identify the commutative, associative and distributive properties of operation of integers
- ↪ perform the operations addition and subtraction on integers.
- ↪ apply integers in the real-life situation.

## *Introduction*

In the previous grades you had already learnt about whole numbers and natural numbers. These numbers together with negative numbers form another set of numbers called **Integers**. In this unit you will learn about integers and the four fundamental operations on integers.

## **2.1. Revision of whole and natural numbers**

### *Learning Outcomes:*

At the end of this unit, learners will be able to:

- ↪ Describe whole and natural numbers
- ↪ Identify the relation between whole and natural numbers.

### Activity 2.1.1

1. List down the first three smallest natural numbers.
2. List down the first five smallest whole numbers.
3. What is the smallest and greatest natural number?
4. What is smallest and the greatest whole number?
5. Define:

A. Natural number

B. Whole number

### Definition 2.1.:

*Whole numbers are numbers contains natural number and zero.*

*Whole number is denoted by  $\mathbb{W}$  and is described by*


$$\mathbb{W} = \{0, 1, 2, 3, \dots\}$$


### Definition 2.2:

*Natural number is a set of numbers denoted by  $\mathbb{N}$  and is described by  $\mathbb{N} = \{1, 2, 3, \dots\}$*

*Natural numbers are also called **counting numbers**.*

Relationship between natural numbers and whole numbers is described below





Revision of operations on natural numbers and whole numbers

**Example 1.** Find the following sums.

A.  $452 + 323$

B.  $3234 + 598$

C.  $423 + 5672$

**Solution:**

$$\begin{array}{r} A. \ 452 \\ + 323 \\ \hline 775 \end{array}$$

$$\begin{array}{r} B. \ 3234 \\ + 598 \\ \hline 3832 \end{array}$$

$$\begin{array}{r} C. \ 423 \\ + 5672 \\ \hline 6195 \end{array}$$

**Example 2.** Find the following differences

A.  $453 - 223$

B.  $4567 - 352$

C.  $456 - 456$

**Solution:**

$$\begin{array}{r} A. \ 453 \\ - 223 \\ \hline 230 \end{array}$$

$$\begin{array}{r} B. \ 4567 \\ - 352 \\ \hline 4215 \end{array}$$

$$\begin{array}{r} C. \ 456 \\ - 456 \\ \hline 0 \end{array}$$

**Example 3.** Find the product of the following numbers.

A.  $46 \times 44 = 2024$

C.  $123 \times 342 = 42066$

B.  $72 \times 78 = 5616$

**Example 4.** Find the quotient of the following

A.  $455 \div 5 = 91$

B.  $4824 \div 12 = 402$

**Note:**

- \* The sum of any two natural numbers is always natural number.
- \* The sum of any two whole numbers is always whole number.
- \* The product of any two natural numbers is always natural number.
- \* The product of any two whole numbers is always whole number.

### Exercise 2.1.1

Calculate the following

A.  $345 + 892$

C.  $6543 - 234$

E  $5427 \div 9$

B.  $342 + 435$

D.  $23 + (654 - 345)$

F.  $342 \times 345$

## 2.2. INTRODUCTION TO INTEGER

### Learning Outcomes:

At the end of this unit, learners will be able to:

- Apply real-life applications of integers in terms of temperature, altitude and money to express positive and negative numbers
- Define the set of integers
- Indicate integers on the number line.
- Describe the relations, among natural numbers, whole numbers and integers  $\mathbb{N} \subset \mathbb{W} \subset \mathbb{Z}$



### Activity 2.2.1

Discuss with your friends/partners

Anders Celsius the Swedish astronomer who lived between 1701 and 1744 A.D. He devised a way of measuring temperature which was adjusted and improved after his death.



Figure 2.2.1

1. The table below shows that the temperature recorded on three consecutive days for Hawassa city on **Table 1**

Days	Temperature in degree Celsius
Monday	16 °c above zero
Tuesday	6°c below zero
Wednesday	1°c below zero



Monday



Tuesday



Wednesday

Figure 2.2.2 Thermometer

- A. What was the temperature on each day using the sign plus or minus?
  - B. How far did the temperature from zero in each day as shown in table and figures above?
  - C. What was the coldest temperature?
  - D. What was the hottest temperature?
2. What is the number that comes after zero?
  3. What is the number that comes before zero?

### Note:

\* Very cold temperatures below zero can be described by number with **minus** sign which is called **negative numbers**.

\* The temperatures above zero can be described by a number with **plus** sign which is called **positive numbers**.

### Example 1

The temperature  $5^{\circ}\text{C}$  below zero can be written as  $-5^{\circ}\text{C}$  and The temperature  $5^{\circ}\text{C}$  above zero can be written as  $+5^{\circ}\text{C}$  or  $5^{\circ}\text{C}$


**Note:**

- \* positive numbers can be written without plus sign. For instance,  $+2$  simply written as  $2$
- \* Do you understand the difference of the concepts of the numbers below zero and above zero?

### Definition 2.3:


An integer is a set of numbers consisting of whole numbers and negative numbers. The set of integers is denoted by

$$\mathbb{Z} = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$$



Negative integer is denoted by  $\mathbb{Z}^-$  and positive integer is denoted by  $\mathbb{Z}^+$ . The set of positive integers is the same as the set of natural numbers. (i.e.  $\mathbb{Z}^+ = \mathbb{N}$ ). Zero is an integer which is neither positive nor negative. Integers, whole numbers and natural numbers are related as follows: in Venn diagram  $\mathbb{N} \subset \mathbb{W} \subset \mathbb{Z}$

Note that decimals and fractions are not integers.



Integer numbers represented on Venn diagram

The number  $-4$  is read as negative four and  $4$  is read as positive four or four

**Example 2:**

- A. List some Positive integers
- B. List five negative integers
- C. Give example of integer which is neither positive nor negative
- D. List four examples of numbers which are not integers

**Solution**

A. 9, 54, 380, 6284, 89123 are Positive integers.

B. -9, -54, -125, -647, -32564 are negative integers

C. 0




D. 2.5,  $\frac{3}{4}$ , 1.08,  $\frac{10}{7}$  are not integers

**Example 3:**

**Represent the following numbers on number line**

- A. From -3 to 3      B. From -4 to 4      C. From -6 to 6


**solution:** First draw horizontal line and mark the numbers on the number line



**Integer numbers represent on a number line**

## Exercise 2.2.1

1. Give the numbers represented by the letters on the number line below



2. Represent the numbers 1, -1, 5, -5, 2 and -2 on a number line
3. Write down four numbers which are not integer.
4. Write the following integers in word  
 A. -3                      B. . 21                      C. -132                      D. 99
5. write the following numbers as numeral  
 A. Positive four                      C. Negative fifty-three  
 B. Negative twenty                      D. one hundred one
6. Represent the following fact by using a numeral + and - signs.  
 A. A loss of Birr 100                      D. Five minutes late  
 B. A rise of temperature                      E. below zero  
 C. A walk of 7 km forward                      F. above zero
7. What is the relationship between natural number and positive integer?
8. Explain the relationship between whole number, natural number and integer
9. From the numbers listed below select integers  
 A.  $2\frac{1}{3}$                       C.  $\frac{2}{5}$                       E. 12                      G. 0.25  
 B. 0                      D. 1.02                      F. -294                      H. 455

## Opposite numbers

### Definition 2.4

*Opposite numbers are numbers that are at the same distance from zero in opposite direction and the opposite of zero is zero.*



Using the above number line, the opposite of some integers is listed in the table below.

Number	-10	-9	-8	-7	-6	1	2	3	4	5
Opposite number	10	9	8	7	6	-1	-2	-3	-4	-5

**Example 4:** find the opposite of the following numbers

A. 12

B. -9

C. 413

D. -564

**Solution**

A. The opposite of 12 is -12

C. The opposite of 413 is -413

B. The opposite of 0 is 0

D. The opposite of -564 is 564

**Note:**

i. The opposite of  $a$  is  $-a$

ii. For integer  $a$ ,  $-(-a) = a$

iii. The opposite of positive number is negative number

iv. The opposite of negative number is positive number

v. The opposite of zero is zero.

### Exercise 2.2.2

1. Find the opposite of each integer given below.

A. 70

B. -23

C. -170

D. 0

2. Find the opposite of the following

A. Gain of Birr 120

B.  $15^{\circ}$  above zero

C. A walk of 5 km backward

3. Ahmed saw on weather report that Hawassa's temperature was  $9^{\circ}\text{C}$  above zero.

- A. write Hawassa's temperature as an integer.
- B. If the temperature in Bale is opposite of Hawassa's temperature.  
What is the temperature in Bale?

## 2.3. COMPARING AND ORDERING INTEGERS

### Learning Outcomes:

At the end of this unit, learners will be able to:

- ↪ Compare and order integers using a number line
- ↪ Determine the predecessor and successor of a given integers



### Activity 2.3.1

*Discuss with your friends/partners*

1. On October, the lowest temperature of three different cities was recorded as follows

City	Temperature in
Debire Birhan	4° below zero
Addis Ababa	4° above zero
Adama	10° above zero

- A. Which city has the lowest (minimum) temperature?
  - B. Which city has the highest temperature?
  - C. Indicate the above temperatures on number line and arrange the temperatures from the smallest to largest
2. Indicate each of the following pairs of numbers on number line and circle the number which is found to the right of the other.
 

A 0 and 6      B. 0 and -6    C. -6 and -4    D. 6 and 4
  3. Circle the bigger number from the following pairs of numbers.
 

A. 0 and 6      B. 0 and -6    C. -6 and -4    D. 6 and 4

To compare integers, draw number line and indicate the numbers on number line then,

♻ The number which is to the **right** of the other is **bigger** number.

♻ The number which is to the **left** of the other is **smaller** number.

We use the symbols to compare numbers.

♻  $a > b$  means  $a$  is greater than  $b$

♻  $a < b$  means  $a$  is less than  $b$

♻  $a = b$  means  $a$  is equal to  $b$

Consider the following number line



♻ A is to the left of B, hence A is smaller than B, (i.e.  $A < B$ )

♻ B is to the right of A, hence B is bigger than A. (i.e.  $B > A$ )

### Example 1.

For each pair of numbers below select the number which is to the right of the other

A. 3 and 6


C. -3 and -6

B. 10 and 5

D. -10 and -5

### Solution:

*First draw a number line and mark the numbers*



**A. 6 is to the right of 3      C. -3 is to the right of -6**


**B. 10 is to the right of 5      D. -5 is to the right of -10**

**Example 2.** Insert ">" or "<" to express the corresponding relationship between the following pairs of integers

- A.  $-10$  \_\_\_\_  $0$     B.  $0$  \_\_\_\_  $-3$     C.  $0$  \_\_\_\_  $-6$     D.  $-8$  \_\_\_\_  $0$

**Solution:**

use a number line to compare numbers



A.  $-10$  is to the left of  $0$ , hence  $-10$  is smaller than  $0$ , symbolically,  $-10 < 0$

B.  $0$  is to the right of  $-3$ , hence  $0$  is greater than  $-3$ , symbolically  $0 > -3$

C.  $0$  is to the right of  $-6$ , hence  $0$  is greater than  $-6$ , symbolically  $0 > -6$

D.  $-8$  is to the left of  $0$ , hence  $-8$  is smaller than  $0$ , symbolically  $-8 < 0$

**Note:**



All negative integers are to the left of  $0$ , hence every negative integer is less than  $0$ .




$0$  is to the right of any negative integer, hence  $0$  is greater than any negative integer.

**Example 3:**

**Insert** ">" or "<" to express the corresponding relationship between the following pairs of integers

- A.  $-4$  \_\_\_\_  $2$     B.  $5$  \_\_\_\_  $-8$     C.  $8$  \_\_\_\_  $-6$     D.  $-10$  \_\_\_\_  $4$

**Solution:** use a number line to compare numbers





A.  $-4$  is to the left of  $2$ , hence  $-4$  is smaller than  $2$ , symbolically  $(-4 < 2)$

B.  $5$  is to the right of  $-8$ , hence  $5$  is greater than  $-8$ , symbolically  $(5 > -8)$

C.  $8$  is to the right of  $-6$ , hence  $8$  is greater than  $-6$ , symbolically  $(8 > -6)$

D.  $-10$  is to the left of  $4$ , hence  $-10$  is smaller than  $4$ , symbolically  $(-10 < 4)$


**Note:**

-  All negative integers are to the left of positive integers; hence every negative integer is less than any positive integer.
-  All positive integers are to the right of any negative integer; hence every positive integer is greater than any negative integer.


**Example 4.** Compare the following integers using the sign " $>$ " " $=$ " or " $<$ ".

A  $-94$  \_\_\_\_  $-100$       B.  $-250$  \_\_\_\_  $-180$       C  $-40$  \_\_\_\_  $-40$

**Solution:** To compare the numbers draw part of number line



**Solution:** draw part of number line




**Solution**

C  $-40$  \_\_\_\_  $-40$

*This two numbers are the same numbers; hence they are equal*

**Example 5.** Name all integers which lie between  $-5$  and  $2$


**Solution:** Draw a number line



*The integers that lie between  $-5$  and  $-2$  are  $-4, -3, -2, -1, 0$  and  $1$*

**Example 6.** List all integers which lie between  $-4$  and  $-3$

**Solution:** Draw a number line



There is no integer between -4 and -3

### Exercise 2.3.1

1. Insert ">" "=" or "<" to express the corresponding relationship between the following pairs of integers.

A. -150 ____ 0	E. 65 ____ -20	I. -120 ____ -80
B. 0 ____ -300	F. -340 ____ -341	J. -360 ____ 185
C. -1200 ____ -74	G. -23 ____ -23	K. 0 ____ 13
D. -34 ____ 34	H. -8 ____ 2	L. 14 ____ 0

2. From each pair of numbers below circle the number which is to the right of the other.

A. 3 and 6	C. 10 and 5	E. -87 and -99	G. 35 and -45
B. -3 and -6	D. -1005 and -105	F. -360 and -1	H. 0 and -12

3. List all integers which lie between the following pairs of numbers

A. -2 and 3	B. -1 and 1	C. -98 and -95	D. -10 and -9
-------------	-------------	----------------	---------------

4. The five integers  $x$ ,  $y$ ,  $z$ ,  $n$ , and  $m$  are represented on the number line below



Using "<" or ">" fill in the blank space

A. $z$ ____ $x$	B. $m$ ____ $x$	C. $z$ ____ $n$	D. $0$ ____ $z$
-----------------	-----------------	-----------------	-----------------

5. Answer the following questions

- What is the greatest negative integer?
- What is the smallest positive integer?
- What is the greatest integer?
- What is the smallest integer?

## Successor and predecessor of integers

**Successor** of an integer is an integer that *comes after* the given integer.

**Predecessor** of an integer is an integer that *comes before* the given integer.

**Example 7:** Find the successor of the given integers

- A. 5      B. 0      C. -4      D. -99      E. 99

**Solution:**

A. The successor of 5 is 6.

D. The successor of -99 is -98.

B. The successor of 0 is 1.

E. -The successor of 99 is 100.

C. The successor of -4 is -3.

**Example 8:** Find the predecessor of the given integers.

- A. 5      B. 0      C. -4      D. -99      E. 99

**Solution:**

A. The successor of 5 is 4.

D. The successor of -99 is -100.

B. The successor of 0 is -1.

E. -The successor of 99 is 98.

C. The successor of -4 is -5.

## Ascending and Descending order of integers

**Ascending order** means arranging the given numbers in increasing order. (i.e. arranging from the smallest to the largest number).

**Descending order** means arranging the given numbers in decreasing order. (i.e. arranging from the largest to the smallest).

**Example 9:** Arrange the following set of numbers in ascending order


A. 10, 2, 5, 0, 8, 120

C. -256, 31, 0, -25, 13, -52, 103


B. -40, -8, 0, -12, -10

**Solution:**


**A.** 0, 2, 5, 8, 10, 120



**B.** -40, -12, -10, -8, 0,



**C.** -256, -52, -25, 0, 13



**Example 10**


Arrange the following set of numbers in descending order

A. 0, -65, -100, -12, -1


B. 55, -75, 0, 15, -35

**Solution**

**A.** 0, -1, -12, -65, -100



**B.** 55, 15, 0, -35, -75



## Exercise 2.3.2

1. Find the successor of the following integers

A. 20

B. -16

C. 999

D. -999

2. Find the predecessor of the following integers

A. 1000

B. -17

C. -999

D. -1000

3. Arrange the following numbers in ascending order

A. 0, -78, 17, -24, 71,

B. -230, -157, -400, 350, 50

4. Arrange the following numbers in descending order

A. -9, -99, 69, 59, -89,      B 200, -300, -757, -445, 400

5. Abebe, Almaz, Hailu and Derartu played basketballs. The results are shown in the table below

	First play	Second play	Final result
Abebe	Loss 5 basket balls	Won 7 basket balls	Won 2 baskets (+2)
Almaz	Won 5 basket balls	Loss 5 basket balls	
Hailu	Loss 3 basket balls	Loss 2 basket balls	
Derartu	Won 1 basket balls	Won 2 basket balls	

- Complete the final result of the players in the above table
- Who was the winner?
- Who was the loser?
- Arrange the final result of the players in descending order.

### Project work:

Register the daily temperature of Addis Ababa, Adama, Hawassa, Debire Birhan, Gambela, Semera, and Bahir Dar from Ethiopian mass media in a given day and answer the following:

- Graph the temperature of each city on number line
- Which city was coldest?
- Which city was warmest?

## 2.4. ADDITION AND SUBTRACTION OF INTEGERS

### Learning Outcomes:

At the end of this unit, learners will be able to:

- Find the sum of integers.
- Find the difference between two integer

## Addition of integers



### Activity: 2.4.1

1. Hussen, Almaz, Hailu and Derartu are selling different articles. Their profit or loss is shown in the table below

	First sell	Second sell	Total profit/ loss
Hussen	Loss Birr 15	Profit Birr 17	
Almaz	Profit Birr 15	Loss Birr 15	
Hailu	Loss Birr 13	Loss Birr 20	
Derartu	Profit Birr 10	Profit Birr 12	

2. Find the sum of the following using number line

- A.  $4+(-8)$       B.  $-2+5$       C.  $-3+(-4)$       D.  $-4+0$

3. Find the sum of the following integers

- A.  $3+2$       B.  $-20+0$       C.  $-3+2$       D.  $4+(-3)$       E.  $-10+10$

### Rules: for addition of integers on number line


1. Start the arrow from the first addend.
2. Move the arrow to the **right** with the same magnitude as the second addend, if the sign of the second addend is **positive**.
3. Move the arrow to the **left** with the same magnitude as the second addend, if the sign of the second addend is **negative**.
4. The sum of the integers is at the point where the arrow ends.

**Example 1:** find the sum of the following integers using number line


- A.  $3+(-6)$       B.  $-3+(-6)$       C.  $-3+6$

### Solution


A.  $3+(-6) = -3$



B.  $-3+(-6) = -9$



C.  $-3 + 6 = 3$



**Example 2** add the following numbers

A.  $-4 + 4$     B.  $6 + (-6)$     C.  $-1 + 1$     D.  $2 + (-2)$

**Solution:**

A.  $-4 + 4 = 0$     B.  $6 + (-6) = 0$     C.  $-1 + 1 = 0$     D.  $2 + (-2) = 0$

**Note:** the sum of any two **opposite** integers is **zero**

### Addition of positive and negative integers without using number line

To find the sum of positive and negative integers, follow the following steps

**Step 1:** Take the sign of a number with greater magnitude.

**Step 2:** Find the **difference** of magnitude of the numbers.

**Step 3:** put the sign in front of the difference

**Example 3:** Find the sum of the following integers

A.  $22 + (-43)$

**Solution:**

$43 > 22$ , the sign of 43 is  $(-)$ , hence the sum is  $(-)$

$$22 + (-43) = -(43-22)$$

$$= -21$$

B.  $625 + (-214)$

**Solution:**

$$\begin{aligned} \underline{\hspace{2cm}} \quad 625 > 214, \text{ the sign of } 625 \text{ is } (+), \text{ hence the sum is } (+) \\ 625 + (-214) &= + (625 - 214) \\ &= 411 \end{aligned}$$

C.  $-1300 + 560$

**Solution:**

$$\begin{aligned} 1300 > 560, \text{ the sign of } 1300 \text{ is } (-), \text{ hence the sum is } (-) \\ -1300 + 560 &= - (1300 - 560) \\ &= - 740 \end{aligned}$$

### Addition of two negative numbers

To find the sum of two negative integers: -

**Step 1:** The sign of the sum is always **negative**.

**Step 2:** Find the **sum** of magnitude of the numbers.

**Step 3:** Put the **negative** sign in front of the sum

**Example 4:** Find the sum of the following

A.  $-15 + (-35)$       B.  $-36 + (-40)$       C.  $-120 + (-76)$

**Solution:**

$$\begin{aligned} \text{A. } -15 + (-35) &= - (15 + 35) = - 50 \\ \text{B. } -36 + (-40) &= - (36 + 40) = - 76 \\ \text{C. } -120 + (-76) &= - (120 + 76) = - 196 \end{aligned}$$

**Example 5:** A person saved Birr 200 and then he spent Birr 150. How much money does he have now?

**Solution:**

*Saved 200 can be represented by + 200*

*Spent 150 can be represented by -150*

*We get:*

***Therefore, he has left Birr 50.***

**Example 6:** A submarine at 21m below sea level suddenly moves up about 6m. At what depth is the submarine located now?

**Solution:**

*21m below sea level can be represented by - 21*

*Moves up about 6m can be represented by +6, then*

*We get  $-21 + (+6) = -(21 - 6) = -15$*

***The submarine is 15m below sea level.***

## Property of addition of Integers

### *I. The sum of two integers is an integer*

**Example**

A.  $2+5= 7$  ,7 is integer

C.  $-4+(-5)=-9$  ,-9 is integer

B.  $4 + (-6) = -2$  , -2 is an integer

D.  $-3+5=2$  ,2 is integer

### *II. Addition has commutative property:*

for any two integers m and n

$$m + n = n + m$$

**Example**

$$\begin{array}{rcl} 5 + (-3) & & -3 + 5 \\ = + (5 - 3) & = & + (5 - 3) \\ = 2 & = & 2 \end{array}$$

Therefore,  $5 + (-3) = -3 + 5$

### III. Associative property:

for any three integers  $x, y$  and  $z$

$$(x + y) + z = x + (y + z)$$

#### Example

$$3 + (4 + (-5)) = (3 + 4) + (-5)$$

$$3 + (-1) = 7 + (-5)$$

$$2 = 2$$

Therefore,  $3 + (4 + (-5)) = (3 + 4) + (-5)$  is **Associative property**

### IV. Property of zero For any integer number

For any integer number  $x$ ,  $x + 0 = x = 0 + x$

### V. Property of opposites:

for any integer  $x$ ,  $x + (-x) = 0$

### Exercise 2.4.1

- Find the sum of the following integers?
 

A. $250 + (-250)$	C. $160 + (-160)$
B. $-170 + 170$	D. $-15 + 15$
- Find the sum of the following using number line.
 

A. $-3 + 7$	C. $6 + (-3)$	E. $-5 + (-2)$
B. $-4 + (-5)$	D. $-10 + 10$	F. $-7 + (4)$
- Find the sum of the following integers
 

A. $23 + (-10)$	D. $230 + (-63)$	G. $12 + (-23)$
B. $-30 + 40$	E. $-325 + 234$	H. $-2250 + (-457)$
C. $-560 + (-50)$	F. $-50 + (-45) + 62$	I. $-20 + (-10) + (210)$
- You have Birr 5 in debt. You borrow Birr 12 more. What is the total amount of your debt?
- An airplane takes off and then climbs 2500 meter. After 20 minutes, the airplane descends 150 meters. What is the airplane's current height?

## Subtraction of Integers



### Activity: 2.4.2

- Write the following differences in the form of sum.
 

A. $3 - 6 = 3 + (-6)$	C. $4 - 2$
B. $-3 - 5 =$	D. $-5 - (-6)$
- Find the following differences by expressing in the form of sum
 

A. $5 - 6 = 5 + (-6) = -1$	C. $4 - 2$
B. $-2 - 2 =$	D. $-7 - (-6)$

## Subtraction of integer

**Subtracting** an integer  $b$  from  $a$  is the same as adding opposite of  $b$  to  $a$  i.e. for any integers and ,

$$a - b = a + (-b)$$

**Example 1:** Find the difference of the following integer by expressing in the form of sum.

- |            |            |            |            |
|------------|------------|------------|------------|
| A. $5 - 2$ | B. $4 - 7$ | C. $2 - 6$ | D. $6 - 2$ |
|------------|------------|------------|------------|

**Solution:**

A.  $5 - 2$ , subtracting 2 means adding the opposite of 2

$$5 - 2 = 5 + (-2) = 3$$

B.  $4 - 7$ , subtracting 7 means adding the opposite of 7

$$\begin{aligned} 4 - 7 &= 4 + (-7) \\ &= -(7-4) = -3 \end{aligned}$$

C.  $2 - 6$ , subtracting 6 means adding the opposite of 6

$$\begin{aligned} 2 - 6 &= 2 + (-6) \\ &= -(6-2) = -4 \end{aligned}$$

D.  $6 - 2$ , subtracting 2 means adding the opposite of 2

$$\begin{aligned} 6 - 2 &= 6 + (-2) \\ &= +(6 - 2) = 4 \end{aligned}$$

### Example 2:

Find the difference of the following integer by expressing in the form of sum.

A.  $25 - (-30)$    B.  $-12 - (-18)$    C.  $-55 - (-20)$    D.  $8 - (-9)$

### Solution:

A.  $25 - (-30)$ , subtracting -30 means adding the opposite of -30

$$25 - (-30) = 25 + 30 = 55$$

B.  $-12 - (-18)$ , subtracting -18 means adding the opposite of -18

$$\begin{aligned} -12 - (-18) &= -12 + 18 \\ &= +(18 - 12) = 6 \end{aligned}$$

C.  $-55 - (-20)$ , subtracting -20 means adding the opposite of -20

$$\begin{aligned} -55 - (-20) &= -55 + 20 \\ &= -(55 - 20) = -35 \end{aligned}$$

D.  $8 - (-9)$ , subtracting -9 means adding the opposite of -9

$$8 - (-9) = 8 + 9 = 17$$

### Note:

*for any two-integer a and b*

$$a - (-b) = a + b$$

**Example3:** Find the difference of the following integer by expressing in the form of sum.

A.  $-21 - 12$

C.  $-24 - 35$

B.  $-38 - 44$

D.  $-248 - 125$

**Solution:**

A.  $-21 - 12$ , subtracting 12 means adding the opposite of 12

$$\begin{aligned} -21 - 12 &= -21 + (-12) \\ &= -(21 + 12) = -33 \end{aligned}$$

B.  $-38 - 44$ , subtracting 44 means adding the opposite of 44

$$\begin{aligned} -38 - 44 &= -38 + (-44) \\ &= -(38 + 44) = -82 \end{aligned}$$

C.  $-24 - 35$ , subtracting 35 means adding the opposite of 35

$$\begin{aligned} -24 - 35 &= -24 + (-35) \\ &= -(24 + 35) = -59 \end{aligned}$$

D.  $-248 - 125$ , subtracting 125 means adding the opposite of 125

$$\begin{aligned} -248 - 125 &= -248 + (-125) \\ &= -(248 + 125) = -373 \end{aligned}$$

**Note:**

**For any two-integer  $a$  and  $b$**

$$-a - b = -(a + b)$$

**Exercise 2.4.2**

1 express the following differences in the form of sum

A.  $3 - (-10)$

C.  $9 - 12$

E.  $-178 - 57$

B.  $-5 - (-6)$

D.  $25 - 17$

F.  $-365 - 1000$

2. Find the following differences.

A.  $0 - 3$

C.  $-75 - 15$

E.  $465 - (-778)$

B.  $-7 - (-8)$

D.  $-1789 - (-1000)$

F.  $25 - 75$

3. The melting point dry ice is  $-109^{\circ}\text{F}$ . The boiling point of dry ice is  $109^{\circ}\text{F}$  then, how many degrees is the boiling point above the melting point.

## 2.5. MULTIPLICATION AND DIVISION OF INTEGER NUMBERS

### Learning Outcomes:

At the end of this unit, learners will be able to:

- ↪ Solve problems on multiplication of integers.
- ↪ Identify the commutative and associative property of multiplication.
- ↪ Identify the distributive property of multiplication over addition.
- ↪ Divide integers whose quotient as expressed in decimals (2 decimal places)

### Multiplication of Integer numbers



#### Activity 2.5.1

1. Express each of the following product as sum and find the product

- A.  $5 \times 2$       B.  $9 \times 3$       C.  $(-5) \times 4$       D.  $(-6) \times 2$

2. Express each sum as product

- A.  $0 + 0 + 0 + 0$       C.  $(-2) + (-2) + (-2)$   
 B.  $3 + 3 + 3 + 3$       D.  $(-10) + (-10) + (-10) + (-10) + (-10) + (-10)$

3. Multiply the following

- A.  $2 \times 4$       C.  $3 \times 51$       E.  $4 \times (-25)$   
 B.  $0 \times 121$       D.  $-10 \times 2$

### Note:

*The product of two positive integers is **positive integer***

**Example 1:** find the product of the following

- A.  $3 \times 5$       B.  $11 \times 6$

### Solution:

using similar ways as product of natural number and whole number

$$A. 3 \times 5 = 5 + 5 + 5 = 15$$

$$B. 11 \times 6 = 66$$

**Example 2:** Find the product of the following numbers expressing in the form of **Sum**

$$A. 4 \times (-5) \quad B. 3 \times -13 \quad C. 6 \times -1 \quad D. 2 \times -365$$

**Solution:**

$$A. 4 \times (-5) = -5 + (-5) + (-5) + (-5) = -20$$

$$B. 3 \times -13 = -13 + (-13) + (-13) = -39$$

$$C. 6 \times -1 = -1 + (-1) + (-1) + (-1) + (-1) + (-1) = -6$$

$$D. 2 \times -365 = -365 + (-365) = -730$$

**Note:**

*for above examples, you can see that the product of positive and negative integer is negative integer.*

**Example 2:** Multiply the following

$$A. 4 \times (-3) \quad B. 12 \times (-11) \quad C. -10 \times 267$$

**Solution:**

$$A. 4 \times (-3) \Rightarrow \text{the result is negative sign (-)}$$

$$\Rightarrow 4 \times 3 = 12$$

$$\text{so } 4 \times (-3) = -12$$

$$B. 12 \times (-11) \Rightarrow \text{The result is negative sign (-)}$$

$$\Rightarrow 12 \times 11 = 132$$

$$\text{so } 12 \times (-11) = -132$$

$$C. -10 \times 267 \Rightarrow \text{The result is negative sign (-)}$$

$$\Rightarrow 10 \times 267 = 2670$$

$$-(10 \times 267) = -2670$$

**Example 4:** find the product of the following

A.  $-2 \times 2 = -4$     B.  $-2 \times 1 = -2$     C.  $-2 \times 0 = 0$     D.  $-2 \times (-1) = ?$

In the above example, the product pattern is  $-4, -2, 0$ , hence  $-2 \times (-1) = 2$

**Note:**

*The product of two negative integers is positive integer*

**Example 5:** Multiply

A.  $-3 \times (-5)$     B.  $-21 \times (-1)$     C.  $-6 \times (-13)$     D.  $-11 \times (-8)$

**Solution: -**

A.  $-3 \times (-5)$   $\Rightarrow$  The result is positive sign (+)

$\Rightarrow 3 \times 5 = 15$

so,  $-3 \times (-5) = 15$

B.  $-21 \times (-1)$   $\Rightarrow$  The result is positive sign (+)

$\Rightarrow 21 \times 1 = 21$

so,  $-21 \times (-1) = 21$

C.  $-6 \times (-13)$   $\Rightarrow$  The result is positive sign (+)

$\Rightarrow 6 \times 13 = 78$

so,  $-6 \times (-13) = 78$

D.  $-11 \times (-8)$   $\Rightarrow$  The result is positive sign (+)

$\Rightarrow 11 \times 8 = 88$

so,  $-11 \times (-8) = 88$

**Summary of product of integers**

Two factors	Sign of Product	Example
Both are positives	Positive	$3 \times 5 = 15$
Opposite sign	Negative	$-3 \times 5 = -15$ or $5 \times (-3)$
Both are negatives	Positive	$-3 \times (-5) = 15$

## Exercise 2.5.1

1. Multiply

A.  $6 \times 12$

C.  $-7 \times 13$

E.  $-17 \times (-103)$

B.  $13 \times 15$

D.  $16 \times (-25)$

F.  $-170 \times (-345)$

### Properties of multiplication of integer numbers



#### Activity 2.5.2

Which of the following statements are true and which are false

A.  $2 \times 4 = 4 \times 2$

C.  $5 \times (3 \times 2) = (5 \times 3) \times 2$

B.  $21 \times (-3) = -3 \times 21$

D.  $-5 \times (3 + 7) = (-5 \times 3) + (-5 \times 2)$

#### 1. Commutative properties of multiplication

For any two integers  $a$  and  $b$ ,

$$a \times b = b \times a \dots \text{Commutative property of multiplication}$$

##### Example 1

$$-2 \times 5 = 5 \times (-2)$$

$$-10 = -10$$

$$-12 \times (-20) = -20 \times (-12)$$

$$240 = 240$$

#### 2. Associative properties of multiplication

For any three integers numbers  $a$ ,  $b$ ,  $c$ , then

$$a \times (b \times c) = (a \times b) \times c \dots \text{Associative property}$$

##### Example 2

$$-2 \times (5 \times 6) = (-2 \times 5) \times 6$$

$$-2 \times 30 = -10 \times 6 \quad (\text{First multiply the bracket})$$

$$-60 = -60$$

#### 3. Distributive properties of multiplication over addition

Let  $a$ ,  $b$ , &  $c$  be any integers, then

$$a \times (b + c) = (a \times b) + (a \times c) \dots \text{Distributive property}$$

**Example 3**  $-6 \times (5 + 3) = (-6 \times 5) + (-6 \times 3)$

$$\begin{aligned} -6 \times (8) &= (-30) + (-18) \text{ (Apply operations in the bracket)} \\ -48 &= -48 \end{aligned}$$

#### 4. Properties of zero and 1 on multiplication

##### Note:

The product of any integer and zero is zero. Let ' $a$ ' be any integer, then or  $a \times 0 = 0$  or  $0 \times a = 0$

**Example 4** Find the product

A.  $2 \times 0$       B.  $0 \times 49$       C.  $-122 \times 0$       D.  $0 \times (-2999)$

##### Solution

$$A. 2 \times 0 = 0$$

$$C. -122 \times 0 = 0$$

$$B. 0 \times 49 = 0$$

$$D. 0 \times (-2999) = 0$$

##### Note:

The product of any integer and 1 is the number itself.

Let ' $a$ ' be any integer then or  $a \times 1 = a$  or  $1 \times a = a$

**Example 5**

A.  $9 \times 1 = 9$       B.  $-25 \times 1 = -25$       C.  $1 \times (-291) = -291$

#### Exercise 2.5.2

1. State the properties for the following

A.  $2 \times (-3 + 54) = (2 \times (-3)) + (2 \times 54)$

D.  $5 \times 1 = 5$

B.  $-525 \times 955 = 955 \times (-525)$

E.  $0 \times (-8) = 0$

C.  $2 \times (-13 \times 5) = (2 \times (-13)) \times 5$

2. Simplify each of the following pairs and compare their result

A.  $3 \times (-2 \times 3)$  and  $(3 \times (-2)) \times 3$

B.  $259 \times 0$  and  $0 \times 259$

C.  $-6 \times (5 + 2)$  and  $(-6 \times 5) + (-6 \times 2)$

D.  $-5 \times (-3 + (-8))$  and  $(-5 \times (-3)) + (-5 \times (-8))$

3. Copy and complete the following table below

a	b	c	$a \times b$	$b \times a$	$a \times (b \times c)$	$(a \times b) \times c$
-7	4	-3				
9	-2	5				
-6	-8	-12				

### Multiplication of three or more integers



#### Activity 2.5.3.

1. Multiply the following numbers and notice the sign of the product.

A.  $-4 \times 5 \times 3 \times 1$

C.  $-4 \times (-5) \times (-3) \times 1$

B.  $-4 \times 5 \times (-3) \times 1$

D.  $-4 \times (-5) \times (-3) \times (-1)$

2. Multiply the following numbers

A.  $-4 \times 5 \times 0 \times 1$

B Activity 1.5.3

What do you conclude from a and b?

The following properties are helpful in simplifying products with three or more factors:-



The product of an even number of negative factors is positive.



The product of an odd number of negative factors is negative.



Product of integers with at least one factor 0 is zero.



If you multiply an integer a by -1, then you get the opposite of a (i.e.  $-1 \times a = -a$ )



When you multiply a number by a variable, you can omit the multiplication sign and keep the number in front of the variable (i.e.  $3 \times y = 3y$ )

**Example 1:** Determine the sign of product

A.  $-1 \times 5 \times (-6)$

C.  $-5 \times (-2) \times 3 \times (-8) \times (-4) \times 7$

B.  $-265 \times (-81) \times 35 \times (-68)$

**Solution:**

A. The number of negative factors is 2 (even), then the product is positive

B. The number of negative factor is 3 (odd), then the product is Negative

C. The number of negative factor is 4 (even) then the product is positive.

**Example 2** Find the product of the following

A.  $-3 \times 5 \times (-4)$

C.  $-5 \times (-2) \times 3 \times (-8) \times (-4) \times 0$

B.  $-5 \times (-5) \times 3 \times (-8)$

**Solution:**

A. The number of negative factor is 2 (even), then the product is positive  $(-3 \times 5) \times (-4) = -15 \times (-4) = 60$

B. The number of negative factor is 3 (odd), then the product is Negative

$$\begin{aligned} \Rightarrow -5 \times (-5) \times 3 \times (-8) &= (-5 \times (-5)) \times (3 \times (-8)) \\ &= 25 \times (-24) \\ &= -600 \end{aligned}$$

C. There is one factor 0 in the product, hence the product is 0

$$-5 \times (-2) \times 3 \times (-8) \times (-4) \times 0 = 0$$

### Exercise 2.5.3

1. Find the sign of the product

A.  $12 \times (-50) \times (-61)$

C.  $(-1) \times (-2) \times (-3) \times (-4) \times (-5)$

B.  $(-125) \times (-3) \times (-52)$

D.  $(3) \times (-6) \times (-9) \times (-12)$

2. Find the product

A.  $9 \times (-3) \times (-6)$

D.  $(3) \times (-6) \times (-9) \times (-12)$

B.  $(-12) \times (-13) \times (-5)$

E.  $15625 \times (-2) \times 0 \times (-15625) \times 98$

C.  $(-1) \times (-2) \times (-3) \times (-4) \times (-5)$

### Division of integers



#### Activity 2.5.4

1. Check the quotient of the following using product, the first is done for you

A.  $20 \div 4 = 5$ , hence  $20 = 5 \times 4$

B.  $-36 \div 9 = -4$

D.  $-8 \div (-2) = 4$

C.  $84 \div (-7) = -12$

E.  $-7 \div 2$

2. Referring the above activity

A. The quotient of negative and positive numbers is (Positive/negative)

B. The quotient of positive and negative numbers is (Positive/negative)

C. The quotient of negative and negative numbers is Positive/negative)

**Division** is an inverse operation of multiplication

For example, divide 12 by 3 means find a number that gives product 12 when it is multiplied by 3. The number is 4 and, because In the division  $4 \times 3 = 12$

• 12 is **dividend**

• 4 is **quotient**

• 3 is **divisor**

**Note:**

For any number  $a$ ,  $b$  and  $c$  where  $b \neq 0$ ,  $a \div b = c$ , if and only if  $a = b \times c$

$a \div b$  read as  $a$  divided by  $b$

$a \div b$  is also denoted by  $\frac{a}{b}$  or  $a/b$

**Example 1:**

Find the quotient by converting in to products

- A.  $8 \div 2$     B.  $12 \div (-4)$     C.  $-10 \div (-5)$     D.  $-8 \div 4$

**Solution:**

A.  $8 \div 2 =$  , means  $8 = 2 \times ?$ , the number is 4, because  $8 = 2 \times 4$  Hence,  $8 \div 2 = 4$

B.  $12 \div (-4) =$  , means  $12 = -4 \times ?$ , the number is -3, because  $12 = -4 \times -3$  Hence,  $12 \div (-4) = -3$

C.  $-10 \div (-5) =$  , means  $-10 = -5 \times ?$ , the number is 2, because  $-10 = -5 \times 2$ , Hence,  $-10 \div (-5) = 2$

D.  $-8 \div 4 =$  , means  $-8 = 4 \times ?$ , the number is -2, because  $-8 = 4 \times (-2)$  Hence,  $-8 \div 4 = -2$

From the above example,



The quotient of **two positive** numbers is **positive**



The quotient of **two negative** numbers is **positive**



The quotient **positive** and **negative** integers is **negative**

To divide integers without converting in to multiplication follow the following procedure:

1. If the sign of dividend and divisor are the same:



The sign of the quotient is (+)



Find the magnitude of the quotient



Put the sign in front of the quotient

2. If the sign of dividend and divisor is different:



The sign of the quotient is (-)



Find the magnitude of the quotient



Put the sign in front of the quotient

### Example 2

*calculate the quotient of the following*

A.  $-368 \div 8$

C.  $175 \div (-7)$

E.  $-1 \div 4$

B.  $\frac{-264}{12}$

D.  $\frac{8211}{-21}$

F.  $24 \div (-5)$

### Solution

A.  $-368 \div 8 \rightarrow$  The result is negative sign (-)

$$368 \div 8 = -48$$

$$\Rightarrow -368 \div 8 = -48$$

B.  $\frac{-264}{12} \rightarrow$  The result is negative sign (-)

$$264 \div 12 = 22$$

$$\Rightarrow \frac{-264}{12} = -22$$

C.  $175 \div (-7) \rightarrow$  The result is negative sign (-)

$$175 \div 7 = 25$$

$$\Rightarrow 175 \div (-7) = -25$$

d.  $\frac{8211}{-21} \rightarrow$  The result is negative sign (-)

$$8211 \div 21 = 391$$

$$\frac{8211}{-21} = -391$$

**Example 3** Calculate the quotient of the following

- A.  $-1 \div 4$       B.  $-24 \div (-5)$       C.  $-5 \div 4$       D.  $-150 \div (-20)$

**Solution**

A.  $-1 \div 4 \rightarrow$  The result is negative sign (-)

$$1 \div 4 = 0.25$$

$$\Rightarrow -1 \div 4 = -0.25$$

B.  $-24 \div (-5) \rightarrow$  The result is negative sign (-)

$$24 \div 5 = 4.8$$

$$\Rightarrow -24 \div (-5) = 4.8$$

C.  $-5 \div 4 \rightarrow$  The result is negative sign (-)

$$5 \div 4 = 1.25$$

$$\Rightarrow -5 \div 4 = -1.25$$

D.  $-150 \div (-20) \rightarrow$  The result is positive sign (+)

$$150 \div 20 = 7.5$$

$$\Rightarrow -150 \div (-20) = 7.5$$

**Example 4** Find the quotient of the following

- A.  $-387 \div (-9)$       B.  $\frac{-1984}{-16}$

**Solution:-**

A.  $-387 \div (-9) \rightarrow$  The result is positive sign (+)

$$387 \div 9 = 43$$

$$\Rightarrow -387 \div (-9) = 43$$

B.  $\frac{-1984}{-16} \rightarrow$  The result is positive sign (+)

$$= -1984 \div 16 = 124 \quad \Rightarrow \frac{-1984}{-16} = 124$$

### Division of integers is summarized in the following table

Dividend and divisor	Sign of quotient	Example
Both positive	Positive	$15 \div 3 = 5$ or $\frac{15}{(3)} = 5$
Opposite sign	Negative	$15 \div (-3) = -5$ or $\frac{15}{(-3)} = -5$
Both negative	Positive	$-15 \div (-3) = 5$ or $\frac{-15}{(-3)} = 5$

### Properties of division of integers

Properties	Examples
1. The quotient of any two integers is not necessarily an integer	$20 \div 8 = 2.5 = \frac{5}{2}$
2. A non-zero integer divided by itself is 1	$20 \div 20 = 1$
3. An integer divided by zero is <b>not defined</b> . i.e Any integer cannot be divided by zero	$\frac{a}{0}$ = undefined is not defined
4. Zero divided by any non-zero integer is zero.	$\frac{0}{20} = 0$
5. An integer divided by 1 is itself the number.	$\frac{20}{1} = 20$

### Exercise 2.5.4

1. Identify dividend, divisor and quotient of the following

A.  $-96 \div 12 = -8$       B.  $\frac{(-56420)}{-124} = 455$

2. Find the quotient by converting in to products

A.  $18 \div 3$       B.  $24 \div (-4)$       C.  $-40 \div (-5)$       D.  $-44 \div 4$

3. Divide

A.  $-130 \div 5$       C.  $(-1048) \div (-1048)$       E.  $768 \div 0$       G.  $0 \div 0$   
 B.  $0 \div (-4965)$       D.  $\frac{-2574}{33}$       F.  $\frac{-17622}{-99}$

4. Fill the blank space

A.  $56 \div \underline{\hspace{1cm}} = 7$

B.  $-1092 \div \underline{\hspace{1cm}} = 14$

C.  $\underline{\hspace{1cm}} \div (-18) = 46$

D.  $\underline{\hspace{1cm}} \div (55) = -101$

5. Divide the following

A.  $-12 \div 5$

B.  $-37 \div (-2)$

C.  $-26 \div (-8)$

D.  $45 \div (-4)$

## 2.6. Even and Odd integers

### Learning Outcomes:

At the end of this unit, learners will be able to:

→ Describe even and odd integers

→ Identify even and odd integers



### Activity 2.6.1

Discuss with your friends/partners

1. Group the following objects by two

A. Picture of six balls

C. Picture of 4 rulers

B. Picture of five books

D. Picture of 3 pens

2. From activity 1a and c, explain what you observe

3. From activity 1b and d, explain what you observe

4. What do we call numbers that can be grouped by two (divisible by two) without leaving remainder?

5. What do we call numbers that can be grouped by two leaving one remainder?

6. Identify the following numbers as even and odd     **6, 5, 4, 3**

Even integers = \_\_\_\_\_

Odd integers = \_\_\_\_\_

**Even integer** is an integer that can be divisible by 2 without leaving remainder.

**Example 1:** The following are single digit even integers

$-8, -6, -4, -2, 0, 2, 4, 6,$  and  $8$

**Odd integer** is an integer that cannot be divisible by 2. When you divide any odd integer by 2, its remainder is 1.

**Example 2:** The following are single digit odd integers

$-9, -7, -5, -3, -1, 1, 3, 5, 7,$  and  $9$

### **Many digits even and odd integers**

For many digits number:

If the unit digit is  $0, 2, 4, 6,$  or  $8$ , then the number is even integer, otherwise it is odd.

### **Example 3:**

Identify the following numbers as even and odd integers

$20, -16, 31, -23, 14, -15, 345, -252$

### **Solution:**

The unit digit of even integer is  $0, 2, 4, 6,$  or  $8$ , hence

$20, -16, 14$  and  $-252$  are even integers.

The rest  $31, -23, -15,$  and  $345$  are odd integers

### **Example 4:**

list three consecutive even integers next to 31

### **Solution:**

$32, 34,$  and  $36$

### Example 5:

list the next three consecutive even integers greater than -6

**Solution:**  $-4, -2, \text{ and } 0$

### Exercise 2.6.1

1. Identify the following integers as even and odd  
 $-36, 55, 0, -71, 103, -180$
2. Identify the following integers as even and oddS  
 $3675, 5568, -7109, 1034, -1807$
3. List the next three consecutive even integers greater than 143
4. List the next three consecutive odd integers greater than -54
5. List an even integer which is neither positive nor negative
6. What is the smallest positive even integer
7. What is the greatest odd negative integer
8. What is the greatest three-digit positive integer
9. What is the smallest two-digit negative integer

### Addition of even and odd integers

**Example 1:** add the following integers

- A.  $-40 + 36$     B.  $24 + (-18)$     C.  $-4 + 4$     D.  $246 + 344$

**Solution:**

- A.  $-40 + 36 = -(40 - 36) = -4$     C.  $-4 + 4 = 0$   
 B.  $24 + (-18) = +(24 - 18) = 6$     D.  $246 + 344 = 590$

From above example, the sum of two even integers is always  
(even / odd)

**Example 2:** add the following integers

- A.  $31 + 25$     B.  $-45 + 23$     C.  $11 + (-33)$     D.  $-21 + (-15)$

**Solution:**

- A.  $31 + 25 = 56$     C.  $11 + (-33) = -(33 - 11) = -22$   
 B.  $-45 + 23 = -(45 - 23) = -22$     D.  $-21 + (-15) = -(21 + 15) = -36$

From above example, the sum of two odd integers is always \_\_\_\_\_  
(even/odd)

**Example 3: add the following integers**

A.  $8 + 19$     B.  $-4 + (-23)$     C.  $17 + (-32)$     D.  $-21 + 30$

***Solution:***

A.  $8 + 19 = 27$

C.  $17 + (-32) = -(32 - 17) = -5$

B.  $-4 + (-23) = -(4 + 23) = -27$     D.  $-21 + 30 = +(30 - 21) = 9$

*From above example, the sum of odd and even integer is always \_\_\_\_ (even/odd)*

***Subtraction of even and odd integers*****Example 4: subtract the following integers**

A.  $8 - 6$     B.  $12 - 20$     C.  $-10 - 10$     D.  $-6 - (-30)$

***Solution:***

A.  $8 - 6 = 2$

B.  $12 - 20 = 12 + (-20) = -(20 - 12) = -8$

C.  $-10 - 10 = -10 + (-10) = -(10 + 10) = -20$

D.  $-6 - (-30) = -6 + 30 = +(30 - 6) = 24$

*From the above example, we can conclude that*

*even - even = \_\_\_\_\_*

**Example 5: Subtract the following integers**

A.  $7 - 5$     B.  $11 - 23$     C.  $-15 - 13$     D.  $-9 - (-11)$

***Solution:***

A.  $7 - 5 = 2$

B.  $11 - 23 = 11 + (-23) = -(23 - 11) = -12$

C.  $-15 - 13 = -15 + (-13) = -(15 + 13) = -28$

$$D. -9 - (-11) = -9 + 11 = +(11 - 9) = 2$$

*From the above example, we can conclude that*

• *odd - odd = \_\_\_\_\_*

**Example 6:** *subtract the following integers*

A.  $12 - 5$     B.  $31 - 50$     C.  $-18 - 15$     D.  $-22 - (-9)$ .

**Solution:**

A.  $12 - 5 = 7$

B.  $31 - 50 = 31 + (-50) = -(50 - 31) = -19$

C.  $-18 - 15 = -18 + (-15) = -(18 + 15) = -33$

D.  $-22 - (-9) = -22 + 9 = -(22 - 9) = -13$

*From the above example, we can conclude that*

• *odd - even = \_\_\_\_\_*                      • *even - odd = \_\_\_\_\_*

**Product of even and odd integers**

**Example 1:** *Multiply the following integers*

A.  $12 \times 4$

C.  $-6 \times 4$

E.  $-8 \times (-4)$

B.  $0 \times (-6)$

D.  $-2 \times 10$

**Solution:**

A.  $12 \times 4 = 48$

C.  $-6 \times 4 = -24$

E.  $-8 \times (-4) = 32$

B.  $0 \times (-6) = 0$

D.  $-2 \times 10 = -20$

*From the above example, we can conclude that even  $\times$  even = \_\_\_\_\_*

**Example 2:** *Multiply the following integers*

A.  $2 \times 3$

C.  $-7 \times 4$

E.  $-9 \times (-4)$

B.  $0 \times (-5)$

D.  $-3 \times 8$

**Solution:**

A.  $2 \times 3 = 6$

C.  $-7 \times 4 = -28$

E.  $-9 \times (-4) = 36$

B.  $0 \times (-5) = 0$

D.  $-3 \times 8 = -24$

From the above example, we can conclude that

•  $odd \times even = \underline{\hspace{2cm}}$

•  $even \times odd = \underline{\hspace{2cm}}$

**Example 3: Multiply the following integers**

A.  $5 \times 3$

C.  $-7 \times 11$

E.  $-9 \times (-5)$

B.  $1 \times (-5)$

D.  $-13 \times 3$

**Solution:**

A.  $5 \times 3 = 15$

C.  $-7 \times 11 = -77$

E.  $-9 \times (-5) = 45$

B.  $1 \times (-5) = -5$

D.  $-13 \times 3 = -39$

From the above example, we can conclude that

•  $odd \times odd = \underline{\hspace{2cm}}$

**Exercise 2.6.2**

1. Fill in the blank space with correct answer

A.  $even + even = \underline{\hspace{2cm}}$

E.  $odd - even - odd + odd = \underline{\hspace{2cm}}$

B.  $even - odd = \underline{\hspace{2cm}}$

F.  $odd \times odd \times even = \underline{\hspace{2cm}}$

C.  $odd \times even = \underline{\hspace{2cm}}$

G.  $even \times odd + even = \underline{\hspace{2cm}}$

D.  $even + odd + even = \underline{\hspace{2cm}}$

H. The difference between any two consecutive even integers is equal to  $\underline{\hspace{2cm}}$ .

i. The difference between any two consecutive odd integers is equal to  $\underline{\hspace{2cm}}$ .

## Summary for unit two

- ☞ Natural number is a set of numbers denoted by  $\mathbb{N}$  and is described by  $\mathbb{N} = \{1, 2, 3, \dots\}$
- ☞ Whole numbers are numbers contains natural number and zero denoted by  $\mathbb{W}$  and is described by  $\mathbb{W} = \{0, 1, 2, 3, \dots\}$
- ☞ An integer is a set of numbers consisting of whole numbers and negative numbers. The set of integers is denoted by  $\mathbb{Z} = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$
- ☞ Every negative integer is less than any positive integer.
- ☞ Every positive integer is greater than any negative integer.
- ☞ Successor of an integer is an integer that comes after the given integer
- ☞ Predecessor of an integer is an integer that comes before the given integer.
- ☞ The sum of any two integers is always an integer.
- ☞ The difference of any two integers is always an integer.
- ☞ The sum of two opposite integers is zero.
- ☞ The product of positive and negative integer is negative integer.
- ☞ The product of an even number of negative factors is positive.
- ☞ The product of an odd number of negative factors is negative.
- ☞ The product of any two integers is always an integer.
- ☞ Any negative integer divided by negative integer is always positive
- ☞ Even integer is an integer that can be divisible by 2 without leaving remainder..

## Review Exercise for unit 2

1. Represent the following integers on number line

A. -12, 3, 6, 8, -5, -9      B. -11, 6, -4, -8      C. -2, 3, 4, -3, -10

2. Show the relation between  $\mathbb{N}$ ,  $\mathbb{W}$   $\mathbb{Z}$ ?

3. List down 3 example of non-integer numbers?

4. Find the opposite of the following integers?

A. -80      C. -34      E. 835      G. -1020      I. -768

B. 34      D. 1002      F. 543      H. -90

5. List down all integers that lie between the following pair of integers

A. -70 and -60      C. -3 and -4      E. 0 and 7

B. -34 and -34      D. -12 and -10      F. -12 and 0

6. Compare the following pair of integers using the sign  $<$ ,  $=$ , and  $>$

A. -65 \_\_\_\_\_ 0      D. 1002 \_\_\_\_\_ -1002      G. -1020 \_\_\_\_\_ 220

B. 34 \_\_\_\_\_ -678      E. -835 \_\_\_\_\_ -835      H. -90 \_\_\_\_\_ -65

C. -34 \_\_\_\_\_ -304      F. 543 \_\_\_\_\_ -56      I. -768 \_\_\_\_\_ 34

7. If  $-5 > x$ , then list down four possible values of  $x$ , where  $x$  is an integer.

8. Arrange the following integers in descending and ascending order

.  $-1, -5, 1, -51, -69, 101, -101, 21, -81$

9. Find the following sum

A.  $0+(-59)$

D.  $-545+511$

G.  $-1020 + 20$

B.  $26+(-52)$

E.  $-215+(-349)$

H.  $-90+(-489)+315$

C.  $-95+124$

F.  $121+(-290)$

I.  $-768+2001$

10. Find the following difference

A.  $0 - 592$

C.  $-89-482$

E.  $-1099-(-412)$

B.  $75-105$

D.  $513-(-109)$

F.  $299-1251-(-191)$

11. Multiply

A.  $10 \times (-24)$

D.  $-4 \times (-25) \times 14(-1)$

B.  $-45 \times (-12)$

E.  $-9 \times (-55) \times 998 \times 0 \times 32$

C.  $-9 \times (-24)$

F.  $-1 \times (-1) \times 1 \times (-1) \times (-1) \times (-1) \times (-1)$

12. Name the property of the following for any integers  $m, n, \&p$ .

A.  $m \times n = n \times m$

E.  $1 \times m = 1$

B.  $m \times (n + p) = (m \times n) + (m \times p)$

D.  $0 \times m = 0$

C.  $m \times (n \times p) = (m \times n) \times p$

13. Find the quotient of the following

A.  $1080 \div 8$

D.  $0 \div 492$

G.  $1 \div 20$

B.  $-214 \div (-214)$

E.  $-492 \div 0$

H.  $-54 \div 4$

C.  $-19225 \div -1$

F.  $-7632 \div 36$

I.  $-110 \div (-20)$

- 14. Write four consecutive negative even integers.*
- 15. Write all odd integers between -106 and -96*
- 16. Decide whether the following statements are Even or odd*
- A. The sum of any two even integers*
  - B. The difference of any two odd integers*
  - C. The product of any two even integers*
  - D. The product of any three odd integers*
  - E. The sum of even and odd integers*
  - F. The difference of odd and even integers*
  - G. The product of odd and even integers*
- 17. Find three integers whose sum is zero.*
- 18. List examples of a pair of integers whose sum is zero.*
- 19. Abel was playing a two-round game in which he could gain or loss points. During the first round he lost 30 points. During the second round gained 13 points. What was his net (total) score at the end of the game?*

# UNIT 3

## RATIO, PROPORTION AND PERCENTAGE

### *Learning Outcomes:*

At the end of this unit, learners will be able to:

- ⇒ understand the notions of ratio and proportions.
- ⇒ solve problems involving ratio and proportion
- ⇒ describe a percentage
- ⇒ solve problems involving percentages.
- ⇒ relate fractions, decimals and percentages to real life situations
- ⇒ apply the concept of percentage in solving real life problems

### *Introduction*

In this unit you will learn the basic mathematical concepts that will be applied in business, like profit, loss, simple and compound interest.

The unit has three sections. The first section deals with the concept of ratio and proportion. The second section deals with percentages and the third section is about application of ratio, proportion and percentage. Here you will see how to calculate loss, profit, simple interest, compound interest, income tax, VAT, and turnover tax.

### 3.1. RATIO AND PROPORTION

At the end of this section, learners will be able to:

- understand the notions of ratio and proportions.
- solve problems involving ratio and proportion
- describe a percentage
- solve problems involving percentages.
- relate fractions, decimals and percentages to real life situations
- apply the concept of percentage in solving real life problems.

#### 3.1.1 RATIO



##### Activity 3.1.1

Discuss with your friends

1. write a simple ratio for each of the following


A. Cats to hens



B. What are the ratio books to pens to pencils?



2. consider the following figure



a. What is the ratio of shaded part to unshaded part?




b. What is the ratio of shaded part to the whole part?

3. Can you define a ratio based on the above activities in your own words?





4. What is the ratio of girls to boys in your class?

### Definition 3.1:

**The method of comparing two or more quantities of the same kind and in the same units by division is known as a ratio.**

-  The symbol to denote the ratio is denoted by :
-  The ratio of one quantity  $a$  to another quantity  $b$  is usually denoted by  $a:b$ . Where,  $a$  is called antecedent and  $b$  is called consequent.
-  The ratio of  $a$  to  $b$  can be written as any one of the following forms:

- i.  $a : b \rightarrow$  read as a ratio  $b$  or  $a$  to  $b$
- ii.  $a/b \rightarrow$  read as  $a$  over  $b$
- iii.  $a \div b \rightarrow$ ,  $a$  divided by  $b$ , provided that  $b \neq 0$

-  In most cases the ratio  $a:b$  is written in simplified form, where  $a$  and  $b$  are natural numbers.
-  To write the ratio of two quantities, the quantities must have the same units of measurement.
-  Quantities having different units of measurement cannot be compared in the form of ratio.
-  Ratio of quantities have no unit (unitless).

**Example 1:** In a class there are 18 boys and 24 girls.

- A. What is the ratio of boys to girls?
- B. What is the ratio of girls to boys?
- C. What is the ratio of girls to the total number of students in the class?
- D. What is the ratio of boys to the total number of students in the class?

**Solution:**

Total number of students = number of boys + number of girls

$$=18+24=42$$

A. Ratio of boys to girls =  $\frac{\text{number of boys}}{\text{number of girls}}$

$$\frac{18}{24} = \frac{18 \div 6}{24 \div 6} = \frac{3}{4} \text{ or } 3:4$$

B. Ratio of girls to boys =  $\frac{\text{number of girls}}{\text{number of boys}}$

$$\frac{24}{18} = \frac{24 \div 6}{18 \div 6} = \frac{4}{3} \text{ or } 4:3$$

C. Ratio of girls to Total number of students =  $\frac{\text{number of girls}}{\text{number of students}}$

$$\frac{24}{42} = \frac{24 \div 6}{42 \div 6} = \frac{4}{7} \text{ or } 4:7$$

D. Ratio of boys to Total number of students =  $\frac{\text{number of boys}}{\text{number of students}}$

$$\frac{18}{42} = \frac{18 \div 6}{42 \div 6} = \frac{3}{7} \text{ or } 3:7$$

### Example 2 :-

Write down the ratio of the first number to the second one, in the Simplest form.

A. 4 to 8      B. 54 to 128      C. 48 to 6      D. 540 to 48

*Solution:*

**A.** Ratio of 4 to 8

$$\frac{4}{8} = \frac{4 \div 4}{8 \div 4} = \frac{1}{2} \text{ or } 1:2$$

**C.** Ratio of 48 to 6

$$\frac{48}{6} = \frac{48 \div 6}{6 \div 6} = \frac{8}{1} \text{ or } 8:1$$

**B.** Ratio of 54 to 128

$$\frac{54}{128} = \frac{54 \div 2}{128 \div 2} = \frac{27}{64} \text{ or } 27:64$$

**D.** Ratio of 540 to 48

$$\begin{aligned} \frac{540}{48} &= \frac{540 \div 2}{48 \div 2} = \frac{270 \div 2}{24 \div 2} \\ &= \frac{135 \div 3}{135 \div 3} = \frac{45}{4} \text{ or } 45:4 \end{aligned}$$

**Example 3** Find the ratio of the following and express simplest form of fraction.

A.  $\frac{3}{4}$  to  $\frac{5}{3}$     B.  $\frac{2}{13}$  to  $\frac{15}{16}$     C. 3.2 to 1.2    D.  $\frac{7}{9}$  to 0.7

**Solution:**

**A. Ratio of  $\frac{3}{4}$  to  $\frac{5}{3}$  is**  $\frac{\frac{3}{4}}{\frac{5}{3}} = \frac{3 \times 3}{4 \times 5} = \frac{9}{20} = 9:20$

**B. Ratio of  $\frac{2}{13}$  to  $\frac{15}{16}$  is**  $\frac{\frac{2}{13}}{\frac{15}{16}} = \frac{2 \times 16}{13 \times 15} = \frac{32}{195} = 32:195$

**C. Ratio of 3.2 to 1.2 is**  $\frac{3.2}{1.2} = \frac{32}{12} \times \frac{10}{10} = \frac{32 \div 4}{12 \div 4} = \frac{8}{3} \text{ or } 8:3$

**D. Ratio of  $\frac{7}{9}$  to 0.7 is**  $\frac{\frac{7}{9}}{0.7} = \frac{\frac{7}{9}}{\frac{7}{10}} = \frac{7 \times 10}{9 \times 7} = \frac{10}{9} = 10:9$

**Example 4. Find the ratio of the following quantities.**

A. 65 cents to 5 Birr

C. 700 grams to 6kg

B. 12 meters to 200cm

D. 600 seconds to 8 minutes

**Solution:** First change the quantities in to the same units

A. 1 Birr=100cent then 5 Birr =500cent

The ratio of 65 cents to Birr 5= Ratio of 65 cents to 500cent

$$\frac{\cancel{65 \text{ cents}}}{500 \cancel{\text{cents}}} = \frac{65 \div 5}{500 \div 5} = \frac{13}{100} = 13:100$$

B. 1m=100cm then 12m=1200cm

The ratio of 12m to 200cm = Ratio of 1200cm to 200cm

$$\frac{\cancel{1200\text{ cm}}}{200\cancel{\text{cm}}} = \frac{12 \div 2}{2 \div 2} = \frac{6}{1} = 6:1$$

C. 1kg=1000grams then 6kg=6000grams

The ratio of 700gram to 6kg = Ratio of 700 gram to 6000gram

$$\frac{\cancel{700\text{ gram}}}{\cancel{6000}\text{gram}} = \frac{7}{60} = 7:60$$

D. 1minute=60seconds then 8minute=480 second

The ratio of 600 second to 8minutes = Ratio of 600 second to 480 second

$$\frac{\cancel{600\text{ second}}}{\cancel{480}\text{second}} = \frac{60 \div 12}{48 \div 12} = \frac{5}{4} = 5:4$$

### Exercise 3.1.1

1. Write down the ratio of the first number to the second one in the simplest form

A. 48 and 80

C. 4.8 and 9.6

E.  $\frac{2}{21}$  and  $\frac{8}{21}$

B. 360 and 72

D. 13.6 and 10.2

F.  $2\frac{11}{12}$  and  $1\frac{2}{3}$

2. Express the following ratios as in their lowest term:

A. 4 Birr to 16 cents

B. 2 liters to 2250 milliliters

C. 5 days to 100 hours

D. 3 hours and 30minutes to 180 minutes

E. 3.5 kg to 6500 grams


F. 3 hectares to 3000 square meters

3. Using the figure below answer the following questions

A. What is the ratio of shaded part to unshaded part?

B. What is the ratio of shaded part to the whole part?

C. What is the ratio of unshaded part to the whole part?



**Note:**

1 liter = 1000 milliliters

1 hectare = 10,000 square meters

1 day = 24 hours

1kg = 1000gm

## Dividing the given quantity in the given ratio

### Examples:

1. Divide Birr 300 with ratio 3: 2

**Solution:** let the first part is  $3x$  and the second part is  $2x$ , then

$$3x + 2x = 300 = \frac{5x}{5} = \frac{300}{5} = x = 60$$

**Therefore,** the first part is  $3x = 3 \times 60 = 180$  and the second part is  $2x = 2 \times 60 = 120$  Birr

2. The ratio of two numbers is 7:3 and their sum is 50, then find the two numbers.

**Solution:** let the first number is  $7x$  and the second number is  $3x$ , then

$$\begin{aligned} 7x + 3x &= 50 \\ \frac{10x}{10} &= \frac{50}{5} = x = 5 \end{aligned}$$

**Therefore,** the first number is  $7x = 7 \times 5 = 35$  and the second number is  $3x = 3 \times 5 = 15$ .

3. Two numbers have ratio 13:10. If their difference is 84, then find the larger number.

**Solution:** let the larger number is  $13x$  and the smaller number is  $10x$ ,

$$\begin{aligned} \text{then } 13x - 10x &= 84 \\ 3x &= 84 \\ x &= 28 \end{aligned}$$

**Therefore,** the larger number is  $13x = 13 \times 28 = 364$

4. A bag contains red and white balls; the ratio of red to white balls is 3:4. If the bag contains 36 white balls, then find the number of red balls.

**Solution:**

The number of red balls is  $3x$  and the number of white balls is  $4x$

$$\begin{aligned} \text{Number of White balls } 4x &= 36 \\ \frac{4x}{4} &= \frac{36}{4} = x = 9 \end{aligned}$$

Therefore, the number of red balls is  $3x = 3 \times 9 = 27$

5. The ratio of the measures of the angles of a triangle is 1:2:3. Then find the measure of each angle.

**Solution:**

*let the angles of the triangle are  $1x, 2x$  and  $3x$*

*$1x + 2x + 3x = 180^\circ$  ... Sum of interior angle of a triangle is  $180^\circ$*

$$\begin{array}{r} 6x = 180^\circ \\ \cancel{6x} = \frac{180^\circ}{6} \\ x = 30^\circ \end{array}$$

**Therefore:** *the first angle is  $1x = 1 \times 30^\circ = 30^\circ$*

*Second angle is  $2x = 2 \times 30^\circ = 60^\circ$*

*Third angle is  $3x = 3 \times 30^\circ = 90^\circ$*

6. If a, b, and c are numbers such that a:b:c = 3:4:5 and b=20, then find a+b+c

**Solution:**

*let  $a = 3x$ ,  $b = 4x$  and  $c = 5x$*

$$b = 4x = 20$$

$$\begin{array}{r} \frac{4x}{4} = \frac{20}{4} \\ x = 5 \end{array}$$

*Therefore,  $a = 3x = 3 \times 5 = 15$ ,*

$$c = 5x = 5 \times 5 = 25$$

$$\text{And } a + b + c = 15 + 20 + 25 = 60$$


### Exercise 3.1.2

- Find two numbers whose ratio is 3 to 5 and whose sum is 192.
- A wire of length 240 cm is cut into 3 pieces, in the ratio 1:2:5. Find the length of each piece.
- Aster, Fatuma, Mohammed and Yared contribute the sum of money to Renaissance dam in the ratio 1:3:5:7. If the largest amount contributed is Birr 1050. Calculate the amount contributed by each person.

4. Two numbers have ratio 12:5. Their difference is 98. Find the larger number.

5. If  $a : b = 3:2$ , find the value of  $\frac{a+b}{a-b}$ .

6.. Find the ratio of the areas of the squares ABCD to that of PQRS where  $AB = 12\text{cm}$  and  $PQ = 8\text{cm}$



7.  $x, y$  and  $z$  are numbers such that  $x:y:z = 1:2:4$  and  $y = 48$ .

Find  $x+y+z$

### 3.1.2 Proportion



#### Activity 3.1.2

Discuss with your friends

1. Compare the following ratios A. 4.6 and 6.9 B.  $\frac{3}{6}$  and  $\frac{40}{80}$
2. What result did you get from above activities?
3. What do you call equality of two ratios?

#### Definition 3.2:

**proportion** is equality of two ratios.

**Example 1** compare the ratio 6:10 and 48:80, do they form proportion?

**Solution:**

A. the simplest form of 6:10 is

$$\frac{6 \div 2}{10 \div 2} = \frac{3}{5} = 3:5$$

B. The simplest form of  $48:80$  is  $\frac{48 \div 16}{10 \div 16} = \frac{3}{5} = 3:5$

Therefore,  $6:10 = 48:80$  is proportion

**Example 2:** compare the ratio  $1:3$  and  $5:10$ , do they form proportion?





**Solution:**

the simplest form of  $5:10$  is  $\frac{5 \div 5}{10 \div 5} = \frac{1}{2} = 1:2$

therefore, the two ratios are not equal (i.e.  $1:3 \neq 5:10$ )

the two ratios do **not form proportion**.

**Note:**

-  If the four quantities  $a, b, c,$  and  $d$  are in proportion, then  $a:b=c:d$ .
-  The proportion  $a:b=c:d$  can be written as  $\frac{a}{b} = \frac{c}{d}$ , here  $a$  and  $d$  are called **extremes** (end terms) and  $b$  and  $c$  are said to be **means** (middle terms)
-  In proportion the product of means is equal to the product of extremes. (i.e. if  $\frac{a}{b} = \frac{c}{d}$ , then  $a \times d = b \times c$ ).
-  you can easily remember this by applying, cross multiplication as follows If  $\frac{a}{b} = \frac{c}{d}$ , then  $a \times d = b \times c$  or  
If  $a:b = c:d$  then  $a \times d = b \times c$

**Example 3:** show that  $20, 45, 8$  and  $18$  are in order of proportion

**Solution:**

to show they are in order of proportion (i.e.  $20 : 45 = 8 : 18$ )

Use cross multiplication (product of means = product of extremes)

$$\text{Product of means} = 45 \times 8 = 360$$

$$\text{Product of extremes} = 20 \times 18 = 360$$

Since, product of means is equal to product of extremes, the numbers 20, 45, 8 and 18 **are in order of proportion.**

**Example 4:** Are the numbers 3, 5, 6 and 12 in order of proportion?

**Solution:**

check  $3:5=6:12$ ?

$$\text{Product of means} = 5 \times 6 = 30$$

$$\text{Product of extremes} = 3 \times 12 = 36$$

Product of means  $\neq$  product of extremes, hence the numbers 3, 5, 6 and 12 are **not in order of proportion.**

**Example 5:** If the numbers 3, 5,  $x$  and 20 are in order of proportion, then find the value of  $x$ .

**Solution:**

$$3:5 = x:20$$

Product of means = product extremes

$$\frac{5x}{5} = \frac{3 \times 20}{5} \quad x = 12$$

**Example 6:** In proportion  $\frac{3}{5} = \frac{9}{y}$ , find the value of  $y$ .

**Solution:**

$$\text{use cross multiplication} \quad \frac{3}{5} = \frac{9}{y} = \frac{3y}{3} = \frac{9 \times 5}{3} = y = 15$$

**Example 7:** The distance from Addis Ababa to Adama is 75 km.

If a map scale is 1:1,000,000, then find the distance between the two cities on a map, in centimeters.

**Solution:**

Map scale = Map distance : ground distance

**Note**

Map scale = map distance: ground distance

1km = 100, 000cm

7.5km = 7,500,000 cm

$$\begin{aligned} 1:1,000,000 &= x:7,500,000 \text{ cm} \\ \frac{\cancel{1,000,000} x}{1,000,000} &= \frac{\cancel{7,500,000} \text{ cm}}{1,000,000} \\ x &= \frac{75}{10} \\ x &= 7.5 \text{ cm} \end{aligned}$$

Therefore, the distance between the two cities on a map 7.5cm

**Exercise 3.1.3**

1. From each pair of ratios below circle the given pair of ratios that are in proportion.

A.  $\frac{2}{3}$  and  $\frac{4}{260}$     B.  $\frac{1}{3}$  and  $\frac{5}{20}$     C.  $\frac{4}{5}$  and  $\frac{12}{20}$     D.  $\frac{8}{4}$  and  $\frac{2}{1}$

2. Find the unknown terms in each of the following

A.  $\frac{3}{5} = \frac{x}{20}$     B.  $x:18 = 2:9$     C.  $\frac{8}{m} = \frac{3.2}{4}$     D.  $\frac{3}{5} = \frac{18}{n} = \frac{y}{50}$

3. Show that the numbers 14,21,2,and 3 are in order of proportion.

4. Given the proportion  $10:18=35:63$ , then find

A. The sum of means

C. the sum of extremes

B. The product of means

D. the product of extremes

5. A map scale is 1:1,000,000. If the distance between the two cities on a map is 3.5cm then find the actual distance between the cities in kilometers.

## Direct and inverse proportionality

### A. Direct proportionality

#### Definition 3.3:

*y is said to be directly proportional to x (written as  $y \propto x$ ), if*

*there is constant  $k$  such that  $y = kx$  or  $k = \frac{y}{x}$*

*The number  $K$  is called constant of proportionality.*

*If  $y \propto x$ , then as  $x$  increases  $y$  also increases or as  $x$  decreases also  $y$  decreases. And the ratio  $\frac{y}{x}$  is constant.*

**Example 1:** The following table shows the number of kilograms of sugar bought and the price paid in birr.

sugar in Kg ( $x$ )	1	2	3	4
price in Birr ( $y$ )	30	60	90	120

- Is  $y$  directly proportional to  $x$ , if so what is the constant of proportionality?
- Write the formula relating  $y$  and  $x$ .
- Using the formula calculate the price of 6.5 Kg sugar.
- How much kg sugar will be bought with Birr 315.

**Solution:**

A. As  $x$  increases  $y$  also increases and the ratio  $\frac{y}{x}$  is constant,

$$k = \frac{30}{1} = \frac{60}{2} = \frac{90}{3} = \frac{120}{4}, \text{ hence } y \propto x$$

B.  $\frac{y}{x} = k$  implies  $\frac{y}{x} = 30$  or  $y = 30x$ .

C.  $y = ?$ , when  $x = 6.5$  Kg,

$$y = 30x$$

$$y = 30 \times 6.5 \text{ Kg} = y = \text{birr } 195$$

D.  $x = ?$ , when  $y = 315 \text{ Birr}$ , then  $\frac{y}{x} = 30$

$$y = 30 X =$$

$$\frac{315}{30} = \frac{30x}{30} \quad x = 10.5 \text{ Kg}$$

**Example 2:** if  $y \propto x$ , then find the value of m

x	12	6
y	48	m

**Solution:**

since  $y \propto x$ , the ratio  $\frac{y}{x}$  is constant

$$\frac{48}{12} = \frac{m}{6} = \frac{12 \times m}{12} = \frac{48 \times 6}{12} = m = 24$$

**Example 3:** If 5kg sweet potato costs Birr 120, then calculate the cost of 8 kg sweet potato.

**Solution:**

as the amount of sweet potato increases, the cost also increases

Hence, the cost of sweet potato is directly proportional to amount of sweet potato.

sweet potato in Kg	5	8
cost in (birr)	120	x

$$\frac{120}{5} = \frac{x}{8} = \frac{5x}{5} = \frac{120 \times 8}{5} = x = 192 \text{ birr}$$

Therefore, the cost of 8kg sweet potato is Birr 192

### Exercise 3.1.4

- If  $y$  is directly proportional to  $x$ ;  $y=24$  when  $x=6$ , then find
  - the constant proportionality.
  - the formula relating  $y$  and  $x$
  - the value  $y$ , when  $x = 3$ .
  - the value  $x$ , when  $y = 15$ .
- Which of the following table shows direct proportional relation ship between  $x$  and  $y$ ?

A.

x	8	5	6
y	24	15	18

B.

x	3	7	4
y	6	14	12

- What is the value of  $m$  from the given table below if  $y \propto x$ ?

x	14	7
y	21	m

$y$  is directly proportional to  $x$ , If  $x=20$  when  $y=160$ , then what is the value of  $x$  when  $y=32$ ?

- If a car travels 120 km in 2 hours, then how long will it take for the car to travel 330 km (assuming the car is moving at constant speed)?

### B. Inverse proportionality

#### Definition 3.4:

$y$  is said to be inversely proportional to  $x$ . (written as  $y \propto \frac{1}{x}$ ), if there  $k$  is constant such that  $y = \frac{k}{x}$  or  $k = y \times x$ .

The number  $k$  is called constant of proportionality.

If  $y \propto \frac{1}{x}$ , then as  $x$  increases  $y$  decreases or as  $x$  decreases  $y$  increases. And the product  $k$  is constant.

#### Example 1:

The following table shows the time taken for a car to move a distance

of 5,000m at various speed.

time in second(x)	1,000	500	250	200
speed in m/s(y)	5	10	20	25

- A. Is y inversely proportional to x , if so what is the constant of proportionality?
- B. Write the formula relating y and x .
- C. Using the formula calculate the time taken by the car to cover the given distance moving at a speed of 8 meter/second.

**Solution:**

A. As x increases, y decreases and the product  $y \times x$  is constant.

$$k = 5 \times 1000 = 10 \times 500 = 20 \times 250 = 25 \times 200 = 5000, \text{ hence } y \propto \frac{1}{x}$$

B.  $y \times x = k$ , implies  $y \times x = 5000$

C.  $x = ?$   $y = 8 \text{ m/s}$

$$y \times x = 5000 = \frac{8 \times x}{8} = \frac{5000}{8} \quad x = 625 \text{ seconds}$$

**Example 2:**

If  $y \propto \frac{1}{x}$ , then find the value of n.

x	12	6
y	15	n

**Solution:**

since  $y \propto \frac{1}{x}$ , the product  $y \times x$  is constant.

$$\frac{6 \times n}{6} = \frac{12 \times 15}{6} \quad n = 30$$

**Example 3:** If 8 persons do a certain job in 9 hours, then how long will it take for 12 persons to do the same job. (Assume all persons do the job at the same rate)

Number of persons	8	12
Time taken to do the job(hr.)	9	x

**Solution:**

As the number of person increases, the time taken to finish the job decreases.

Hence, the number of **persons** is **inversely** proportional to **time** taken to finish the job

$$12 \times x = 8 \times 9$$

$$\frac{12 \times x}{12} = \frac{8 \times 9}{12} \quad x = 6hr$$

**Therefore, it takes 6 hr for 12 persons to do the job.**

**Exercise 3.1.5**

- If  $y \propto \frac{1}{x}$  and  $y=6$  when  $x = 4$ , then find the constant of proportionality.
- $y$  is inversely proportional to  $x$ . If  $x = 25$ , then  $y=8$ . What is the value of  $y$  when  $x = 10$ .
- Which of the following table is inversely proportional in the relationship between  $x$  and  $y$ ?

A.

$x$	2	4	8
$y$	36	18	9

B.

$X$	5	10	15
$Y$	20	10	5

- It takes 8 days for 35 laborers to harvest coffee on a plantation. How long will 20 laborers take to harvest coffee on the same plantation?
- $x$  is inversely proportional to  $y$ . If  $x= 25$ , then  $y= 15$ . Find  $y$  when  $x=30$
- 100 men working in a factory produce 6000 articles in 10 days. How long it takes (assume all workers do the job at the same rate)
  - 50 men to produce 6000 articles
  - 150 to produce 6000 articles
- A contractor appoints 36 workers to build a wall. They could finish the task in 12 days. How many days will 16 workers take to finish the same task?

### 3.2. REVISION ON PERCENTAGES

At the end of this section, learners will be able to:

- Visualizes percent and its relationship to fractions, ratios, and decimal numbers using bar models.
- Identifies the base, percentage, and rate in a problem.



#### Activity 3.2.1

Discuss with your friends

1. consider the following


- A. count the green shaded part and the total part
- B. What is the ratio of green part to total part?
- C. express the above ratio in percent form
- D. define percent in your own words



2. write the following numbers in percent form

A.  $\frac{81}{100}$    B.  $\frac{2}{100}$    c. 0.75   D.  $\frac{1}{5}$

3. Use the following bar model to relate the given fraction and percent



#### Definition 3.5:

The word **percent** means “for every hundred” or “per 100”.  
We use the symbol % to denote percent.

For example,

5 Percent symbolically written as 5 % and 5 % Means 5 per 100 or  $\frac{5}{100}$ . The concept of percent is widely used in a variety of mathematical applications.

- Interest
- Sales tax

- Change of Profit and loss
- Inflation

## Converting a given percent in fractional form

### Example 1:

Change 30 % in to fraction

### Solution:

30% means 30 per 100. So,  $30\% = \frac{30}{100} = \frac{3}{10}$

### Note:

To change a given percent to fraction form

- Drop the percent sign from the number
- Put 100 as the denominator of the number and write the fraction in the lowest term.

### Example 2:

Convert each of the following percent form in to fraction:

A. 1%      B.. 45%      C.. 90%      D. 4.8%

### Solution:

$$A. 1\% = \frac{1}{100}$$

$$C. 90\% = \frac{\cancel{90}}{\cancel{100}} = \frac{9}{10}$$

$$B. 45\% = \frac{45 \div 5}{100 \div 5} = \frac{9}{20}$$

$$D. 4.8\% = \frac{4.8}{100} = \frac{4.8 \times 10}{100 \times 10} \\ = \frac{48 \div 8}{1000 \div 8} = \frac{6}{125}$$

## Converting a given percent in decimal form

To convert a given percent in to decimal form:

First write the percent in fraction form, and then convert it in to decimal by division.

**Example 3:** Express each of the following percent form into decimal form. A. 30%      B. 7.38 %      C. 114 %      D. 11%

### Solution:

$$A. 30\% = \frac{30}{100} = 0.3$$

$$C. 114\% = \frac{114}{100} = 1.14$$

$$B. 7.38\% = \frac{7.38}{100} = 0.0738$$

$$D. 11\% = \frac{11}{100} = 0.11$$

### Exercise 3.2.1

1. Convert each of the following percent forms to fractions?

A. 60%

C. 125%

E.  $5\frac{3}{4}\%$

B. 2.6%

D.  $\frac{3}{4}\%$

F. 0.045%

2. Convert each of the following percent forms in to decimals?

A. 80%

C. 12%

E.  $5\frac{1}{4}\%$

B. 26%

D.  $\frac{2}{5}\%$

F. 0.45%

### Converting a given number to percent

**Example 4:** convert  $\frac{2}{5}$  to percent form

**Solution:**

*to convert it in to percent, try to write equivalent form of the number, so that its denominator is 100.*

$$\frac{2}{5} = \frac{2 \times 100}{5 \times 100} = \frac{40}{100} = 40\%$$

**Note:**

*To convert a given number to percent form, multiply the given number*

$$\frac{100}{100} \text{ or by } 100\%$$

**Example 5:** Convert each of the following fractions as percent form:

A.  $\frac{1}{2}$

B.  $\frac{4}{5}$

C.  $\frac{3}{4}$

D.  $\frac{3}{8}$

**Solution:**

$$A. \frac{1}{2} = \frac{1}{2} \times 100\% = \frac{100}{2} = 50\% \quad C. \frac{3}{4} = \frac{3}{4} \times 100\% = \frac{300}{4} = 75\%$$

$$B. \frac{4}{5} = \frac{4}{5} \times 100\% = \frac{400}{5} = 80\% \quad D. \frac{3}{8} = \frac{3}{8} \times 100\% = \frac{300}{8} = 37.5\%$$

**Example 6:** Express each of the following decimal into percent form

A. 0.5

B. 0.8

C. 0.24

D. 2.09

**Solution:**

$$A. 0.5 = 0.5 \times 100\% = 50\%$$

$$C. 0.24 = 0.24 \times 100\% = 24\%$$

$$B. 0.8 = 0.8 \times 100\% = 80\%$$

$$D. 2.09 = 2.09 \times 100\% = 209\%$$

**Example 7 :**

Express each of the following ratio into percent form

A. 75 seconds to 5minutes

C. 4 km to 500 meters

B. 24 hours to 8days

D. 5litres to 4000mililiters

**Solution:**

A. 75 seconds to 5minutes =

( since 5 minutes= 300 seconds)

$$75 : 300 = \frac{75}{300} = \frac{1}{4} = \frac{1}{4} \times 100\% = \frac{100}{4} = 25 \%$$

B. 24 hours to 8 days

( since 8 days= 192 hours)

$$= 24 : 192 = \frac{24}{192} = \frac{1}{8} = \frac{1}{8} \times 100\% = \frac{100}{8} = 12.5 \%$$

C. 4 km to 500metres

( since 4Km= 4000 meters)

$$= 4000:500 = \frac{4000}{500} = \frac{40}{5} = 8$$

$$8 \times 100\% = 800 \%$$

D. 5liters to 4000 milliliters


( since 5liters= 5000 milliliters)

$$= 5000 : 4000$$

$$\frac{5000}{4000} = \frac{5}{4} = \frac{5}{4} \times 100\% = \frac{500}{4} = 125\%$$

### Exercise 3.2.2

- Convert each of the following fractions as percent form?  
A.  $\frac{3}{5}$     B.  $\frac{1}{6}$     C.  $2\frac{3}{4}$     D.  $\frac{13}{6}$
- Express each of the following decimals in percent form.  
A. 0.12    B. 7.5    C. 3.65    D. 0.0012
- Convert each of the following ratios in percent form?  
A. 2m to 5m    C. 50mililiters to 3liters  
B. 2hr to 25minutes    D. 5kg to 5000g
- Express the shaded part in percent form



### Base, rate and percentage



#### Activity 3.2.2

**Discuss** with your friends

- Calculate 10% of 200.
- 5% of 60 is calculated as follows

$$50\% \text{ of } 60 = \frac{5}{100} \times 60 = \frac{300}{100} = 3$$

In this calculation

- A. 5% is called \_\_\_\_\_    C. 3 is called \_\_\_\_\_
- B. 60 is called \_\_\_\_\_

### Definition 3.6:

**Base (B)** is the number that represents 100 % or total value of something.

**Percentage (p)** is the number or the amount that represents a part of the whole.

**Rate (R)** is the ratio of percentage to base, written as percent.

*Note:*

**Base** is usually preceded by “of” in the given statement

**Rate** is expressed in percent

**Example:** 20% of 150 is 30. In this statement determine base, rate and percentage.

**Solution:**

The number expressed in percent is rate, hence 20% is rate.

The whole part or the number preceded by of is base, hence 150 is base. The part of whole is percentage, hence 30 is percentage.

### Formula relating base, rate and percentage

**Example:**

calculate 5% of 120

**Solution:**


$$5\% \text{ of } 120 = \frac{5}{100} \times 120 = 6$$

$\downarrow \quad \quad \downarrow \quad \quad \downarrow$   
 $R \times B = P$

From above example you can see that base, rate and percentage are related by the formula

$$P = R \times B$$

You can also use the following triangle to easily remember the relation between P, R and B.



**Note:**

*R is expressed in percent, so  $\frac{P}{B}$  is multiplied by 100%*

**Example 1:** calculate the following

A. 80% of birr 260

B. 9.6% of 7.2 liters

C. 90% of 264 cm

D. 34% of 7000 Kg

**Solution:**

*A. 80% of birr 260*

$$P = R \times B$$

$$P = \frac{80}{100} \times 260 \text{ Birr}$$

$$p = 80\% \times 260 \text{ Birr}$$

$$P = 8 \times 26 \text{ Birr}$$

$$P = \text{Birr } 208$$

*B. 9.6% of 7.2 liters*

$$P = R \times B$$

$$p = \frac{9.6}{100} \times 7.2 \text{ liters}$$

$$p = 9.6\% \times 7.2 \text{ liters}$$

$$P = 0.0096 \times 7.2 \text{ liters}$$

$$P = 0.6912 \text{ liters}$$

*C. 90% of 264 cm*

$$P = R \times B$$

$$p = 90\% \times 264 \text{ cm}$$

$$P = \frac{90}{100} \times 264 \text{ cm}$$

$$P = 0.9 \times 264 \text{ cm}$$

$$P = 237.6 \text{ cm}$$

*D. 34% of 7000 Kg*

$$P = R \times B$$

$$p = 34\% \times 7000 \text{ kg}$$

$$P = \frac{34}{100} \times 7000 \text{ Kg}$$

$$P = 0.34 \times 7000 \text{ Kg}$$

$$P = 2380 \text{ kg}$$

**Example 2:** Eleni answered 80 % of questions correctly in mathematics test. If the test has 25 questions, then how many questions she answered correctly?

**Solution:**

$$R=80\%, \quad B=25, \quad P=?$$

$$P = R \times B$$

$$P = 80 \% \times 25$$

$$P = \frac{80}{100} \times 25$$

$$\mathbf{P=20}$$

**Therefore,** she answered 20 questions correctly

### Exercise 3.2.3

- Calculate each of the following.
  - What is 10% of 160?
  - What is 60% of Birr 300?
  - Find 4.5% of 9km.
  - Find  $\frac{3}{4}$  % of Birr 5000.
- In a class there are 60 students. If 20% of the class are girls, then how many girls and boys are there?
- 25% of people in Addis Ababa city watched the foot ball final game on TV. How many people watched the football final game if the population of the city is 5000 000

**Example 3:**

calculate the unknown variables in each of the following questions

- |                        |   |
|------------------------|---|
| A. Birr 36 is 9% of x  | C. 16 minutes is $13\frac{1}{3}$ % of T |
| B. 15 cents is 5% of x | D. 90cm is 180% of y                    |

**Solution:**

A.  $R=9\%$  ,  $B= x$  and  $P = 36$

$$B = \frac{P}{R}$$

$$B = \frac{36 \text{ birr}}{9/100} \times 25$$

$$= 36 \times \frac{100}{9} = \frac{3600 \text{ Birr}}{9} = 400 \text{ birr}$$

B.  $R=5\%$ ,  $B=x$  and  $P=15\text{cent}$

$$B = \frac{P}{R} \quad B = \frac{15}{5\%}$$

$$B = \frac{15 \text{ cents}}{5/100} = 15 \text{cents} \times \frac{100}{5}$$

$$B = \frac{1500 \text{cents}}{5} = 300 \text{ cents}$$

C.  $R=13\frac{1}{3}\% = \frac{40}{3}\% = \frac{400}{3}$ ,  $B=T$  and  $P=16\text{minutes}$

$$B = \frac{P}{R}, \quad B = \frac{16 \text{minutes}}{40/300}$$

$$= 16 \text{minutes} \times \frac{300}{40} = 120 \text{minutes}$$

D.  $R=180\%$ ,  $B=y$  and  $P=90\text{cm}$

$$B = \frac{P}{R}, \quad B = \frac{90 \text{cm}}{180\%}$$

$$= 90 \text{cm} \times \frac{100}{80} = \frac{900 \text{ cm}}{80} = 50 \text{ cm}$$

**Example 4:** A women saves 20% of what she earns. If she saves Birr 300 a month, how much does she earn a month?

**Solution:**

$R=20\%$

*A part of money she earns is saved  $P=\text{Birr } 300$*

*The whole amount she earns a month? i.e.  $B=?$*

$$B = \frac{P}{R}, \quad B = \frac{300 \text{birr}}{20\%}$$

$$B = \frac{300 \text{birr}}{20/100} = 300 \text{birr} \times \frac{100}{20} = 1500 \text{birr}$$

**Therefore, the woman earns Birr 1500 a month.**

### Exercise 3.2.4

- Calculate the following
  - 6 is 24% of a number, what is the number?
  - What is the total amount whose 35% is Birr 700.
  - If 15% of a number is 18, then what is this number?
  - 40% of a number is 110, what is number?
- If 30% of a man's salary is Birr 6300, what is the amount of his full salary?
- In a class where the number of boys is 36 % of the total number of students. If there are 18 boys, then how many students are there in the class?

**Example 5:** calculate the following

- What percent of 400 is 50?
- What percent of 2 kg is 400gm?
- What percent of 1 hr is 30 minutes?
- What percent of 60 m is 72cm ?

**Note:**  
 1kg=1000gm  
 1hr=60 minutes  
 1m=100cm

**Solution:**

- A. B=400 ,P= 50 and R = ?

$$R = \frac{P}{B} \times 100\% = R = \frac{50}{400} \times 100\%$$

$$R = \frac{50}{400} \times 100\% = \frac{50}{4}\% = 12.5\%$$

- B. B = 2Kg ,P =400gm and R= ?

$$R = \frac{P}{B} \times 100\% = R = \frac{400 \text{ gm}}{2000 \text{ gm}} \times 100\%$$

$$= \frac{400}{20}\% = 20\%$$

- C. B= 1hr = 60 minutes, P = 30 minutes, R= ?

$$R = \frac{P}{B} \times 100\% = R = \frac{30 \text{ minutes}}{60 \text{ minutes}} \times 100\%$$

$$= \frac{300}{6}\% = 50\%$$

$$D. B = 60 \text{ m} = 6000\text{cm}, \quad P = 72\text{cm} \quad \text{and} \quad R = ?$$

$$R = \frac{P}{B} \times 100\% = R = \frac{72\cancel{\text{cm}}}{6000\cancel{\text{cm}}} \times 100\%$$

$$= \frac{72}{60} \% = 1.2\%$$

**Example 6:** In a certain village where farmers use two different types of fertilizers, fertilizer type A and fertilizer type B. 10000 farmers were asked which of the fertilizers they use. It was found that 2500 farmers use type A, 4550 farmers use type B and the remaining farmers use both types A and B. Find the percent of the farmers that

- Use fertilizer type A
- Use fertilizer type B
- use both types of fertilizers

**Solution:**

*A. The total 10000 farmers is base, B = 10000*

*2500 is part of total farmers that use type A, hence it is percentage,*

$$P = 2500 \quad R = ?$$

$$R = \frac{P}{B} \times 100\% =$$

$$R = \frac{2500}{10000} \times 100\% = 25\%$$

*Therefore, 25% of farmers use type A fertilizer*

*B. Total 10000 farmers is base, B = 10000*

*4550 is part of total farmers that use type B, hence it is percentage,*

$$P = 4550 \quad R = ?$$

$$R = \frac{P}{B} \times 100\%$$

$$R = \frac{4550}{10000} \times 100\% = \frac{455}{100}\% = 45.5\%$$

*Therefore, 45.5% of farmers use type B fertilizer*

C. Total 10000 farmers is base,  $B = 10000$  the remaining farmers that use both type of fertilizers is  $P = 10000 - (2500 + 4550) = 2950$ ,

$$P = 2950 \quad R = ?$$

$$R = \frac{P}{B} \times 100\%$$

$$R = \frac{2950}{10000} \times 100\% = \frac{295}{10}\% = 29.5\%$$

Therefore, 29.5% of farmers use both type of fertilizer.

### Exercise 3.2.5

- Evaluate the following
  - What percent of 60 is 30?
  - What percent of 24 is 6?
  - What percent of 800 is 500?
- A Woman saves Br.300 from her monthly salary. If her monthly salary is Br.7500, then find her saving in percent?
- The following table gives the number of students in a given class according to their age and sex.

Age	Sex	
	Male	Female
13	15	18
14	6	9
Total	21	27

- What percent of the class are females?
  - What percent of the class are males?
  - What percent of class are 13 years old?
  - What percent of class are 14 years old female students ?
- In a class of 48 students 6 of them were absent on Monday. What percent of the class was absent and what percent of class was attended on that day?

### **Mixed problems containing base, rate and percentage**

1. A student scored 16 out of 25 in mathematics mid exam. What is the student's score in percent?
2. There are 50 students in a class. If 8% are absent on a particular day, find the number of students present in the class.
3. In a basket of oranges, 20% of them are defective and 76 are in good condition. Find the total number of oranges in the basket.
4. Hawa was able to sell 220 kilograms of her vegetables before noon. If Hawa had 400 kg of vegetables in the morning, what percent of Kilograms of vegetables was she sold in the morning?
5. A cow gives 24 liters milk each day. If the milkman sells 75% of the milk in one day, how many liters of milk is left with him?
6. Tolosa sold 540 eggs. If these are 36% of total eggs, then how many eggs are not sold?
7. A factory has 2400 workers of which 900 are male and the rest are female. What percent of the workers are female?

### **3.3. APPLICATION OF RATIO, PROPORTION AND PERCENTAGE**

**At the end of this section, learners will be able to:**

⇒ Apply the concept of percentage to solve real life problems.

In our day to day life, we come across a number of situations wherein we use the concept of percent. In the following section, we discuss the application of percentage in different fields like problems in percent change, profit and loss, discount, simple interest, compound interest, Value added tax (VAT), turn of tax (TOT), and income tax.

**Percent increase and decrease****Activity 3.3.1****Discuss with your friends**

1. A quantity increases from 400 to 600
  - A. Find the increased amount
  - B. Find the ratio of the increased amount to original quantity
  - C. Express the answer in  $b$  in the form of percent (find the percent increase)
  - D. How do you calculate percent increase?
2. A quantity decreases from 500 to 300
  - A. Find the decreased amount
  - B. Find the ratio of decreased amount to original quantity
  - C. Express the answer in  $b$  in the form of percent (find the percent decrease)
  - D. How do you calculate percent decrease?

**Note:**

1. Percent increase is calculated as:

$$\text{percent increase} = \frac{\text{increased amount}}{\text{original quantity}} \times 100\%$$

2. Percent decrease is calculated as:

$$\text{percent decrease} = \frac{\text{decreased amount}}{\text{original quantity}} \times 100\%$$

**Example 1:** Find the percent increase from 250 to 400.

**Solution:**

$$\text{Increased amount is } 400 - 250 = 150$$

*Original quantity is 250*

$$\text{percent increase} = \frac{\text{increased amount}}{\text{original quantity}} \times 100\%$$

$$\text{percent increase} = \frac{150}{250} \times 100\% = 60\%$$

**Therefore,** the percent increase from 250 to 400 is 60%

**Example 2:** Find the percent decrease from 400 to 250.

**Solution:**

*decreased amount is  $400 - 250 = 150$*

*Original quantity is 400*

$$\text{percent decrease} = \frac{\text{decreased amount}}{\text{original quantity}} \times 100\%$$

$$\text{percent decrease} = \frac{150}{400} \times 100\% = 37.5\%$$

**Therefore,** the percent decrease from 250 to 400 is 37.5%

**Example 3:** The student's population of a school is 4000 in 2013 E.C.

If the student's population increased to 4620 in 2014 E.C then, find the percent increase between the two years.

**Solution:**

*Increased amount is  $4620 - 4000 = 620$*

*Increased amount is 620*

$$\text{percent increase} = \frac{\text{increased amount}}{\text{original quantity}} \times 100\%$$

$$\text{percent increase} = \frac{620}{4000} \times 100\% = 15.5\%$$

**Example 4:** 20% of an article is damaged and thrown away and only 20kg is left. Find its original weight.

**Solution:**

let the original weight is  $x$ . The weight of the article decreased from  $x$  to 20 Kg. Percent decrease is 20%

$$\text{percent decrease} = \frac{\text{decreased amount}}{\text{original quantity}} \times 100\%$$

$$20\% = \frac{x - 20}{x} \quad \text{no need of multiplying by } 100\%, \text{ percent decrease is given.}$$

$$\frac{20}{100} = \frac{x - 20}{x}, \quad 20x = 100(x - 20) = 20x = 100x - 2000$$

$$20x - 100x = -2000 = \frac{-80x}{-80} = \frac{-2000}{-80}$$

$$x = 25\% \text{ Kg}$$

**Therefore,** the original weight is 25% Kg

### Exercise 3.3.1

- Find the percent change
  - From 80 to 100
  - From 800 to 500
  - From 500 to 300
  - From 6000 to 9000
- While measuring a line segment 5cm, it was measured 5.2cm by mistake. Find the percent error in measuring this line segment
- The number of student fail in mathematics test decreased from 20 to 12. What is the percent decrease?
- Last year Samuel's salary was Birr 8000. If he gets a 10% increment this year, what is his current salary?
- A T – shirt, discounted by 30% for a clearance sale. If the discounted price is Birr 399, then calculate
  - The original price of T – shirt
  - The amount of discount

### 3.3.1 Calculating profit and loss percentage



#### Activity 3.3.2

#### Discuss with your friends

1. A shop keeper buys a pair of shoes for Birr 500 and sells it for Birr 600. Find:-  
 A. The cost price                      C. Profit  
 B. The selling price                  D. Profit percent
2. The price at which an article is purchased is called \_\_\_\_\_.
3. The price at which an article is sold is called \_\_\_\_\_.
4. Profit is made, when selling price is \_\_\_\_\_ than cost price.
5. Loss is made, when selling price is \_\_\_\_\_ than cost price.

#### Definition:

**Cost Price (C.P):** *The Price, at which an article is purchased, is called its cost price.*

**Selling Price (S.P):** *The Price, at which an article is sold, is called its selling price.*

**Profit (Gain):** *When, then there is profit, and When  $S.P > C.P$ , then there is profit, and*

$$\text{Profit} = S.P - C.P$$

**Loss:** *When  $S.P < C.P$ , then there is loss, and*

$$\text{Loss} = C.P - S.P$$

## The formula to calculate profit and loss percent

### 1. Profit percent:

$$\text{profit \%} = \frac{\text{profit}}{\text{cost price}} \times 100\%$$

$$\text{profit \%} = \frac{\text{selling price} - \text{cost price}}{\text{cost price}} \times 100\%$$

$$\text{Profit \%} = \frac{S.P. - C.P}{C.P} \times 100\%$$

### 2. loss percent:

$$\text{Loss \%} = \frac{\text{Loss}}{\text{cost price}} \times 100\%$$

$$\text{Loss \%} = \frac{\text{Cost price} - \text{Selling price}}{\text{cost price}} \times 100\%$$

$$\text{Loss \%} = \frac{C.P. - S.P}{C.P} \times 100\%$$

**Example 1:** A shop keeper buys a jacket for Birr 500, and gives it clean, then sells it for Birr 640. Calculate his profit percent.

*Solution:*

$$C.P = 500, \quad S.P = 640, \quad \text{profit \%} = ?$$

$$\text{Profit \%} = \frac{S.P - C.P}{C.P} \times 100\%$$

$$\text{Profit \%} = \frac{640 - 500}{500} \times 100\%$$

$$\frac{140}{500} \times 100\% = 28\%$$

**Therefore,** his profit percent is 28 %.

**Example 2:** Girma bought 200 eggs for Birr 8 each. 20 eggs are damaged and sold the remaining eggs at Birr 8.5 each; calculate his profit/loss percent.

**Solution:**

The cost of 1 egg is Birr8, so the cost of 200 eggs is  $200 \times 8 = \text{birr } 1600$   
 $= \text{C.P}$  20 eggs are damaged, so  $200 - 20 = 180$  eggs are sold at  
 Birr 8.5 each.

$$SP = 180 \times 8.5 = \text{Birr } 1530.$$

Since,  $S.P. < C.P.$ , there is loss and his loss percent is 100%

$$\text{Loss \%} = \frac{C.P - S.P}{C.P}$$

$$\text{Profit \%} = \frac{1600 - 1530}{1530} \times 100\%$$

$$\frac{70}{16} \times 100\%$$

$$= 4.375\%$$

Therefore, his loss percent is 4.375%.

**Example 3:** A dealer gained 10% profit by selling an article for birr 330. What was the original price of the article?

**Solution:**

profit % = 10%, S.P = 330, C.P=?

$$\text{Profit \%} = \frac{S.P - C.P}{C.P} \times 100\%$$

$$10\% = \frac{330 - x}{x}$$

$$\frac{10}{100} = \frac{330 - x}{x}$$

no need of multiplying by  
 100%, profit % is given and  
 $C.P = x$

$$10x = 100(330 - x)$$

$$10x = 33000 - 100x$$

$$10x + 100x = 33000$$

$$x = 300$$

Therefore, the original price of the article is Birr 300

### Exercise 3.3.2

1. Find profit or Loss percent when:
  - A. Cost Price = Birr 2500 and Selling Price = Birr 2000
  - B. Cost Price = Birr 3500 and Selling Price = Birr 4500
2. Find cost price when:
  - A. Selling Price = Birr 7950 and Gain = 6%
  - B. Selling Price = Birr 980 and Loss = 12%
3. Find selling price when
  - A. Cost Price = Birr 870 and Gain = 10%
  - B. Cost Price = Birr 750 and Loss = 15%
4. An article was bought for Birr 2000 and sold by Birr 2200. Find the profit or loss percent.
5. A shop keeper bought a jacket for Birr 1500 and gives it clean, then sold it for Birr 1800. What is his profit percent?
6. Estifanos bought goods by Birr 5200 and sold it at a profit of 12%. Find the selling price.
7. By selling a camera for Birr 24000, a man loss 4%. Find the cost price?
8. A trader bought a TV set for Birr 2500 and sold it at a loss of 8%. what was the selling price?
9. A man bought 100 eggs for Birr 800 and sells them for Birr 10 each, then find his profit percent.

### 3.3.2. Simple interest



#### Activity 3.3.3

#### Discuss with your friends

1. If you deposit Birr 1000 in a bank, then
  - A. After a year Birr 1000 increased by a certain amount, why?






B. By what amount Birr 1000 increases within 1 year, if the bank pays you 10% per year for using your money?

C. By what amount Birr 1000 increases within 2 years, if a bank pays you 10% per year on money you deposited initially?

2. Define the following by your own words

A. Simple interest      B. Interest

A person needs to borrow some money as a loan from credit & saving association bank, his friends, relatives etc. He promises to return it after a specified time period along with some extra money for using the money of lender.

-  The amount of money borrowed or invested is called **principal (p)**.
-  The extra money paid for the use of money is called **interest (I)**.
-  The time length for which money is borrowed or lent is called **Time (T)**.
-  The total amount of money including interest at the end of time period is called **Amount (A)**.
-  The percentage charged on the total amount you borrow or save is called **interest rate(R)**.

**Example 1:** If Birr 2000 is deposited in a bank at a rate of 7% per year, for 1 year then,

- A. What is the principal?      C. What is the time period?  
B. What is the interest rate?      D. Calculate interest paid in one year?

**Solution:**

Money deposited is Birr 2000, so, principal(P) is Birr 2000

The interest rate(R) is 7% per year


The time period(T) is 1 year

The interest(I) in one year is 7% of Birr 2000, so  $I = 7\% \times 2000$

$$I = \frac{7}{100} \times 2000 = \text{Birr } 140$$

The interest paid on original principal only during the whole interest period is called **simple interest**.

**Simple interest is calculated by the formula:**

<p><b><math>I = PRT</math></b></p> <p>Where,</p> <p><math>I</math> = simple interest  <math>P</math> = principal  <math>T</math> = Time</p> <p>And amount (<math>A</math>) is given by:  <math>A = P + I</math></p>	<p>You can use the following triangle to easily remember the relation between <b><math>I, P, R, T</math></b>.</p> <div style="text-align: center;">  </div> <p> <math>P = \frac{I}{RT}</math>  <math>I = PRT</math>  <math>T = \frac{I}{PR}</math> and  <math>R = \frac{I}{PT} \times 100\%</math> </p>
---	--

- Example 2:** If Birr 4000 is deposited in a bank at a rate of 7% per year, for 4 years then, calculate
- Simple interest at the end of 4 years
  - Amount at the end of 4 years

**Solution:**

$$P = \text{Birr } 4000, \quad R = 7\% \text{ per year}, \quad T = 4 \text{ year}$$

A.

$$I = PRT$$

$$I = 4000 \text{ Birr} \times \frac{7}{100} \times 4$$

$$I = \text{Birr } 1,120$$

B.

$$A = P + I$$

$$A = 4000 + 1120$$

$$A = \text{Birr } 5120$$

**Example 3:** Find the simple interest on Birr 700 at 12 % per year for 3 months.

**Solution:**

$P = \text{Birr}700$  ,  $R = 12\%$  Per year,  $T = 3 \text{ Months}$ , since the interest rate is per year, the time must be converted to years.

$$T = 3 \text{ Months} = \frac{3}{12} \text{ Year} = \frac{1}{4} \text{ Year}$$

$$I = PRT$$

$$I = \text{Birr } 700 \times \frac{12}{100} \times \frac{1}{4}$$

$$I = \text{Birr } 21$$

**Example 4:** If Birr 1000 grows to 1200 in 4 years at simple interest rate, then calculate the simple interest rate.

**Solution:**  $P = \text{Birr}1000$  ,  $R = ?$ ,  $T = 4 \text{ Year}$

$$I = PRT \text{ so } R = \frac{I}{PT} \times 100\%$$

$$\frac{200}{1000 \times 4} \times 100\% = 5\% \text{ per year}$$

### Exercise 3.3.3

1. Michael invests Birr 2,000 in the bank that pays simple interest rate of 5% per year for 6 years. Then how much interest will he get in 6 years?
2. Birr 24,000 invested at 11% simple interest per annum, then what is the amount after 8 years?
3. How long will it take for Birr 15,000 to double itself, if it is invested at simple interest rate of 10% per year?
4. If Birr 20,000 grows to Birr 28,000 after 20 years. Then what is the simple interest rate?

5. In how many years will Birr 12, 000 yield an interest of Birr 1,200 at 5% simple interest?
6. An investment earned Birr 2,100 interest after 6 years. If the simple interest rate is 7% per year. What was the principal?

### 3.3.3. Compound interest



#### Activity 3.3.4

**Discuss with your friends**

1. If Birr 1,000 is invested at a rate of 10% per year, then calculate
  - A. The interest at the end of first year
  - B. The interest on the second year. (take second year principal to be initial principal plus first year's interest)
  - C. The interest on third year. (take third year principal to be initial principal plus first year interest plus second year interest)
  - D. The interest in 3 years
2. What is compound interest?

**Compound interest** is the interest on a loan calculated based on the initial principal plus the accumulated interest from previous periods

**Example 1:** If Birr 2000 is invested at a rate of 10 % compounded annually for 3 years, then calculate

- A. Compound interest for 3 years
- B. Amount at the end of third year

**Solution:**

A. Compound interest for 3 years Is

**Interest for first year ( $I_1$ )**

$$P_1 = 2000, \quad R = 10\%, \quad T = 1 \rightarrow \text{year for year one only}$$

$$I_1 = P_1RT,$$

$$\text{so } I_1 = 2,000 \times 10\% \times 1$$

$$I_1 = 2000 \times \frac{10}{100} \times 1$$

$$I_1 = \text{Birr } 200$$

**Interest for second (Year-2) ( $I_2$ )**

Principal for second year is initial principal plus interests on previous years.

$$P_2 = P_1 + I_1$$

$$P_2 = 2,000 + 200 = P_2 = 2,200$$

$$P_2 = 2,200, \quad R = 10\%, \quad T = 1 \text{ year} \rightarrow \text{for year two only}$$

$$I_2 = P_2RT$$

$$\text{so } I_2 = 2,200 \times 10\% \times 1$$

$$I_2 = 2200 \times \frac{10}{100} \times 1$$

$$I_2 = \text{Birr } 220$$

**Interest for third (Year-3) ( $I_3$ )**

Principal for third year is initial principal plus interests on previous years.

$$P_3 = P_1 + I_1 + I_2$$

$$P_3 = 2,000 + 200 + 220 = P_3 = 2,420$$

$$P_3 = 2,420, \quad R = 10\%, \quad T = 1 \text{ year} \quad \text{for year three only}$$

$$I_3 = P_3RT$$

$$\text{so } I_3 = 2,420 \times 10\% \times 1$$

$$I_3 = 2420 \times \frac{10}{100} \times 1$$

$$I_3 = \text{Birr } 242$$

**Therefore, the interest for 3 years is**

$$I = I_1 + I_2 + I_3$$

$$I = 200 + 220 + 242$$

$$I = 662$$

**B. Amount at the end of third year**

$$A = P + I$$

$$A = 2,000 + 662$$

$$A = 2,662$$

**Note:**

*For compound interest amount can be calculated by the formula of*

$$A = P (1 + R)^t$$

Where,  $A =$  amount,

$R =$  interest rate,  $P =$  principal and  $T =$  time

And  $I = A - P$

Using the formula, the above example is done as follows

$$P = 2000, \quad R = 10\%, \quad t = 3 \text{ Years}, \quad A = ?, \quad I = ?$$

$$A = P (1 + R)^t$$

$$A = 2000(1 + 10\%)^3$$

$$A = 2000(1 + 0.1)^3$$

$$= 2000 \times 1.1^3$$

$$= 2000 \times 1.331$$

$$A = \text{Birr } 2662$$

*interest,*  $I = A - P$

$$I = 2662 - 2000 = \text{Birr } 662$$

**Example 2:** what sum of money amounts to Birr 24,200 in 2 years at 10% per annum, compounded annually?

**Solution:**

$$P = ?, \quad R = 10\%, \quad t = 2 \text{ Years}, \quad A = \text{Birr } 24,200,$$

$$A = P(1 + R)^t$$

$$24,000 = P(1 + 10\%)^2$$

$$24,000 = P(1 + 0.1)^2$$

$$24,000 = 1.21 \times P$$

$$\frac{1.21p}{1.21} = \frac{24200}{1.21} \quad p = 20,000$$

### Exercise 3.3.4

1. Find the compound interest on Birr 8000 for 2 years at 5% per annum, compounded annually.
2. Compare the simple interest and compound interest on Birr 8,000 at 10 % per annum for three years if the interest is compounded annually.
3. Find the difference between the simple and the compound interest on Birr 5000 for 2 years at 6% per annum.
4. Robel obtained a loan of Birr 250,000 from the commercial bank of Ethiopia. If the rate of interest is 8% per annum compounded annually and repaid Birr 50, 000 at the end of first year. What amount will he has to pay to the bank after 2 years to discharge his debt?
5. What sum of money will amount to Birr 21296 in 3 years at 10% per annum, compounded annually?

### 3.3.4 Ethiopian Income Tax, Turn over Tax, VAT

#### Activity 3.3.5

#### **Discuss with your friends**

1. Why do governments collect tax?
2. List out the different types of tax.

**Taxes** are imposed by governments on their citizens to generate income for undertaking projects to boost the economy of the country. Tax raises the revenue of the government to cover cost of

- |                  |             |
|------------------|-------------|
| ✓ Administration | ✓ Education |
| ✓ Defense        | ✓ Housing   |
| ✓ Health etc.    |             |

**We should pay tax!**

To boost our

**country's economy**

Taxes help to raise the standard of living in a country.

In this section we will try to see three different types of tax, VAT, turn over tax and income tax.

#### **Value added tax (VAT)**

**Value added tax (VAT)** is a tax imposed by government on sales of some goods and services.

#### **Note:**

1. In Ethiopia VAT rate is 15%
2.  $VAT \text{ amount} = 15\% \text{ Original Cost}$
3.  $\text{Cost including VAT} = \text{Original Cost} + VAT \text{ amount}$

#### **Example**

**1:** the price of a machine is Birr 3000 before VAT.

A. Calculate the amount of VAT

B. Calculate the total cost of machine including VAT

**Solution:**

A.  $VAT = 15\%$  of Original Cost.

$$= \frac{15}{100} \times 3000 = \text{Birr } 450$$

$$\begin{aligned} \text{B. total cost of machine} &= 3000 + 450 \\ &= \text{Birr } 3450 \end{aligned}$$

**Example 2:** The cost of TV stand including VAT is Birr 2300.

A. Calculate the cost of TV stand before VAT

B. Calculate the amount of VAT

**Solution:** let the cost before VAT is  $x$

A. *cost including VAT = cost before VAT + VAT amount*

$$2300 = x + 15\% \text{ of } x$$

$$2300 = x + 0.15x$$

$$2300 = 1.15x$$

$$x = \frac{2300}{1.15} = \text{Birr } 2000$$

Note

$$\text{cost before VAT} = \frac{\text{cost including VAT}}{1.15}$$

*Therefore, the cost before VAT is Birr 2000*

B. the amount of VAT

$$VAT \text{ amount} = 15\% \text{ of original cost}$$

$$= \frac{15}{100} \times 2000 = \text{Birr } 300$$

### Turn over Tax (TOT)

*Turn over tax is imposed on merchants, who are not required to register for VAT, but supply goods and services in the country.*

In Ethiopia turn over tax rate is on goods sold and services rendered locally.

**Note:**

1. A merchant his annual income below Birr 500,000 will be registered to collect turn over tax.
2. A merchant whose annual income above Birr 500,000 will be registered to collect turn over tax

**Example 1:** calculate turn over tax on sales of Birr 10,000

**Solution:** Turn over tax = 2% of 10,000

$$= \frac{2}{100} \times 10,000 = \text{Birr } 200$$

**Example 2:** Elsa stationery has daily sales of Birr 205. The stationery is open 250 days per year. How much is the turn over tax payable by Elsa?

**Solution:** sales per year =  $205 \times 250 = \text{Birr } 51,250$

Turn over tax = 2% of sales

$$= \frac{2}{100} \times 51,250 = \text{Birr } 1,025$$

## Employment income tax

Employer deduct income tax from the employee before paying monthly salary based on the following tax rate (according to Ethiopian income tax rate)

Employment income (per month) in Birr	Employment in- come Tax rate
0 - 600	0%
601-1650	10%
1651-3200	15%
3201-5250	20%

5251-7800	25%
7801-10900	30%
over 10,900	35%

**Note:**

*the above table shows the tax rate for each interval of total income*

**Example1:** the income tax for Birr 850 is calculated as follows

**Solution:** Birr 850 falls on 10%

For interval 0 to 600 , the tax rate is 0%, hence  $Tax_1 = 0$

For interval 601 to 850 , the tax rate is 10%, hence tax on this interval is  $Tax_2 = 10\% \text{ of } (850-600)$

$$Tax_2 = \frac{10}{100} \times 250$$

$$Tax_2 = 25$$

$$\text{Income tax} = Tax_1 + Tax_2$$

$$= 0 + 25 = \text{Birr } 25$$

**Note:**

for interval for 601 to 850, any value above 600 up to 850 is taken, so The difference is 850-600. It is not 850 - 601

**Therefore,** the income tax on Birr 850 is 25

**Example 2:** Alemu earns Birr 4850 per month, calculate

- The income tax he has to pay.
- The net income after deducting income tax

**Solution:**

**A. Birr 4850 falls on 20%**

➤ For interval 0 to 600 , the tax rate is 0%, hence  $Tax_1 = 0$

➤ For interval 601 to 1650,, the tax rate is , hence tax on this interval is  $Tax_2 = 10\% \text{ of } (1650-600)$

$$Tax_2 = \frac{10}{100} \times 1050 = \text{Birr } 105$$

- For interval 1651 to 3200, the tax rate is 15%, hence tax on this interval is  $Tax_3 = 15\%$  of  $(3200 - 1650)$

$$Tax_3 = \frac{15}{100} \times 1550 = \text{Birr } 232.5$$

- For interval 3201 to 4850, the tax rate is 20%, hence tax on this interval is  $Tax_4 = 20\%$  of  $(4850 - 3200)$

$$Tax_4 = \frac{20}{100} \times 1650 = \text{Birr } 330$$

$$\begin{aligned} \text{Income tax} &= Tax_1 + Tax_2 + Tax_3 + Tax_4 \\ &= 0 + 105 + 232.5 + 330 = \text{Birr } 667.5 \end{aligned}$$

**Therefore,** the income tax on Birr 4850 is Birr 667.5

B. net income is  $4850 - 667.5 = \text{Birr } 4182.5$

### Example 3:

*Derive a formula to calculate income tax and on Birr  $x$ , in terms of  $x$ . Where  $x$  falls on the interval 1651 to 3200*

**Solution:** Birr  $x$  falls on 15%

For interval 0 to 600, the tax rate is 0%, hence  $Tax_1 = 0$

For interval 601 to 1650, the tax rate is 10%, hence tax on this interval is  $Tax_2 = 10\%$  of  $(1650 - 600)$

$$Tax_2 = \frac{10}{100} \times 1050 = \text{Birr } 105$$

For interval 1651 to  $x$ , the tax rate is 15%, hence tax on this interval is  $Tax_3 = 15\%$  of  $(x - 1650)$

$$Tax_3 = 0.15x - 247.5$$

**Therefore,** the income tax

$$Tax = Tax_1 + Tax_2 + Tax_3 = 0 + 105 + 0.15x - 247.5$$

$$Tax = 0.15x - 142.5$$

### Exercise 3.3.5

- Find the income tax and net income of the following employees of commercial Bank of Ethiopia
  - Ato Ahmed with monthly salary of Birr 7,500
  - w/ro Mekdes with monthly salary Birr 11,600
- A shoe dealer purchased shoe from a shoe company that worth Birr 8000. Find the amount he should pay to the company including VAT

### SUMMARY FOR UNIT 3

- The method of comparing two or more quantities of the same kind and in the same units by division is known as a ratio.
- Proportion is equality of two ratios.
- In proportion the product of means is equal to the product of extremes. (i.e. if  $\frac{a}{b} = \frac{c}{d}$ , then  $a \times d = b \times c$ )
- y is said to be directly proportional to (written as  $y \propto x$ ), if there is constant k such that  $y = kx$  or  $k = \frac{y}{x}$
- y is said to be inversely proportional to (written as  $y \propto \frac{1}{x}$ ), if there is constant k such that  $y = \frac{k}{x}$  or  $k = y \times x$ .
- The word percent means “for every hundred” or “per 100”. We use the symbol % to denote percent.
- **Base (B)** is the number that represents 100% or total value of something.
- **Percentage (p)** is the number or the amount that represents a part of the whole.
- **Rate(R)** is the ratio of percentage to base, written as percent.
- **Percentage, base and rate are related by the formula,**  

$$P = R \times B, \quad B = \frac{P}{R} \text{ and } R = \frac{P}{B}$$

➤ Percent increase is calculated as:

$$\text{percent increase} = \frac{\text{increased amount}}{\text{original quantity}} \times 100\%$$

➤ Percent decrease is calculated as:

$$\text{percent decrease} = \frac{\text{decreased amount}}{\text{original quantity}} \times 100\%$$

➤ The Price at which an article is purchased, is called its **cost price**.

➤ The Price at which an article is sold, is called its **selling price**.

➤ When  $S.P > C.P$ , then there is profit, and Profit =  $S.P - C.P$

➤ When  $S.P < C.P$ , then there is loss, and loss =  $C.P - S.P$

➤ Profit or loss percent is given by the formula

$$\text{profit \%} = \frac{\text{profit}}{\text{cost price}} \times 100\% \quad \text{and} \quad \text{Loss \%} = \frac{\text{Loss}}{\text{cost price}} \times 100\%$$

➤ The interest paid on original principal only during the whole interest period is called **simple interest**.

$$\text{Interest} = \text{principal} \times \text{Rate} \times \text{Time} \quad \text{Or} \quad I = PRT$$

➤ **Compound interest** is the interest on a loan calculated based on the initial principal plus the accumulated interest from previous **periods**.

$$A = P(1+R)^T \quad \text{and} \quad I = A - P$$

➤ **Value added tax (VAT)** is a tax imposed by government on sales of some goods and services.

➤ In Ethiopia VAT rate is 15%

➤ Turn over tax is imposed on merchant, who are not required to register for VAT, but supply goods and services in the country.

➤ In Ethiopia turn over tax rate is 2% on goods sold and services rendered locally.

### Review exercise for unit 3

#### I. Write True for the correct statements and False for the incorrect statements.

- Ratio can be defined as “for every hundred terms”.
- $4:3x=32:72$  if  $x=3$
- The ratio of 15 meters to 2000 centimeters is 3:4
- $Loss \% = \frac{Loss}{cost\ price} \times 100\%$
- If 250 divided in the ratio 3:7, the smaller part is 25.
- Proportion is the equality of two ratios.
- If  $y$  is inversely proportional to  $x$ , then as  $x$  increases  $y$  also increases.
- Profit is made when the selling price is less than the cost price

#### II. Choose the correct answer from the given alternatives.

- In a school, 58% of the total number of students is boys. If the Number of girls is 840, how many students are there in the class?  
A. 2000      B. 1800      C. 1500      D. 2100
- Birr 300 is invested at 6% simple interest per annum. How long will it take for the interest to be Birr 180?  
A. 12 years      B. 10 years      C. 8 years      D. 24 years
- The decimal form of  $\frac{3}{4}\%$  is  
A. 0.75      B. 0.075      C. 0.0075      D. 0.057
- The simple interest on Birr 400 invested for 4 months was Birr 12. What is the annual (yearly) rate of interest?  
A. 36      B. 4.8      C. 7      D. 9
- If  $a$ ,  $b$ , and  $c$  are numbers such that  $a:b:c=1:3:4$  and  $c=28$  then, find the sum of  $a+b+c$ ?

- A. 56                      B. 60                      C. 72                      D. 84

14. If  $A:B = 7:8$  and  $B:C = 12:7$ , find  $A: C$  in its simplest form.

- A. 2:3                      B. 3:2                      C. 19:6                      D. 4:5

15. Three numbers  $m, n, p$  are in the ratio  $3:6:4$ . Which of the following is the value of  $\frac{4m-n}{n+2p}$

- A.  $\frac{7}{3}$                       B.  $\frac{3}{7}$                       C.  $\frac{2}{3}$                       D.  $\frac{4}{3}$

16. The ratio of the number of pigs to the number of horses in a farm is 2 to 3. If there are 24 horses, what is the number of pigs?

- A. 72                      B. 48                      C. 36                      D. 16

17. 24,  $x$ , 18 and  $(x-1)$  are in proportion. Find the value of  $x$ .

- A. 6                      B. 4                      C. 12                      D. 8

18. If 85% of 'a' is 'b', then which of the following is true?

- A.  $a=b$                       B.  $a < b$                       C.  $20a = 17b$                       D.  $a > b$

19. which one of the following is not equal to the rest?

- A. 2% of 150      B.  $\frac{3}{2}\%$  of 400                      C. 5% of 60                      D. 6% of 50

20. Which one of the following greater?

- A. 20% of 45      B. 25% of 60                      C. 2% of 800                      D. 74% of 20

### III. Work out problems

21. Find  $\frac{7}{12}$  of 8400?

22. There are 40 questions on a final exam and Yohannes answered 75% of them correctly. How many questions correctly answered?

23. If 'b' is directly proportional to 'a' and  $b=30$  when  $a=20$ , then

- A. Find the equation relating  $a$  and  $b$                       B. Find  $a$  when  $b=42$

24. In a given class room, the number of girls is 10 greater than the number of boys. If the ratio of the number of girls to the number of boys is 7:5 then find

- A. The number of girls                      C. The total number of students

- B. The number of boys
25. A company has 420 workers of which 189 are males and the rest are females. What percent of the workers are female?
26. A trader bought a TV set for Birr 22000 and sold it at loss of 5%. What was the selling price?
27. Mussa received a 6% pay raise. He now earns Birr 8268, what was his salary prior to the salary increase?
28. Birr 6000 is invested at a rate of 5% compound interest compounded annually, Find a. the amount at the end of 2 years b. The interest at the end of 2 years
29. A person wants to buy a car from TOYOTA Company. If the price car including VAT is Birr 5,750,000 then,
- A. What is the price of car before VAT?
- B. What is the value of VAT?
30. Calculate the income tax on salary of Birr 5200.  
Derive a formula to calculate income tax and on Birr  $x$ , in terms of  $x$ . Where  $x$  falls on the interval 5251 to 7800

# UNIT 4

## LINEAR EQUATIONS

### *Learning Outcomes:*

At the end of this unit, learners will be able to:

- ↪ identify variables, terms and variables in algebraic expressions
- ↪ simplify algebraic expressions
- ↪ develop their skills on rearranging and solving linear equations
- ↪ apply the rules of transformation of linear equations for solving problems
- ↪ draw a line through the origin whose equation is given.
- ↪ apply real-life situations in solving linear equations

### *Introduction*

In this unit you will learn the importance of variables in mathematics, apply variables in solving linear **equations**, like linear **equations** with **one** variable involving brackets and linear equation involving fractions.

The unit has four sections. The first section deals with the use of variables in formula and the second section deals with solving linear equation, the third section deals with Cartesian coordinate system and the last section deals about application of linear equations. Here the student will see how to use variables in formula and how to solve linear equation with one variable involving brackets and fractions.

## 4.1. ALGEBRAIC TERMS AND EXPRESSIONS

### *Learning Outcomes:*

At the end of this unit, learners will be able to:

- Describe algebraic terms and expressions
- Simplify algebraic expressions with and without brackets

### 4.1.1. USE OF VARIABLES IN FORMULA



#### Activity 4.1.1

1. Identify the variables

23,  $x$ , 4,  $y$ ,  $z$ , 0.5,  $3x$ ,  $\frac{3}{4}$ ,  $a$ ,  $3b\frac{1}{2}$


2. What is variable?

3. Express the following using variables.

- A. A number  $x$  plus 4
- B. The difference 5 and  $y$ , where  $y$  is greater than 5.
- C. The product of 3 and  $x$ .
- D. I start with  $x$ , add 8 and then, double the result.

4. From the below figure

- A. what is the length of AB
- B. what is the length of BC
- C. Find the perimeter of the figure in terms of  $x$  and  $y$ .



### Definition 4.1:

*a variable is any letter or a symbol that represent some unknown number or value. Such as  $x, y, z, n, m \dots$*

**Example1:** In  $x+2$ ,  $x$  is variable

**Example2:** Describe the following using variables

- |   |                                    |
|---|------------------------------------|
| A. Twice a number.                        | B. Three times a number.           |
| C. Two more than a number.                | D. Half a number.                  |
| E. Two less than a number.                | F. The product of a number and 10. |
| G. Product of ten and sum of two numbers. |                                    |

**Solution:**

- A. let the number is  $x$ , then twice the number  $= 2x$ .
- B. let the number is  $z$ , then three times a number  $= 3z$ .
- C. let the number  $y$ . then two more than a number  $= y+2$ .
- D. let the number is  $n$ , then half the number  $= \frac{n}{2}$ .
- E. let the number is  $m$ , then two less than a number  $= m-2$
- F. let the number is  $k$ , then product of a number and 10  $= 10k$
- G. let the two numbers are  $x$  and  $y$ , then product of ten and sum of two number  $= 10(x+y)$

**Example 3:** If you are 13 years old now. How old will/were you

- A. after  $x$  years ?      B. after  $z$  years?      C.  $y$  years ago?

**Solution:**

A.  $13+x$

B.  $13+z$

C.  $13-y$

### Exercise 4.1.1

**1. Identify** the variables in each of the following.

A.  $2+4x$

B.  $5x+x$

C.  $2y+b$

D.  $\frac{x}{2}-p$

**2. Describe** the following using variables

- A. Four times a number. E. The product of a number and 20.  
 B. One third of a number. F. Five less than a number.  
 C. Ten more than a number. G. Ten more than twice a number.  
 D. The sum of two different numbers. H. Six less than twice a number.

**3. Express** the following in words.

- A.  $x-1$       B.  $7x$       C.  $3+4x$       D.  $\frac{x}{3}-1$

**4. Find** the value of  $y$  in the relation  $y = 2x + 3$  when

- A.  $x = 1$       B.  $x = 5$       C.  $x = -4$       D.  $x = \frac{2}{3}$

**4.1.2. Variables, Terms and Expressions****Activity 4.1.2****1.** Which of the following are terms and which are not terms?

- A.  $x+y$     B.  $2x + 3$     C.  $2x$     D.  $-3x$     E.  $\frac{3}{y}$     F.  $4$

**2.** Find the numerical coefficient of the following terms.

- A.  $-3x$     B.  $y$     C.  $-xy^2$     D.  $\frac{xy}{4}$

**3.** Identify whether each pair of the following are like terms or unlike terms?

- A.  $5x^2$  and  $3y^2$       C.  $2xy$  and  $yx$   
 B.  $3x$  and  $x$       D.  $8x^2 y$  and  $4xy^2$

**Definition 4.2:**

*a constant (a number), a variable, product or quotient of a number and variable is called a **term**.*

**Example:**

- 4 is a term      •  $12xy-4$  is not a term      •  $x-y$  is not a term
- $xy$  is a term      •  $\frac{3x}{y}$  is a term

**Definition 4.3:**

*In the product of a number and variable, the numerical factor of the term is called **numerical coefficient**.*

**Example:**

- A. The numerical coefficient of  $6ab$  is 6
- B. The numerical coefficient of  $-3xy$  is  $-3$
- C. The numerical coefficient of  $x$  is 1 .
- D. The numerical coefficient of  $-\frac{2ab}{3}$  is  $-\frac{2}{3}$

**Definition 4.4:**

**like terms** are terms whose variable and exponent of variable are exactly the same but they may differ in their numerical coefficients. Terms which are not like terms are called **unlike terms**.

A term which do not contain a variable is called constant term. All constant terms are like terms.

**Example:**

- 2 and  $-5$  are like terms, because they are constant terms.
- $\frac{3}{5}a^2b^2$  and  $-\frac{2}{3}b^2a^2$  are like terms.
- $-20abc$  and  $acb$  are like terms
- $3xyz$  and  $x^2y^2z^2$  are unlike terms
- $-2a^3b^3c^3$  and  $-2a^3b^2c^3$  are unlike terms
- $3xy$  and  $3yx$  are like terms

**Exercise 4.1.2**

1. Identify whether each pair of the following are like or unlike terms.

- |                              |                                    |
|------------------------------|------------------------------------|
| A. $3x$ and $-5x$            | E. $xy^2$ and $y^2x$               |
| B. $20xy$ and $ab$           | F. $3x^2$ and $2x^3$               |
| C. $-80ab$ and $15a^2b^2$    | G. $xyz^2$ and $z^2xy$             |
| D. $5ab$ and $\frac{1}{4}ab$ | H. $\frac{3}{x}$ and $\frac{5}{x}$ |

2. Which of the following are terms and which are not terms?  
 A.  $5x-4y$    B.  $8x^2$    C.  $\frac{2x-y}{5}$    D.  $-9abc$    E.  $\frac{8y}{x}$    F. 0
3. In each of the following expressions, determine the numerical coefficient.  
 A.  $2bm$    C.  $-4x$    E.  $\frac{y}{2}$   
 B.  $5yx^2$    D.  $\frac{3x}{4}$    F.  $-z$

**Definition 4.5:**

*Algebraic expressions are formed by using numbers, letters and the operation of addition, subtraction, multiplication and division.*

**Example:**  $12, 2x-3, a+10, 3x, \frac{3}{4}a, 3x, 3x+2a-5$ , etc. are algebraic expressions.

**Note:**

*The terms of an algebraic expression are parts of the expression that are connected by plus or minus signs.*

**Example:** In the expression  $3x+2a-5$ , the terms are  $3x, 2a$  and  $-5$ . and the constant term is  $-5$ .

**Definition 4.6:**

an algebraic expression in algebra which contains one term is called a **monomial**.

Example:  $5x, 12ab, 3xyz, 0.2ab \dots$  are monomial.

**Definition 4.7:**

an algebraic expression in algebra which contains two terms is called a **binomial**.

Example:  $5x+3y, 12-ab, 3x+yz, x-b \dots$  are binomial.

### Definition 4.8:

*an algebraic expression in algebra which contains three terms is called a **trinomial**.*

**Example:**  $2x+3y-3$ ,  $12+a-b$ ,  $x-y-z$ ,  $ab+6+z$  are trinomial.

## Simplifying Algebraic Expressions

*To simplify any algebraic expression, follow the following basic steps*



**Remove** brackets using distributive property.



**Collecting** like terms and



**Add** or subtract like terms

### Note:

*1. When you add or subtract like terms, add or subtract their numerical coefficients.*

*Example:*  $-3x+5x = (-3+5)x=2x$

*2. Unlike terms cannot be added or subtracted*

*Example:*  $5x+2y$  cannot be added.

*3. When you remove brackets using distributive property be careful of signs. Example:*

$$A. -2(x-4) = -2x - (-2) \times 4 = -2x + 8$$

$$B. -3(-2x-1) = -3(-2x) - (-3)(1) = 6x + 3$$

*4. When you multiply a number with variable, you can write the number as coefficient without multiplication sign*

*Example:*       $A. 4 \times y = 4y$                        $B. 5 \times (3y) = 15y$

**Example:** Simplify the following algebraic expression

A.  $3x + 2y - 4x + 5y$

D.  $4(a + 2b) - 3(2a - b) + 6a - 7b$

B.  $3(x + 4) - 5(x - 2)$

E.  $3(2a + 3) - 2(a + 8)$

C.  $20 + 4(x + 3y) - 4x - 8y$

**Solution:**

$$a(b+c)=ab+ac$$

A.  $3x + 2y - 4x + 5y$

- *collecting like terms*

$$= 3x - 4x + 2y + 5y$$

- *adding/subtracting coefficient of like terms*

$$= -x + 7y$$

B.  $3(x + 4) - 5(x - 2)$

- *distributive*

$$= 3x + 12 - 5x + 10$$

- *collecting like terms*

$$= 3x - 5x + 12 + 10$$

- *combine constants*

$$= (3 - 5)x + 22$$

- *Simplify*

$$= -2x + 22$$

C.  $20 + 4(x + 3y) - 4x - 8y$

- *distributive*

$$= 20 + 4x + 12y - 4x - 8y$$

- *collecting like terms*

$$= 4x - 4x + 12y - 8y + 20$$

- *combine constants*

$$= (4 - 4)x + (12 - 8)y + 20$$

- *Simplify*

$$= 4y + 20$$

D.  $4(a + 2b) - 3(2a - b) + 6a - 7b$

- *distributive*

$$= 4a + 8b - 6a + 3b + 6a - 7b$$

- *collecting like terms*

$$= 4a - 6a + 6a + 8b + 3b - 7b$$

- *combine constants*

$$= (4 - 6 + 6)a + (8 + 3 - 7)b$$

- *Simplify*

$$= 4a + 4b$$

E.  $3(2a + 3) - 2(a + 8)$

- *distributive*

$$= 6a + 9 - 2a - 16$$

- *collecting like terms*

$$= 6a - 2a + 9 - 16$$

- *combine constants*

$$= (6 - 2)a + (9 - 16)$$

- *Simplify*

$$= 4a - 7$$

**Exercise 4.1.3**

Categorize the following expressions as monomial, binomial or trinomial and determine the constant terms in each expression.

A. 24

B.  $5x$

C.  $-21t - 2x$

D.  $5x - 3y - 4$

E.  $20w + w^2$

F.  $3x - 2y + 1$

G.  $z - y - 25x$

H.  $y$

I.  $2b - 4a$

2. Simplify the following expressions by collecting like terms

A.  $3x + 35y - 8x$

B.  $15 + 21b - 32b + 3$

C.  $3(3x - 6y) + x + 18y - 4$

D.  $3z - 12 - 11y + x + y - 48z - 3$

E.  $-3(-a + 6) + x - (z + 4) - 2x$

F.  $xy + ab - cd + 2xy - ab + dc$

G.  $3x^2 + 4x + 6 - x^2 - 3x - 3$

H.  $.5 + 2y + 3y^2 - 8y - 6 + 2y^2$

I.  $2x^2 - 3x + 8 + x^2 + 4x + 4$

J.  $.5(2 - x^2) - 3(x^2 - 4y)$

## 4.2. SOLVING LINEAR EQUATIONS

### Learning Outcomes:

At the end of this unit, learners will be able to:

- ⇒ Define a linear equation
- ⇒ Solve linear equations involving brackets
- ⇒ Solve linear equations involving fractions

### 4.2.1. Linear Equations Involving Brackets



#### Activity 4.2.1

Discuss with your friends

1. Connect the following a pair of expressions by using '=' sign

A.  $2x$  and  $2$

B.  $3x - 1$  and  $6$


C.  $y + 4$  and  $5 - y$

2. Using the following balance, find the unknown variables.

A.



B.



C.



3. What replacement of  $x$  will make the following equality is true?

A.  $x - 1 = 1$       B.  $x + 13 = 0$       C.  $4x = 8$       D.  $2x = 0$

4. Determine whether each the following are equations or not.

A.  $2 + x$       B.  $x = 2$       C.  $x > 2$       D.  $2x - 1 = 0$

5. What is equation?

### Definition 4.9:

Two different algebraic expressions connected by equal sign ( $=$ ) is called **equation**.

### Definition 4.10:

A **Linear equation** in one variable  $x$  is an equation which can be written in standard form,  $ax + b = 0$ , where  $a$  and  $b$  are constant numbers with  $a \neq 0$

### Note:

An equation of a single variable in which the highest **exponent** of the variable one is a **linear equation**.

$3x$  is the same as  $3x^1$  ← Exponent is 1

$5x^2$  ← Exponent is 2

**Example:** which of the following equations are linear and which are not linear.

A.  $2x = 2$

C.  $2x^2 + 3 = 2x - 1$

B.  $x + 3 = 2x - 1$

D.  $x = 2x^3 - 6$

## Solution:

$a$  and  $b$  are linear equation, because the highest exponent of variable is one.

$c$  and  $d$  are not linear equation, because the highest exponent of the variable is not one.

## Note:




All equations have two sides with respect to the equal sign. Left hand side and (L.H.S) and right-hand side (R.H.S)

**Example:** In the equation  $3x+5 = x-1$

L.H.S is  $3x+5$  and R.H.S is  $x-1$




## Rules of transforming equations to simpler form

Consider the following balances,

A.  =  add 4 kg to both sides 




**Rule 1:** adding the same number to both sides of equation will not affect the equation.

i.e. if  $a=b$ , then  $a+c=b+c$ , where  $a, b$ , and  $c$  are any constant number.

B.  =  subtracting 2kg from both sides 





**Rule 2:** subtracting the same number to both sides of equation will not affect the equation.

i.e. if  $a=b$ , then  $a - c = b - c$ , where  $a, b, c$  any constant number

C.  =  multiplying both sides by 2 

**Rule 3:** multiplying both sides of equation with the same none zero number will not affect the equation.

If  $a=b$ , then  $a \times c = b \times c$ , where  $a, b, c$  any constant number and  $c \neq 0$

D.  =  dividing both sides by 2  = 

**Rule 4:** dividing both sides of equation with the same none zero number will not affect the equation.

If  $a=b$ , then  $\frac{a}{c} = \frac{b}{c}$ , where  $a, b, c$  any constant number and  $c \neq 0$

**Example:** Find the value of  $x$  by using rules of transformation

A.  $x - 10 = 6$

B.  $10 - x = 6$

C.  $x + 4 = -6$

**Solution:**

A.  $x - 10 = 6$

$$\begin{array}{r} x - 10 = 6 \\ +10 \quad +10 \\ \hline \end{array}$$

$$x = 6 + 10$$

$$\underline{x = 16}$$

B.  $10 - x = 6$

$$\begin{array}{r} 10 - x = 6 \\ -10 \quad -10 \\ \hline \end{array}$$

$$-x = 6 - 10$$

$$-x = -4$$

$$\underline{x = 4}$$

C.  $x + 4 = -6$

$$\begin{array}{r} x + 4 = -6 \\ -4 \quad -4 \\ \hline \end{array}$$

$$x = -6 - 4$$

$$x = -10$$

$$\underline{x = -10}$$

**Example:** Find the value of  $y$  by using rules of transformation

A.  $8y = 72$

B.  $\frac{y}{5} = -2$

**Solution:**

A.  $8y = 72$ ,

to find the value of  $y$  divide both sides by 8

$$\frac{8y}{8} = \frac{72}{8}$$

$$\underline{y = 9}$$

B.

to find the value multiply both sides by 5

$$\cancel{5} \times \frac{y}{\cancel{5}} = -2 \times 5$$

$$\underline{y = -10}$$

### Exercise 4.2.1

- Decide each of the following either linear equation or not.
 

A. $2y - 2$	C. $2x^2 = 4$	E. $3x = 0$
B. $x - 1 = 5$	D. $x - 3 + 4x = 2x - 2$	F. $x > 0$
- By what transformation was the second equation obtained from the first?
 

A. $x + 1 = 2$ $x = 1$	B. $-8x = 12$ $x = -1.5$	C. $2x - 4 = 6$ $2x = 10$	D. $\frac{2x}{3} = 5$ $2x = 15$
---------------------------	-----------------------------	------------------------------	------------------------------------
- Find the value of unknown variable by using rules of transformation
 

A. $-8x = 12$	B. $\frac{2x}{5} = 10$	C. $20 - x = 15$	D. $2x - 4 = 16$
---------------	------------------------	------------------	------------------

### Solving equations containing variable on both sides of the equation

Consider the equation  $x - 5 = 10$ , in solving this equation you add 5 on both sides and get  $= 10 + 5$ , this is the same as shifting  $-5$  to the other side of the equation and it becomes  $+5$ . i.e.

$$x - 5 = 10 + 5, \quad \text{so } x = 10 + 5 = 15$$

Shifting  $-5$  to the other side, becomes  $+5$

**Note:**

*when you shift a variable (term) or number from one side of the equation to the other side its **sign changes**.*

For example, in the equation  $3x - 4 = 2x - 1$ , shifting  $2x$  to the left side you get

$$3x - 4 = +2x - 1 \quad \text{So, it becomes } 3x - 2x - 4 = -1$$

In solving equation containing variable on both sides, collect the terms containing variables on one side and numbers on the other side and do not forget changing the signs when you shift a term from one side to the other.

**Example 3:** solve the following equations and check the result

A.  $2x = x + 4$

B.  $2x - 8 = 4$

C.  $5x - 10 = 6 - 3x$

D.  $x + 5 - 4x = 3 + 2x - 6 - 4x$

**Solution:**

A.  $2x = x + 4$

(shifting  $x$  to the left it becomes  $-x$ )

$$2x - x = 4$$

$$x = \underline{4}$$

**Check:**  $2x = x + 4$ ,

substitute 4 in the place of  $x$

$$2 \times 4 = 4 + 4$$

$$8 = 8, \text{ (True)}$$

Hence,  $x = 4$  is the solution

C.  $5x - 10 = 6 - 3x$

(Transform the variable to the left side & the constant to the right)

$$5x + 3x = 6 + 10$$

$$8x = 16$$

(divide both sides by 8)

$$x = \underline{2}$$

**Check:**  $5x - 10 = 6 - 3x$ ,

substitute 2 in place of  $x$

$$(5 \times 2) - 10 = 6 - (3 \times 2)$$

$$0 = 0 \text{ (True)}$$

Hence,  $x = 2$  is the solution

B.  $2x - 8 = 4$

(Shifting  $-8$  to the right becomes  $+8$ )

$$2x = 12 + 8$$

(divide both sides by 2)

$$x = \underline{6}$$

**Check:**  $2x - 8 = 4$

substitute 4 in place of  $x$

$$(2 \times 6) - 8 = 4$$

$$12 - 8 = 4$$

$$4 = 4 \text{ (True)}$$

Hence,  $x = 6$  is the solution

D.  $x + 5 - 4x = 3 + 2x - 6 - 4x$

(Collect and simplify like terms on the same side)

$$x - 4x + 5 = 3 - 6 + 2x - 4$$

$$-3x + 5 = -3 - 2x$$

$$-3x + 2x = -3 - 5$$

$$-x = -8$$

divide both sides by  $-1$ )

$$x = 8$$

**Check:**  $x + 5 - 4x = 3 + 2x - 6 - 4x$

$$8 + 5 - (4 \times 8) = 3 + (2 \times 8) - 6 - (4 \times 8)$$

$$13 - 32 = 3 + 16 - 6 - 32$$

$$-19 = -19 \text{ (True),}$$

Hence,  $x = 8$  is the solution

### Exercise 4.2.2

1. solve the following equations and check the result

- A.  $3x-9=4x+5$    C.  $12x+7=5-3x+17$    E.  $3x-9=-x+19$    G.  $10=3x-5+2x$   
 B.  $5x-3-4x=13$    D.  $-x+3=x+9$    F.  $8x+5=2x+17$    H.  $x-5=2x+13 + x$

### 4.2.2 Solving Linear equations involving brackets

#### Activity 4.2.2

#### Discuss with your friends

1. Simplify the following by using distributive property

- A.  $3(x + 2)$    C.  $4(x - 1) + 2(1 - 2x)$   
 B.  $-(-2x + 9)$    D.  $4(x - 1) - 2(x - 9)$

2. Solve the following linear equations by using distributive property.

- A.  $3(x + 2) = 0$    C.  $4(x - 1) = 2(1 - 2x)$   
 B.  $-(-2x + 9) = 1$    D.  $4(x - 1) = 2(x - 9)$

#### To solve linear equations involving brackets

1. Remove the brackets using distributive property
2. Collect the terms containing variables on one side and constants on the other side, do not forget to change sign while shifting the terms.

**Note:**

For any numbers  $a$ ,  $b$  and  $c$

1.  $a+(b+c) = a+b+c$    3.  $a - (b + c) = a - b + c$    5.  $a(b-c)=ab-ac$   
 2.  $a-(b+c) = a-b-c$    4.  $a(b+c) = ab+ac$

#### Example 1:

Remove the bracket and simplify the following expressions

- A.  $3 - (2x - 5)$    C.  $3m - 2(m - 3)$   
 B.  $2y + (4 - y)$    D.  $-5n + 3(4 - 2n)$

**Solution:**

$$\begin{aligned}
 A. \quad & 3-(2x-5) \\
 &= 3 - 2x + 5 \\
 &= 3 + 5 - 2x \\
 &= 8 - 2x \\
 &= -2x + 8
 \end{aligned}$$

□	Note:	□
	$a-b=-b+a$	
□	$8-2x=-2x+8$	□

$$\begin{aligned}
 B. \quad & 2y+(4-y) \\
 &= 2y+ 4 - y \\
 &= 2y - y + 4 \\
 &= y + 4
 \end{aligned}$$

$$\begin{aligned}
 C. \quad & 3m - 2 ( m - 3 ) \\
 &= 3m - 2m + 6 \\
 &= m + 6
 \end{aligned}$$

$$\begin{aligned}
 D. \quad & -5n + 3( 4 - 2n ) \\
 &= -5n + 12 - 6n \\
 &= -5n - 6n + 12 \\
 &= -11n + 12
 \end{aligned}$$

**Example 2:** solve each of the following equations.

$$A. \quad 4(2x+3)=36$$

$$C. \quad -10y-(-2y+3)=21$$

$$B \quad 6(5x-7)=4(3x+7)$$

$$D. \quad 4(x-1)+3(x+2)=5(x-4)$$

**Solution:**

$$A. \quad 4(2x+3)=36$$

(Removing the bracket)

$$4 \times 2x + 4 \times 3 = 36$$

$$8x + 12 = 36$$

(Shifting +12 in to right side,  
becomes -12)

$$8x = 36 - 12$$

( Divide both sides by 8)

$$\begin{aligned}
 \frac{8x}{8} &= \frac{24}{8} \\
 x &= 3
 \end{aligned}$$

$$B. \quad 6(5x-7)=4(3x+7)$$

( Removing brackets)

$$30x - 42 = 12x + 28$$

(Shifting variables to the left and  
constants to the right)

$$30x - 12x = 28 + 42$$

(Divide both sides by 18)

$$\begin{aligned}
 \frac{18x}{18} &= \frac{70}{18} \\
 x &= 35/9
 \end{aligned}$$

$$C. -10y - (-2y + 3) = 21$$

(Removing the brackets)

$$-10y + 2y - 3 = 21$$

$$-8y - 3 = 21$$

$$-8y = 21 + 3$$

(Divide both sides by -8)

$$\frac{-8y}{-8} = \frac{24}{-8}$$

$$y = -3$$

$$D. 4(x - 1) + 3(x + 2) = 5(x - 4)$$

(Removing the brackets)

$$4x - 4 + 3x + 6 = 5x - 20$$

(Collecting like terms in one sides)

$$4x + 3x - 4 + 6 = 5x - 20$$

$$7x + 2 = 5x - 20$$

(Simplify in one side)

$$7x - 5x = -20 - 2$$

(Divide both sides by 2)

$$\frac{2x}{2} = \frac{22}{2}$$

$$x = -11$$

### Exercise 4.2.3

1. Solve each of the following equations

A.  $(x - 1) = 11$

E.  $7 - (x + 1) = 9 - (2x - 1)$

B.  $6(8y + 3) = 4(7y + 5)$

F.  $5(1 - 2a) - 3(4 - 4a) = 0$

C.  $8(2y - 6) = 5(3y - 7)$

G.  $3x + 7 + 3(x - 1) = 2(2x + 6)$

D.  $3 - 2(2x + 1) = x - 14$

H.  $5(2x + 1) + 3(3x - 4) = 4(5x - 6)$

2. Solve for x in terms of m and n for

A.  $m(x - 1) = 0$

C.  $n(x - 2) = m + x, n \neq 1$

B.  $m(x + n) = mn$

D.  $x(n - 1) = 4 - x$

### 4.2.3. Solving linear equation involving fractions



#### Activity 4.2.3

1. Find the LCM of the following numbers

A. 2 and 3

B. 4 and 6

C. 3, 4 and 6

D. 4 and 8

2. Multiply both side of the equation by the LCM of the denominator, simplify and Solve the equation

A.  $\frac{x}{2} = \frac{5}{2}$

B.  $\frac{3}{4} = \frac{x}{8}$

C.  $\frac{x}{3} + 8 = 7$

D.  $\frac{3}{4} + \frac{x}{2} = \frac{x}{3}$

**Note:****To solve linear equations involving fractions, follow the following steps:**

1. Find the list common multiple (LCM) of denominators
2. Multiply both sides by LCM
3. Solving linear equation by shifting the terms containing variable to one side and constants to the other side.

**Example: Solve the following linear equations**

A.  $\frac{x+2}{4} = \frac{5}{3}$

B.  $\frac{x}{8} + \frac{7}{4} = \frac{5}{6}$

C.  $\frac{x+1}{3} + \frac{x-1}{10} = 12$

D.  $\frac{2x+3}{3} + \frac{x-1}{2} + \frac{x+1}{4} = 8$

**Solution:**

A.  $\frac{x+2}{4} = \frac{5}{3}$

- The LCM of 4 and 3 is 12, therefore multiply both side by 12

$$12\left(\frac{x+2}{4}\right) + 12\left(\frac{x-1}{2}\right) + 12\left(\frac{x+1}{4}\right) = 12(8)$$

- Simplify

$$= 3(x+2) = 4(5)$$

- removing brackets

$$= 3x+6= 20$$

- collecting like terms =

$$3x = 20-6$$

- Simplify

$$3x = 14$$

- dividing both sides by 6,

$$\frac{3x}{3} = \frac{14}{3}$$

$$x = \frac{14}{3}$$

B.  $\frac{x}{8} + \frac{7}{4} = \frac{5}{6}$

- The LCM of 8, 4 and 6 of is 24, therefore multiply both side by 24.

$$24\left(\frac{x}{8} + \frac{7}{4}\right) = 24\left(\frac{5}{6}\right)$$

- distributive property

$$24\left(\frac{x}{8}\right) + 24\left(\frac{7}{4}\right) = \left(24\frac{5}{6}\right)$$

- Simplify,

$$3x + 42 = 20$$

- collecting like terms,

$$3x = 20- 42$$

- Simplify,

$$3x = - 22$$

- dividing both side by 3,

$$\frac{3x}{3} = \frac{-22}{3}$$

$$x = - \frac{22}{3}$$

C.  $\frac{x+1}{3} + \frac{x-1}{10} = 12$

- The LCM of 3 and 10 is 30, therefore multiplying both sides by 30

$$30\left(\frac{x+1}{3} + \frac{x-1}{10}\right) = 12 \times 30$$

- distributive property

$$30\left(\frac{x+1}{3}\right) + 30\left(\frac{x-1}{10}\right) = 360$$

- Simplify,

$$10(x+1) + 3(x-1) = 360$$

- removing brackets,

$$10x + 10 + 3x - 3 = 360$$

- collecting like terms

$$10x + 3x = 360 - 10 + 3$$

- Simplify

$$13x = 353$$

- dividing both sides by 13

$$\frac{13x}{13} = \frac{353}{13}$$

$$x = \frac{353}{13}$$

D.  $\frac{2x+3}{3} + \frac{x-1}{2} + \frac{x+1}{4} = 8$

- The LCM of denominator is 12, therefore multiplying both sides by 12

$$12\left(\frac{2x+3}{3} + \frac{x-1}{2} + \frac{x+1}{4}\right) = 8 \times 12$$

- distributive property

$$12\left(\frac{2x+3}{3}\right) + 12\left(\frac{x-1}{2}\right) + 12\left(\frac{x+1}{4}\right) = 96$$

- Simplify,

$$4(2x+3) + 6(x-1) + 3(x+1) = 96$$

- removing brackets,

$$8x + 12 + 6x - 6 + 3x + 3 = 96$$

- collecting like terms

$$8x + 6x + 3x = 96 - 12 + 6 - 3$$

- Simplify

$$17x = 87$$

- dividing both sides by 17

$$\frac{17x}{17} = \frac{87}{17}$$

$$x = \frac{87}{17}$$

### Definition 4.11:

Equivalent equations are two or more equations that have the same solution.

**Example 1:** Show that  $2x - 1 = 11$  and  $x + 12 = 3x$  are equivalent equations.

### Solution:

$$2x - 1 = 11 \quad \text{and} \quad x + 12 = 3x$$

$$2x = 11 + 1 \quad \text{and} \quad x - 3x = -12$$

$$2x = 12 \quad \text{and} \quad -2x = -12$$

$$x = 6 \quad \text{and} \quad x = 6$$

Hence,  $2x - 1 = 11$  and  $x + 12 = 3x$  are equivalent equations.

**Example 2:** show that  $2(x-5) = x+6$  and  $2x-4=10$  are not equivalent

**Solution:**

$$2(x-5)=x+6 \quad \text{and} \quad 2x-4=10$$

$$2x-10=x+6 \quad \text{and} \quad 2x-4=10$$

$$2x-x=6+10 \quad \text{and} \quad 2x=10+4$$

$$x=16 \quad \text{and} \quad x=7$$

**Therefore,  $2(x-5)=x+6$  and  $2x-4=10$  do not have the same solution, so they are not equivalent.**

**Exercise 4.2.4**

1. Solve the following

A.  $\frac{5x}{13} + \frac{5x}{26} = 1$

B.  $\frac{3x}{7} + \frac{35x}{8} = 10$

C.  $\frac{2x}{5} - \frac{2}{3} = \frac{x}{2} + 6$

D.  $\frac{-3}{5} + \frac{x}{10} = \frac{-1}{5} - \frac{x}{5}$

E.  $\frac{9w}{7} - 6 = \frac{5w}{7} + 12$

F.  $\frac{7}{10}x + \frac{3}{2} = \frac{3}{5}x + 2$

G.  $\frac{1}{2}(4x + 6) = \frac{1}{3}(9x - 24)$

H.  $\frac{n+1}{2} + \frac{n+2}{3} + \frac{n+4}{4} = 2$

I.  $\frac{1}{3}(x+6) - \frac{1}{2}(3x-4) = 5$

2. Which of the following pairs of equations are equivalent?

A.  $x=0$  and  $2x-1=-1$

C.  $5x=10$  and  $14=2x+4$

B.  $2x-6=2$  and  $-x=-5$

D.  $x=0$  and  $324x+55=55$

**4.3. CARTESIAN COORDINATE SYSTEM****Learning Outcomes:**

**At the end of this unit, learners will be able to:**

- ✎ Describe the Cartesian coordinate system
- ✎ Draw the four quadrants of the Cartesian plane and mark the origin, x-axis and y-axis
- ✎ Plot points on the Cartesian coordinate plane given their coordinates
- ✎ Draw graph of linear equations like  $y = mx$ , in a Cartesian coordinate plane using table values manually and computer applications

### 4.3.1. The Four Quadrants of the Cartesian coordinate Plane



#### Activity 4.3.1

#### Discuss with your friends


1. Draw horizontal number line and mark numbers.
2. Draw vertical number line perpendicular to horizontal number line that pass through zero.
3. Mark the numbers above and below zero on the vertical line.
4. The horizontal number line is called \_\_\_\_\_.
5. The vertical number line is called \_\_\_\_\_.
6. The two intersecting number lines (vertical and horizontal) set up a plane is called \_\_\_\_\_.

#### Definition 4.12:

The two perpendicularly intersecting horizontal and vertical number lines together set up a plane is called **Cartesian coordinate plane**.

- The horizontal number line is called ***x-axis***.
- The vertical number line is called ***y-axis***.
- The point where, *x-axis* and the *y-axis* intersect is called ***origin***.

#### Coordinate Plane or Cartesian Plane



### Locating points on the coordinate plane

- Points on Cartesian plane are described by two numbers that are called **Coordinate**.
- The first number  $a$ , is the horizontal position of the point from the origin. It is called  **$x$ -coordinate (abscissa)**.
- The second number is  $b$ , the vertical position of the point from the origin. It is called  **$y$ -coordinate (ordinate)**.

To locate a point  $(a, b)$  on a coordinate plane follow the following steps.

**Step 1:** Move  $a$  unit in the horizontal direction from the origin.


- Move  $a$  unit to the right if  $a$  is positive.
- Move  $a$  unit to the left if  $a$  is negative.

**Step 2:** Then, move  $b$  unit in the vertical direction

- Move  $b$  unit up if  $b$  is positive.
- Move  $b$  unit down if  $b$  is negative.


**Example 1:** locate the point A  $(3, 2)$  on the coordinate plane

**Solution:**



**Example 2:** locate the point A  $(-3, 2)$  on the coordinate plane

**Solution:** move 3 units to the left from the origin and then 2 units up.



**Example 3:** locate the point A  $(2, 0)$  on the coordinate plane

**Solution:**

*move 2 units to the right from the origin and then 0 unit in the vertical direction (no move in the vertical direction)*


**Example 4:** locate the point A  $(0, -4)$  on the coordinate plane.

**Solution:**


*start from origin and no move on horizontal, then move 4 units down in vertical direction.*

**Note:**

1. The coordinate of all points on the x-axis has the form




2. The coordinate of all points on the  $y$ -axis has the form




### **Determining the coordinate of a given point**

**To determine the coordinate of a point .**

- First draw vertical line passing through the point  $P$  crossing  $x$  - axis. Let it crosses at  $a$ , then the coordinate of  $P$  is  $a$
- Then, draw horizontal line passing through  $P$  crossing  $y$ -axis. Let it crosses at " $b$ ", then the  $y$  coordinate of  $P$  is  $b$
- So, the coordinate of  $P$  is  $(a, b)$ .



**Example:** Determine the coordinates of the points indicated below



**Solution:**

to determine coordinate of A, draw vertical line passing through A. it crosses x -axis at 3, hence its x coordinate is 3. Then draw horizontal line passing through A, it crosses y – axis at 4, hence its y coordinate is 4.


Therefore, the coordinate of A is  $(3, 4)$ . In similar method you can determine the coordinate of the rest points.

$B(-3, 4)$ ,  $C(3, -4)$ ,  $D(-3, -4)$ ,  $E(-1, 0)$ ,  $F(0, -1)$

## Quadrants

The x-axis and y-axis divide the Cartesian plane into 4 regions known as **quadrants**.

The four quadrants (1<sup>st</sup> quadrant, 2<sup>nd</sup> quadrant, 3<sup>rd</sup> quadrant, and 4<sup>th</sup> quadrant) are shown in the diagram below.



- Quadrant I contains **positive x** values and **positive y** values.

- Quadrant 2 contains **negative x** values and **positive y** values.
- Quadrant 3 contains **negative x** values and **negative y** values.
- Quadrant 4 contains **positive x** values and **negative y** values.

**Example 2:** In which quadrant or axis, the following points lie?

A. (4, 1) B. (-4, 1) C. (0, 9) D. (-3, -3) E. (-3, 0) F. (1, -3)

**Solution:**

A. both x and y values are positive then (4, 1) lie 1<sup>st</sup> quadrant

B. x is **negative** and y is **positive** then (-4, 1) lie on 2<sup>nd</sup> quadrant

C. x coordinate is 0, then (0, 9) lie on y-axis

D. both x and y values are **negative**, then (-3, -3) lie on 3<sup>rd</sup> quadrant

E. y coordinate is 0 then (-3, 0) lie on x-axis.

F x is **positive** and y is **negative** then (1, -3) lie on 4<sup>th</sup> quadrant

### Exercise 4.3.1


1. **Locate** the following points on the same Cartesian coordinate plane.

- |              |               |                             |
|--------------|---------------|-----------------------------|
| A. M (-1, 4) | D. P (-3, -2) | G. S (-2, 1.5)              |
| B. N (4, 6)  | E. Q (-5, 0)  | H. T ( $-\frac{3}{2}$ , -5) |
| C. O (4, -1) | F. R (0, -3)  | I. U (0, 0)                 |

2. Based on the coordinate plane below answer the following questions.

A. Write the coordinates of the points F, T, P, M, N, Q and k

B. Which points has the coordinate (-2, 3)



3. In which quadrant or axis lie the following coordinate points?

A. (5, 212)

D. (-45, -4)

G.  $(\frac{5}{2}, 0)$

B. (-3, 27)

E. (10, 0)

H. (35, -12.5)

C. (0, -16)

F. (12, -30)



I.  $(-\frac{7}{4}, -\frac{13}{5})$

### 4.3.2 Coordinates and graph of linear equations



#### Activity 4.3.2

1. Write the coordinate of the points that lie on the lines  $x=4$  and  $y=5$  indicated below



2. Plot the following points on coordinate plane and connect them using straight line.  $(-2, 0)$ ,  $(-2, -3)$ ,  $(-2, 3)$

3. Plot the following points on coordinate plane and connect them using straight line.  $(0, -2)$ ,  $(-3, -2)$ ,  $(3, -2)$

4. What is the equation of the line in activity 2?

5. What is the equation of the line in activity 3?

**Graph of an equation of the form  $x = a$ , where "a" is constant**

To draw the graph of a line  $x = a$ , follow the following steps:

**Step 1:** Prepare table of values relating  $x$  and  $y$ . In the case of  $x = a$ , for any different possible values of  $y$ , the value of  $x$  is constant. It is  $x = a$ .

**Step 2:** plot the coordinate of the points in step 1.

**Step 3:** Join all points in step 2 using straight line.

**Example1:** Draw the graph of the following equation of lines on the same Cartesian coordinate plane.

A.  $x=-4$

B.  $x-3=0$

C.  $x=\frac{3}{2}$

**Solution:**

A.  $x=-4$ , for any possible values of  $y$ , the value of  $x$  is  $-4$

x	-4	-4	-4	-4	-4	-4	-4
y	-3	-2	-1	0	1	2	3
(x,y)	(-4,-3)	(-4,-2)	(-4,-1)	(-4,0)	(-4,1)	(-4,2)	(-4,3)


B.  $x+3=0$

$x = -3$ , for any possible values of  $y$ , the value of  $x$  is  $-3$

x	-3	-3	-3	-3	-3	-3	-3
y	-3	-2	-1	0	1	2	3
(x,y)	(-3,-3)	(-3,-2)	(-3,-1)	(-3,0)	(-3,1)	(-3,2)	(-3,3)

C.  $x=\frac{3}{2}$ , for any possible values of  $y$ , the value of  $x$  is  $\frac{3}{2}$

x	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$
y	-3	-2	-1	0	1	2	3
(x,y)	$(\frac{3}{2}, -3)$	$(\frac{3}{2}, -2)$	$(\frac{3}{2}, -1)$	$(\frac{3}{2}, 0)$	$(\frac{3}{2}, 1)$	$(\frac{3}{2}, 2)$	$(\frac{3}{2}, 3)$



**Note:**

- The graph of the equation of the line  $x=a$ ,  $a$  is constant is a line parallel to  $y$ -axis at a distance of  $a$  unit from the origin.
- If  $a$  is positive, then the line lies to the **right** of the  **$y$ -axis**.
- If  $a$  is negative, then the line lies to the **left** of  **$y$ -axis**.
- If  $a=0$ , then the line lies **on  $y$ -axis**.

### Graph of an equation of the form $y=b$ , where $b$ is constant

To draw the graph of a line  $y=b$ , follow the following steps:

**Step 1:** Prepare table of values relating  $x$  and  $y$ . In the case of  $y=b$ , for any different possible values of  $x$ , the value of  $y$  is constant. It is  $y=b$ .

**Step 2:** plot the coordinate of the points in step 1.

**Step 3:** Join all points in step 2 using straight line.

**Example 1:** Draw the graph of the following equation of lines on the same Cartesian coordinate plane.

A.  $y = -4$

B.  $y = 3$

C.  $y - \frac{3}{2} = 0$

**Solution:**

A.  $y = -4$


x	-3	-2	-1	0	1	2	3
y	-4	-4	-4	-4	-4	-4	-4
(x,y)	(-3,-4)	(-2,-4)	(-1,-4)	(0,-4)	(1,-4)	(2,-4)	(3,-4)

B.  $y=3$

X	-3	-2	-1	0	1	2	3
Y	3	3	3	3	3	3	3
(x,y)	(-3,3)	(-2,3)	(-1,3)	(0,3)	(1,3)	(2,3)	(3,3)

C.  $y - \frac{3}{2} = 0$   
 $y = \frac{3}{2}$

x	-3	-2	-1	0	1	2	3
y	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$
(x,y)	$(-3, \frac{3}{2})$	$(-2, \frac{3}{2})$	$(-1, \frac{3}{2})$	$(0, \frac{3}{2})$	$(1, \frac{3}{2})$	$(2, \frac{3}{2})$	$(3, \frac{3}{2})$

**Note:**

- The graph of the equation of the line  $y=b$ ,  $b$  is constant is a line parallel to  $x$ -axis at a distance of  $b$  unit from the origin.
- If  $b$  is positive, then the line lies **above** the  $x$ -axis.

- If  $b$  is negative, then the line lies below the **x-axis**.
- If  $b = 0$ , then the line lies **on x-axis**.

### Exercise 4.3.2

- Draw the graph of the following equations on the same coordinate plane.  
A.  $x = 7$       B.  $y = 5$       C.  $x = \frac{1}{2}$       d.  $y = 0$
- Which of the following points lie on a line  $y=1$   
A.  $(-1, 1)$       C.  $(0, 0)$       E.  $(-3, 1)$       G.  $(2, 2)$       I.  $(1, 15)$   
B.  $(1, 1)$       D.  $(0, 1)$       F.  $(1, 2)$       H.  $(-12, 1)$
- A point  $(m,5)$ ,  $(m,6)$  and  $(m,-2)$  lie on the line  $x=-8$ , then find the value of  $m$ ?
- Draw the graph of a line that passes through the following points.  
A.  $(2, 3), (1, 3), (-2, 3)$       C.  $(0, 0), (0, 1), (0, -2)$   
B.  $(-1, 0), (-1, -1), (-1, 4)$
- List any three points that lie on the line  $x = -2$
- List any three points that lie on the line  $y = -3$

### Graph of the equation of the form $y=mx$ where $m$ is constant



#### Activity 4.3.3

- If  $y=x$ , then complete the given table below.

x	-3	-2	-1	0	1	2	3
y							
(x,y)							

- Draw the graph of the line  $y=x$  using table values above.

**Steps:** To draw the graph of  $y= mx$

**Step 1:** Choose some value of  $x$  and find corresponding value of  $y$ .

**Step 2:** plot the coordinate of the points in step 1.

**Step 3:** Join all points in step 2 using straight line.

**Example:** Draw the graph

A.  $y = 2x$

B.  $y = -2x$

**Solution:**

A.  $y = 2x$

Choose some values of  $x$ , and determine the corresponding values of  $y$  using the equation  $y = 2x$

choose the value of  $x = -3, -2, -1, 0, 1, 2, 3$

when  $x = -3$ ,  $y = 2(-3) = -6$

when  $x = -2$ ,  $y = 2(-2) = -4$

when  $x = -1$ ,  $y = 2(-1) = -2$


when  $x = 0$ ,  $y = 2(0) = 0$

when  $x = 1$ ,  $y = 2(1) = 2$

when  $x = 2$ ,  $y = 2(2) = 4$

when  $x = 3$ ,  $y = 2(3) = 6$

x	-3	-2	-1	0	1	2	3
y	-6	-4	-2	0	2	4	6
(x,y)	(-3,-6)	(-2,-4)	(-1,-2)	(0,0)	(1,2)	(2,4)	(3,6)



B.  $y = -2x$

Choose some values of  $x$ , and determine the corresponding values of  $y$  using the equation  $y = -2x$

Choose the value  $x = -3, -2, -1, 0, 1, 2, 3$

when  $x = -3, y = -2(-3) = 6$

when  $x = -2, y = -2(-2) = 4$

when  $x = -1, y = -2(-1) = 2$

when  $x = 0, y = -2(0) = 0$


when  $x = 1, y = -2(1) = -2$

when  $x = 2, y = -2(2) = -4$

when  $x = 3, y = -2(3) = -6$

x	-3	-2	-1	0	1	2	3
y	6	4	2	0	-2	-4	-6
(x,y)	(-3,6)	(-2,4)	(-1,2)	(0,0)	(1,-2)	(2,-4)	(3,-6)

$y = -2x$



**Note:**

- The graph of  $y = mx$  passes through 1<sup>st</sup> and 3<sup>rd</sup> quadrant when  $m > 0$ .
- The graph of  $y = mx$  passes through 2<sup>nd</sup> and 4<sup>th</sup> quadrant when  $m < 0$ .
- For any values of  $m$  the graph of  $y = mx$  passes through the origin.
- In the equation of the line  $y = mx$ ,  $m$  is called **slope** of a line.

### Exercise 4.3.3

1. If  $y = 4x$ , then complete the table below and draw the graph of  $y = 4x$

$x$	-3	-2	-1	0	1	2	3
$y$							
$(x, y)$							

2. Which of the following equation of lines passes through 1<sup>st</sup> and 3<sup>rd</sup> quadrant
- A.  $y = 7x$                       C.  $y - 12x = 0$                       E.  $y + 6x = 0$   
 B.  $y = -10x$                       D.  $y - x = 0$                       F.  $y = -\frac{3}{2}x$
3. Draw the graph of the following equations.
- A.  $y = 5x$                       C.  $y - 3x = 0$                       E.  $2y + 6x = 0$   
 B.  $y = -5x$                       D.  $y + x = 0$                       F.  $y = \frac{3}{2}x$
4. If a point (2,8) lie on a line  $y = mx$ , then find the value of  $m$ .

## 4.4. APPLICATIONS OF LINEAR EQUATIONS



### Activity 4.4.1

#### Discuss with your friends

Translate each of the following phrases in to mathematical equations.

- A. A number increased by 5 is 11.  
 B. A number decreased by 5 is 11.  
 C. The product of a number and 5 is 30.  
 D. The sum of a number and 7 is 29.  
 E. Half of a number is 20.  
 F. 10 less than a number is 12.  
 G. 5 more than four times a number is 30.

## Solving word problems

*To solve word problems, follow the following important steps*

- 1. Read the problem carefully*
- 2. Select variables for unknown quantities*
- 3. Write a mathematical equation*
- 4. Solve the equation*
- 5. Interpret the result and write the final answer in words.*
- 6. Check the answer*

**Example 1:** A number increased by 10 is 23. What is the number?

**Solution:**

*Let  $y$  be the unknown number*

*Mathematical equation*  $y + 10 = 23$

*solve the mathematical equation*  $y + 10 = 23$

$$y = 23 - 10$$

$y = 13$   $\Leftarrow$  solution

*Hence the unknown number is 13.*

**Example 2:** 26 less than three times a number is 46. What is the number?

**Solution:**

*Let  $x$  be the unknown number*

*Mathematical equation*  $\Rightarrow 3x - 26 = 46$

*solve the mathematical equation*  $3x - 26 = 46$

$$3x = 46 + 26$$

$$3x = 72$$

$x = 24$   $\Leftarrow$  solution

*Hence the unknown number is 24.*

**Example 3:** A number decreased by 11 is 35. What is the number?

**Solution:**

Let  $x$  be the unknown number

Mathematical equation  $x - 11 = 35$

solve the mathematical equation  $x - 11 = 35$

$$x = 35 + 11$$

$$x = 46 \quad \Rightarrow \text{Solution}$$

Hence the unknown number is 46.

**Example 4:** The sum of One seventh of a number and 15 is 19. What is the number?

**Solution:**

Let  $x$  be the unknown number

Mathematical equation  $\Rightarrow \frac{x}{7} + 15 = 19$

Solve the mathematical equation  $\frac{x}{7} + 15 = 19$

$$\frac{x}{7} = 19 - 15$$

$$\frac{7x}{7} = 7 \times 4$$

$$x = 28 \leftarrow \text{Solution}$$

Hence the unknown number is 28.

### Exercise 4.4.1

1. A number increased by 7 gives 20, what is the number?
2. A number decreased by 3 gives 13. What is a number?
3. The difference between a positive number  $x$  and 3 is 21. What is a number?

4. 15 more than twice a number is 37. What is a number?
5. A number is increased by 7, twice the result is 41. Find the number.
6. If the sum of  $\frac{1}{4}$  number and 15 is 13. What is a number?
7. 8 less than three times a number is 55. What is a number?
8. A number is doubled and the result is increased by 8. If the final result is 36. What is a number?

### Example 5: (Application involving consecutive integers)

Find three consecutive integers whose sum is 156.

#### Solution:

Let  $x$  be the smaller integer, then

The three consecutive integers are  $x, x+1$ , and  $x+2$

Mathematical equation  $x + (x+1) + (x+2) = 156$

Solve the mathematical equation  $x + (x+1) + (x+2) = 156$

$$x + x + 1 + x + 2 = 156$$

$$3x + 3 = 156$$

$$3x = 156 - 3$$

$$3x = 153$$

$$x = 51 \leftarrow \text{Smallest number}$$

Hence the unknown integers are 51, 52 and 53.

### Example 6: (applications involving ages)

The sum of the ages of a man and his wife is 96 years. The man is 6 years older than his wife. How old is his wife?

#### Solution:

- let his wife's age is  $y$ .
- The man's age will be  $y+6$  (he is 6 years older)

- The sum of their age is 96, so

$$y + (y + 6) = 96$$

$$2y + 6 = 96$$

$$2y = 96 - 6$$

$$2y = 90$$

$$y = 45 \quad \text{his wife's age}$$

Hence, the age of his wife is 45 years.

### Example 7:

A farmer has sheep and hen. The sheep and hens together have 100 heads and 356 legs. How many sheep and hens does the farmer have?

### Solution:

Since the total heads is 100, the sheep and hen together are 100 in number.

Let the farmer has  $x$  sheep. And the number hens in this case is  $100 - x$

A sheep has 4 legs, so  $x$  sheep have  $4x$  legs.

A hen has 2 legs, so  $(100 - x)$  hens have  $2(100 - x)$  legs

The total number of legs is 356, so

$$\text{number of legs of sheep} + \text{number of legs of hens} = 356$$

$$4x + 2(100 - x) = 356$$

$$4x + 200 - 2x = 356$$

$$4x - 2x = 356 - 200$$

$$2x = 156$$

$$x = 78$$

Therefore, the farmer has  $x = 78$  sheep and  $100 - x = 100 - 78 = 22$  hens

**Exercise 4.4.2**

1. The sum of three consecutive integers is 345. What are the numbers?
2. The sum of two consecutive even integers is 170. What are the integers?
3. The sum of two consecutive odd integers is 144. What are the numbers?
4. The sum of the ages of a man and his wife is 83. The man is 3 years older than his wife. How old is a man and his wife?
5. In a class there are 45 students. The number of girls is 1.5 times the number of boys. How many boys and how many girls are there in the class?
6. There are 26 girls and 22 boys in class room. Every student contributes equal amount of Birr for the renaissance dam (Hidassie Gidib). If Birr 2400 is collected from all students, then what amount of Birr contributed by  
A. Each student?    B. boys?            C. girls?
7. A government has 87 hospitals. If the government plan to build 2 hospitals in a year. How many years will it takes so that the total number of hospitals be 109.
8. In a triangle **ABC**, angle **A** is twice as large as angle **B** and angle **B** is 20 more than angle **C**. What is the measure of each angle? (hint sum of interior angle of a triangle is  $180^\circ$ )

**SUMMARY FOR UNIT 4**

- **A variable** is any letter or a symbol that represent some unknown number or value.
- a constant (a number), a variable, product or quotient of a number and variable is called a **term**.
- In the product of a number and variable, the numerical factor of the term is called **numerical coefficient**.
- **like terms** are terms whose variable and exponent of variable are exactly the same but they may differ in their numerical coefficients. Terms which are not like terms are called **unlike terms**.
- **Algebraic expressions** are formed by using numbers, letters and the

operation of addition, subtraction, multiplication and division.

- Two different algebraic expressions connected by equal sign is called **equation**.
- A linear equation in one variable  $x$  is an equation which can be written in standard form  $ax+b=0$ , where  $a$  and  $b$  are constant numbers with  $a \neq 0$ .
- To transform linear equations to simpler form, you can apply the following rules.
  - i. if  $a=b$ , then  $a+c=b+c$ , where  $a, b$ , and  $c$  are any constant numbers.
  - ii. if  $a=b$ , then  $a-c=b-c$ , where  $a, b$ , and  $c$  are any constant numbers.
  - iii. If  $a=b$ , then  $a \times c = b \times c$ , where  $a, b, c$  are any constant number and  $c \neq 0$
  - iv. If  $a = b$ , then  $\frac{a}{b} = \frac{b}{c}$ , where  $a, b, c$  any constant number and  $c \neq 0$
- To solve linear equations involving brackets, remember the following key properties. For any numbers  $a, b$  and  $c$ 

$$a+(b+c)=a+b+c \qquad a-(b-c) = a-b+c \qquad a(b-c)=ab - ac$$

$$a-(b+c) = a-b-c \qquad a(b+c) = ab+ac$$
- Points on Cartesian plane are described by two numbers  $(a, b)$  that are called coordinates.
  - i. The first number  $a$ , is the horizontal position of the point from the origin. It is called  $x$ -coordinate (abscissa).
  - ii. The second number  $b$  is the vertical position of the point from the origin. It is called  $y$ -coordinate (ordinate).
- The graph of the equation of the line  $x=a$ ,  $a$  is constant is a line parallel to  $y$ -axis at a distance of  $a$  unit from the origin.
- The graph of the equation of the line  $y=b$ ,  $b$  is constant is a line parallel to  $x$ -axis at a distance of  $b$  unit from the origin.
- The graph of  $y=mx$  passes through 1st and 3rd quadrant when  $m>0$ .
- The graph of  $y=mx$  passes through 2nd and 4th quadrant when  $m<0$ .
- In the equation of the line  $y=mx$ ,  $m$  is called slope of a line.
- For any values of  $m$  the graph of  $y=mx$  passes through the origin.
- In solving word problems follow the following steps
  - i. Read the problem carefully

- ii. Select variables for unknown quantities
- iii. Write a mathematical equation
- iv. Check the answer

### Review Exercise for unit 4

#### I. Write True for the correct statements and False for the incorrect statements.

- For any integers  $a, b$  and  $c$ ,  $a(b+c) = ab+ac$ .
- An algebraic expression  $2x^2 + 2xy$  is trinomial algebraic expression.
- $3a^2 b$  and  $3ab^2$  are like terms.
- If  $a=b$ , then  $a-b= 0$
- The equation of the line  $x+y=0$  is vertical line.
- 4 is slope of equation of line  $y-4x=0$ .
- The numerical coefficient of  $3xy^2$  is 2.
- If  $c=d$ , then  $a+c=d+a$ .

#### II. Choose the correct answer from the given alternatives.

- Which one of the following pair terms are like terms?  
A.  $3x^2$  and  $3x$     B.  $-x^2$  and  $-x$     C.  $2ab^2$  and  $2a^2b$     D.  $-4xy$  and  $9yx$
- Which of the following is equivalent to the equation  $-3(2x-1)=2x+12$ ?  
A.  $-6x+3=2x+12$     B.  $-8x=8$     C.  $x=-1$     D.  $-6x=2x-4$
- The solution of  $\frac{x}{2} - x = \frac{3}{8}x - 11$  is  
A.  $\frac{7}{2}$     B.  $\frac{7}{11}$     C.  $\frac{88}{7}$     D.  $-\frac{11}{7}$
- Which of the following value of  $x$  satisfies the equation  $2x=3x-1$ ?  
A. -1    B. 1    C. 0    D. 2
- Which of the following is the slope of the line  $y-3x=0$ ?  
A. -3    B. 3    C.  $\frac{1}{3}$     D. has no slope
- The sum of the age of a girl and her brother is 34. The girl is 8 years older than her brother, how old is her brother?  
A. 21    B. 13    C. 11    D. 23

15. Which one of the following equations is equation of horizontal line?

- A.  $y=4x$    B.  $y+1=0$    C.  $x-2=0$    D.  $x=0$

### III. Work out problems

16. Write the following word phrases in to algebraic expression

- A. Six less than a number  
 B. Nine more than twice of a number  
 C. A number increased by five  
 D. 35 more than half of a number

17. Simplify the following algebraic expressions

- A.  $-5x+4-21x-21$    C.  $2(6y-x)-10y+2(y+x)$   
 B.  $5x-(3x-1)-6(7-x)$

18. Solve each of the following linear equations.

- A.  $4x+36=86-8x$    E.  $\frac{2x+7}{3} - \frac{x-9}{2} = \frac{5}{2}$   
 B.  $12x-8+2x-17=3x-4-8x+74$    F.  $\frac{x+3}{6} - \frac{x-5}{4} = \frac{3}{8}x$   
 C.  $2(6y-18)-102=78-18(y+2)$    G.  $\frac{x-2}{3} - \frac{x+3}{8} = \frac{5x}{6} - 6$   
 D.  $7(x+26+2x)=5(x+7)$

19. Draw the graph of each of the following equation of the line on the same Cartesian coordinate plane

- A.  $y+4x=0$    B.  $y+1=0$    C.  $x=0$    D.  $y=x$

20. Find the equation of the line passes through the origin with slope 4.

21. What is the slope of a line  $y = \frac{x}{2}$

22. 16 less than four times a number is 12, what is a number.

23. The sum of one fifth of a number and 12 is 30.

24. I think a number. If I subtract 8 from it and multiply the difference by 3, the result is 21. What is a number?

25. Bezawite is 8 years older than her sister. How old is Bezawit if the sum of their ages is 32.

26. In mathematics final exam Kaliab scord 10 more than twice the lowest score. If Kaliab scored 46 points, what was the lowest score?



# UNIT 5

# PERIMETER AND AREA OF PLANE FIGURES

## *Learning Outcomes:*

**At the end of this unit, learners will be able to:**

- ↪ classifies the different kinds of triangles.
- ↪ constructs and describe properties of four- sided figures
- ↪ find the perimeter of triangle, parallelograms, trapezium, rhombus and composite shapes
- ↪ derive formula for area of triangle, parallelograms, trapezium and rhombus.
- ↪ calculate areas of triangle, parallelograms, trapezium, rhombus and composite shapes
- ↪ apply the concept of area and perimeter of Plane figures in solving real life problem

## *Introduction*

In this unit you will learn the different kinds of geometric figures and their properties such as triangle, rectangle, square, rhombus, and trapezium. In addition to their properties you will learn, how to calculate the area and perimeter of these geometric figures.

The unit has five sections. The first section deals with revision of triangles, second section deals with four - sided figures, the third section deals with Perimeter and Area of four-sided figures, the fourth section deals with Perimeters and Areas of triangles and the last section deals about application of the concept of area and perimeter of Plane figures in solving real life problems.

## 5.1. REVISION OF TRIANGLES

### *Learning Outcomes:*

At the end of this unit, learners will be able to:


- classifies the different kinds of triangles.






### *Activity 5.1.1*

#### *Discuss with your friends/partners*




1. Define triangle
2. Name triangle below.



3. Classify the following triangles based on their sides.




4. Classify the following triangle based on their angles



5. Draw a triangle and measure the interior angle of this triangle by using protractor. Find the sum of all interior angles?

### Definition 5.1:

**A triangle** is a simple closed plane figure made of three-line segments.



- The line segments forming the triangle are **sides** of the triangle.  
 $\overline{AC}$ ,  $\overline{AB}$  and  $\overline{BC}$  are the sides of the triangle..
- The point where the sides of the triangle meet is called **vertex** (plural form vertices) A, B and C are the vertices of the triangle.
- Angles of a triangle are formed by two sides of the triangle meeting at a point (vertex).
- The interior angles of the above triangle are:
  - The angle at vertex A,  $\angle BAC$  or  $\angle A$
  - The angle at vertex B,  $\angle ABC$  or  $\angle B$
  - The angle at vertex C,  $\angle BCA$  or  $\angle C$
- The sum of interior angles of a triangle is  $\angle A + \angle B + \angle C = 180^\circ$
- A triangle is named by using its vertices.

The above triangle is named as triangle ABC, symbolically,  $\triangle ABC$ . It can also be named using any other sequence of vertices like triangle BCA, symbolically,  $\triangle BCA$ .

### Note:

- A triangle has three sides, three angles, and three vertices.
- The sum of all interior angles of a triangle is always equal to  $180^\circ$ .

## Types of triangles

Triangles can be classified in 2 different ways:

- Classification of triangle according to interior angles
- Classification of triangle based on side length

## Classification of triangle based on side length


Based on side length, triangles are classified in to three:

1. Scalene Triangle
2. Isosceles Triangle
3. Equilateral Triangle

### 1. Scalene Triangle

A triangle in which all the three sides are unequal in length is called a **scalene triangle**.


All sides are different in length,  
so  $\triangle ABC$  scalene triangle



### 2. Isosceles Triangle:

A triangle in which two of its sides are equal is called **isosceles triangle**.

In triangle  $\triangle PQR$ ,  $PQ = PR = 6$  cm,  
then  $\triangle PQR$  is an isosceles triangle



### 3. Equilateral Triangle:

A triangle in which all its three sides are equal in length is called an **equilateral triangle**.

In triangle given above  
 $LM = MN = NL = 5.5$  cm, then  
 $\triangle LMN$  is equilateral triangle.



## Classification of a triangle according to interior angles

Based on the measure of interior angles, triangles are classified in to three:

1. Acute Angled Triangle
2. Right-Angled Triangle

### 1. Acute Angle Triangle

A triangle with all three angles less than  $90^\circ$  is an **acute angle triangle**.




Figure 5.1.3 acute angle triangle

In the above triangle ABC, measure of all angles are less than  $90^\circ$  (acute angle), so triangle ABC is acute angle triangle

### 2. Right-Angle Triangle

A triangle with one angle measuring exactly  $90^\circ$  is a right-angle triangle.

- The other two angles of a right-angle triangle are acute angles.
- The side opposite to the right angle is the largest side of the triangle and is called the **hypotenuse**.




Figure 4.1.2 right angle triangle

**Example:** In triangle ABC,  $\angle ACB = 90^\circ$  is right angle, so triangle ABC is **right angle triangle**.

### 3. Obtuse angle triangle

An **obtuse angle triangle** is a triangle with one obtuse angle (greater than  $90^\circ$ ) and two acute angles.




Figure 4.1.3 obtuse angle triangle

**Example:**

In triangle ABC,  $96^\circ$  is obtuse angle, then Triangle ABC is obtuse angle triangle.

**Exercise 5.1.1**

1. Fill in the blanks:

- A. The triangle in which all its sides are equal is called \_\_\_\_\_.
- B. \_\_\_\_\_ is a triangle in which all its sides are different length.
- C. Each angle of equilateral triangle is \_\_\_\_\_.
- D. \_\_\_\_\_ is a triangle with two equal sides.
- E. In obtuse angled triangle one angle is \_\_\_\_\_ than  $90^\circ$ .
- F. Each angle of an acute triangle is \_\_\_\_\_ than  $90^\circ$ .
- G. The sum of interior angles of a triangle is equal to \_\_\_\_\_.

2. Consider the following triangle.

- A. List the sides, angles and vertices of the triangle

- B. Name the triangle



3. Classify the following triangle:

- A. Sides of triangle are 4 cm, 4 cm and 7 cm
- B. Angles of triangle is  $90^\circ$ ,  $60^\circ$  and  $30^\circ$
- C. Angles of triangle is  $110^\circ$ ,  $40^\circ$  and  $30^\circ$
- D. Sides of triangle are 5 cm, 13 cm and 12 cm
- E. Angles of triangle is  $60^\circ$ ,  $60^\circ$  and  $60^\circ$

**5.2. FOUR - SIDED FIGURES****Learning Outcomes:**

At the end of this unit, learners will be able to:

- ✎ *construct and describe properties of four-sided figures*



### Activity 5.2.1

Discuss with your friends/partners:



1. considering the geometric figure below and answer the following question

A. The geometric figure has \_\_\_\_ sides

B. Name all its sides.

C. Name all its vertices.

D. Name all its interior angles.

E. The geometric figure is called \_\_\_\_.

F. Name the geometric figure using the four vertices.

G. The vertex opposite to D is \_\_\_\_

H. Name the adjacent vertex to D.

I. Name the opposite side to  $\overline{AD}$ .

J. Name the adjacent side to  $\overline{AD}$ .

#### Definition:

**A quadrilateral** is a four-sided geometric figure bounded by line segments.

- A quadrilateral is named by using its four vertices in clock wise or anti clock wise direction.
- The above quadrilateral is named as quadrilateral ACDB or BACD or CDBA and so on, but it cannot be named as ABCD or ADCB.
- The sides of the quadrilateral ABDC above are  $\overline{AB}$ ,  $\overline{AC}$ ,  $\overline{CD}$ , and  $\overline{BD}$

- i. Opposite side to  $\overline{AB}$  is  $\overline{CD}$
  - ii. Opposite side to  $\overline{AC}$  is  $\overline{BD}$
  - iii. The adjacent sides to  $\overline{AB}$  are  $\overline{AC}$  and  $\overline{BD}$
- The point where the sides of a quadrilateral meet is called Vertex (plural form vertices). The vertices of the quadrilateral ABDC above are A, B, D and C.
  - i. The opposite vertex to A is D.
  - ii. The vertexes adjacent to A are C and B.
  - The angles formed by adjacent side of the quadrilateral are **interior angles** of the quadrilateral. The interior angles of quadrilateral ABDC above are  $\angle A$ ,  $\angle B$ ,  $\angle D$  and  $\angle C$
  - i. The opposite angle to  $\angle A$  is  $\angle D$ .
  - ii. The angle adjacent to  $\angle A$  are  $\angle C$  and  $\angle B$ .
  - line segment that connects two opposite vertices of the quadrilateral is called **diagonal**. The diagonals of quadrilateral ABDC above are  $(\overline{AD})$  and  $(\overline{CB})$ .

### Construction and properties of trapezium




#### Activity 5.2.1

#### Discuss with your friends/partners:

Consider the quadrilateral below and answer the following questions

1. Which sides of this quadrilateral are parallel?
2. What do we call such type of quadrilateral?



3. Measure  $\angle A$ ,  $\angle B$ ,  $\angle D$  and  $\angle C$  calculate


i.  $\angle A + \angle D =$  \_\_\_\_\_

iii.  $\angle A + \angle B + \angle C + \angle D =$  \_\_\_\_\_

ii.  $\angle C + \angle B =$  \_\_\_\_\_

### Definition:

**A trapezium** is a special type of quadrilateral in which exactly one pair of opposite sides are parallel.




- The parallel sides of trapezium are called the **bases** of trapezium

In the above trapezium, the parallel sides  $\overline{AB}$  and  $\overline{DC}$  are **bases**.

- The distance between the bases is called the **height (or altitude)** of the trapezium.  $\overline{AE}$  is the **height**.

- The non-parallel sides of the trapezium are called **legs** of the trapezium. The non-parallel sides  $\overline{AD}$  and  $\overline{BC}$  are **legs**.



*Isosceles trapezium*

- If the legs of trapezium are congruent, then trapezium is called **isosceles** trapezium.

### Construction of trapezium

- To perform geometric construction, you need tools like:

- Ruler                      - Compass                      - Protractor

**Ruler** is used to construct line or line segments.

**Compass** is used to construct circles or arcs.

**Protractor** is used to measure or construct angles.



**Example:** construct a trapezium ABCD using the information given below.

Given:  $(\overline{AB}) \parallel (\overline{CD})$ ,  $AB = 8\text{cm}$ ,  $BC = 5\text{cm}$ ,  $m(\angle A) = 60^\circ$ ,  $m(\angle B) = 85^\circ$ ,

**Solution:**


**Step 1:** draw a line segment  $AB = 8\text{cm}$




**Step 2:** construct  $m(\angle A)$  and  $m(\angle B)$  with the given measures.



**Step 3:** Mark point C on the side of such that  $\angle B$  such that  $BC = 5\text{cm}$



**Step 4:** Draw a line through C parallel to AB, intersecting the side of at  $\angle D$



### Exercise 5.2.1

1. Fill in the blank space with the correct answer.
  - A. Four-sided geometric figure is called \_\_\_\_\_
  - B. A line segment that joins opposite vertex of a quadrilateral is \_\_\_\_\_
  - C. The point where the sides of a quadrilateral meet is called \_\_\_\_\_

D. The tools used for geometric construction are \_\_\_\_\_, \_\_\_\_\_ and \_\_\_\_\_

2. Which one of the following is **not** a trapezium?

3.



4. Considering the quadrilateral below answer the following questions


A. the opposite side to FG is \_\_\_\_\_

B. the adjacent sides to HG are \_\_\_\_\_, \_\_\_\_\_

C. the opposite angle to  $\angle F$  is \_\_\_\_\_

D. The quadrilateral is named as \_\_\_\_\_

E. the diagonal of the quadrilateral are line segments joining \_\_\_\_\_ and \_\_\_\_\_ or \_\_\_\_\_ and \_\_\_\_\_




5. construct a trapezium EFGH using the information given below.

Given:  $(\overline{EF}) \parallel (\overline{GH})$ ,  $EF = 10\text{cm}$ ,  $FG = 6\text{ cm}$ ,  $m(\angle E) = 30^\circ$ ,  
 $m(\angle F) = 80^\circ$ ,

### Construction and properties of parallelogram

#### Definition:

*A parallelogram is a quadrilateral in which both pair of opposite sides are parallel.*




*In the figure  $(\overline{AB}) \parallel (\overline{CD})$  and  $(\overline{AD}) \parallel (\overline{BC})$ , thus ABCD is a parallelogram.*

### Construction of parallelogram


- Construct a parallelogram ABCD using the information given below  
Given:  $(\overline{AB}) \parallel (\overline{CD})$  and  $AB=6\text{cm}$ ,  $BC=4\text{cm}$ ,  $m(\angle A)=80^\circ$ ,  $m(\angle B)=100^\circ$

**Solution:**

**Step 1:** Draw a line segment  $AB=6\text{cm}$




**Step 2:** construct  $\angle A$  and  $\angle B$  so that  $(\angle A)=80^\circ$  and  $m(\angle B)=100^\circ$



**Step 3:** mark point C on side of  $\angle B$

such that  $BC=4\text{cm}$ . And draw a line through C parallel to AB so that it meets the side of  $\angle A$  at point D.

Finally, the parallelogram ABCD satisfying the given condition is



*Using the constructed parallelogram above answer the following questions.*

- Measure  $\angle D$  and  $\angle C$ . compare  $\angle D$  with  $\angle B$ , and  $\angle A$  with  $\angle C$  From the result in 1
  - Opposite angles of a parallelogram are \_\_\_\_\_

2. Find the value of

i.  $m(\angle A) + m(\angle B) = \underline{\hspace{2cm}}$

iii.  $m(\angle C) + m(\angle D) = \underline{\hspace{2cm}}$

ii.  $m(\angle B) + m(\angle C) = \underline{\hspace{2cm}}$

iv.  $m(\angle A) + m(\angle D) = \underline{\hspace{2cm}}$

From the result in 2:

- Consecutive (or adjacent) angles of a parallelogram are  $\underline{\hspace{2cm}}$

3. Measure side AD and DC. Compare AD with BC and AB with DC.

From the result in 3:

- Opposite sides of a parallelogram are  $\underline{\hspace{2cm}}$

4. Construct diagonals AC and DB, so that they intersect at O.

Compare the length of DO with OB and AO with OC.

From the result in 4:

- The diagonals of parallelogram  $\underline{\hspace{2cm}}$  each other.

From the activity above the properties of parallelogram is summarized as follows:

1. Opposite sides of parallelogram are congruent.

*i.e.  $AD = BC$  and  $AB = DC$*

2. Opposite angles of a parallelogram are congruent.


*i.e.  $m(\angle A) = m(\angle C)$  and  $m(\angle B) = m(\angle D)$*

3. Consecutive (or adjacent) angles of a parallelogram are supplementary.

*i.e.  $m(\angle A) + m(\angle B) = 180^\circ$  and  $m(\angle C) + m(\angle D) = 180^\circ$*

4. The diagonals of parallelogram bisect each other.

*i.e.  $AO = OC$ , and  $DO = OB$*



**Note:**

**Bisect** means “divides exactly in to two equal parts

**Supplementary** angles are angles whose sum is  $180^\circ$

**Congruent** means equal in length.

## Construction and properties of special parallelograms

Rectangle, Rhombus and square are special parallelograms. They all satisfy the properties of parallelogram.

### A. Rectangle

#### Definition:

*rectangle is a parallelogram with all its angles are right angle.*




Fig. rectangle

### Construction of Rectangle

Construct a rectangle PQRS using the information given below

Given:  $\overline{PQ} \parallel \overline{RS}$  ,  $PQ=6\text{cm}$ ,  $QR=7\text{cm}$ ,  $m(\angle P)=90^\circ$


#### Solution:

**Step 1:** construct a line segment PQ with length 6 cm.




**Step 2:** construct

$m(\angle P)=90^\circ$  and  $m(\angle Q)=90^\circ$



**Step 3:** mark point R and S, such that  $PS=QR=7\text{cm}$  and connect RS parallel to PQ



*Using the constructed rectangle above answer the following questions.*

1. Does it satisfy all properties of parallelogram?
2. Compare the diagonals PR and SQ.

The properties of rectangle are summarized as follows:

1. Rectangle satisfies all properties of parallelogram.
2. The diagonals of rectangle are equal in length and bisect each other.
3. All angles of rectangle are right angle.


**Note:**

1. *Right angle is an angle that measures  $90^\circ$*
2. *All rectangles are parallelogram, but all parallelograms are not rectangles*
3. *A quadrilateral with congruent diagonals is not necessarily rectangle*
4. *A parallelogram with congruent diagonals is rectangle.*

## B. Rhombus

### Definition:

*A rhombus is a parallelogram in which all its sides are congruent.*



### Properties of rhombus

- i. *All properties of parallelogram are properties of rhombus.*
- ii. *All sides of rhombus are congruent (i.e.  $AB=BC=CD=AD$ ).*
- iii. *The diagonals of rhombus are perpendicular to each other (i.e.*

$AC \perp DB$ )

iv. The diagonal of rhombus bisects the angles at the vertices. For example, in the above rhombus  $\angle CDA$  is bisected by the diagonal  $DB$ , so that  $m(\angle CDB) = m(\angle ADB)$ .


**Note:**

1. All rhombs are parallelogram, but all parallelograms are not rhombus.
2. The diagonals of rhombus are perpendicular and bisect each other.

### C. Square

#### Definition:

*square is a parallelogram with four congruent sides and four right angles.*




#### Properties of square

1. Square satisfies all properties of parallelogram, rectangle and rhombus.
2. The diagonals of square are:
  - Perpendicular to each other
  - Bisect each other and
  - Congruent
3. The diagonals bisect the angles at the vertices. Hence, the diagonal forms  $45^\circ$  with its side.

### D. Kite

#### Definition:

*Kite is a quadrilateral that has two pairs of consecutive congruent sides, but opposite sides are not congruent.*





### Properties of kite

1. The diagonals of kite are perpendicular to each other.

But they do not bisect each other.

2. One pair of opposite angles of kite are congruent.



### Exercise 5.2.2

- Write true if the statement is correct and write false if the statement is wrong.
  - All rectangles are parallelogram.
  - The diagonals of rhombus are congruent
  - The opposite sides of kite are congruent
  - All squares are rhombus
  - A quadrilateral with congruent diagonals is necessarily rectangle
  - All trapeziums are quadrilateral
- Fill in the blank space with the correct answer
  - A quadrilateral with both pair of opposite sides are parallel is \_\_\_\_\_
  - A quadrilateral with adjacent sides congruent, but opposite sides are not congruent is \_\_\_\_\_
  - A rhombus with congruent diagonals is \_\_\_\_\_
  - A quadrilateral with exactly one pair of opposite sides parallel is \_\_\_\_\_
  - A parallelogram with all four angles are right angle, but the diag-

onals are not perpendicular is \_\_\_\_\_

F. A parallelogram in which its diagonals are congruent, perpendicular and bisect each other is \_\_\_\_\_

3. Answer the following questions


- A. Describe similarity and difference of rhombus and kite.
- B. Describe similarity and difference of square and rectangle.

### 5.3. AREA AND PERIMETER OF TRIANGLE

- solve routine and non-routine problems involving Perimetre and area of triangle.

#### A. Area of right angled triangle

The formula of area of right angled triangle is derived from area of rectangle. The diagonal of a rectangle divides the rectangle in to two equal right-angled triangles.




**Area of Rectangle**  $A = bh$

**The area of triangle is half of the area of Rectangle, so the area of triangle is:**  $A = \frac{1}{2}bh$


#### B. Area of acute angled triangle

to calculate area of such triangle, draw perpendicular Line from one of the vertices to the opposite side.



The perpendicular line segment from the vertex to the opposite side is called altitude or **height(h)** and the opposite side is the **base(b)**.

AD is height(h) and BC is base(b). The area of  $\triangle ABC$  is the sum of area of Area of  $\triangle ABD$  and  $\triangle ADC$



Area of  $\triangle ABC$  = area of  $\triangle ABD$  + area of  $\triangle ADC$

$$A = \frac{1}{2}BD \times AD + \frac{1}{2}AD \times DC$$


$$A = \frac{1}{2}AD \times (BD + DC)$$

$$A = \frac{1}{2}AD \times BC \dots\dots (BD + DC = BC)$$

$$A = \frac{1}{2}bh \dots\dots \text{where AD is height (h) and BC is base(b)}$$

### C. Area of obtuse angled triangle

to calculate area of obtuse angled Draw height from vertex A to side Extension of base BC



Area of  $\triangle ABC$  is equal to area of  $\triangle ABD$  minus area of  $\triangle ADC$ .

*Area of  $\triangle ABC$  = area of  $\triangle ABD$  - area of  $\triangle ADC$ .*

$$A = \frac{1}{2}BD \times AD - \frac{1}{2}AD \times DC$$


$$A = \frac{1}{2}AD \times (BD - DC)$$

$$A = \frac{1}{2}AD \times BC \dots\dots (BD - DC = BC)$$

$$A = \frac{1}{2}bh \dots\dots \text{where AD is height (h) and BC is base(b)}$$

Note:


The area of triangle whose base b and altitude(height) to this base h is given by  $A = \frac{1}{2}bh$






**Note**

If the length of sides of a triangle are  $a, b$  and  $c$ , then the perimeter  $p$  of the triangle is

$$P = a + b + c.$$



**Example 1:** Calculate the area of the following triangles



**Solution:**

A.  $A = \frac{1}{2} b h$

$$A = \frac{1}{2} \times 7 \text{ cm} \times 4 \text{ cm}$$

$$A = \frac{28 \text{ cm}^2}{2}$$

$$A = 14 \text{ cm}^2$$

B.  $A = \frac{1}{2} b h$

$$A = \frac{1}{2} \times 9 \text{ cm} \times 5 \text{ cm}$$

$$A = \frac{45 \text{ cm}^2}{2}$$

$$A = 22.5 \text{ cm}^2$$

C.  $A = \frac{1}{2} b h$

$$A = \frac{1}{2} \times 10 \text{ cm} \times 9 \text{ cm}$$


$$A = \frac{90 \text{ cm}^2}{2}$$

$$A = 45 \text{ cm}^2$$

**Example 2:**

Using the figure below, calculate

- the area of rectangle
- the area of shaded region
- the area of unshaded region



**Solution:**

A.

$$\begin{aligned}\text{Area of rectangle} &= bh \\ &= 60\text{cm} \times 20\text{cm} \\ &= \underline{1200\text{cm}^2}\end{aligned}$$

B.


$$\begin{aligned}\text{Area of shaded part} \\ A &= \frac{1}{2}bh \\ A &= \frac{1}{2} 60\text{cm} \times 20\text{cm} \\ &= \underline{600\text{cm}^2}\end{aligned}$$

C.

$$\begin{aligned}\text{Area of unshaded part} &= \\ &= \text{area of rectangle minus area of} \\ &= \text{shaded part} \\ &= 1200\text{cm}^2 - 600\text{cm}^2 \\ &= \underline{600\text{cm}^2}\end{aligned}$$

**Example 3:** what is the area and perimeter of the triangle below

**Solution:**



$$\begin{aligned}A &= \frac{1}{2}bh. \\ A &= \frac{1}{2} 6\text{cm} \times 8\text{cm} \\ A &= \frac{48}{2} \\ A &= \underline{24\text{cm}^2}\end{aligned}$$

Perimeter of the triangle is sum of each side:

$$P = 8\text{cm} + 10\text{cm} + 6\text{cm} = \underline{24\text{cm}}$$

**Example 4:** the area of a triangle is  $64\text{cm}^2$ . If the base is 16 cm long, then calculate the height of the triangle.

**Solution:**

$$A = 64\text{cm}^2, b = 16\text{cm}, h = ?$$

$$A = \frac{1}{2}bh.$$

$$64\text{cm}^2 = \frac{1}{2} 16\text{cm} \times h$$

$$= (64\text{cm}^2) = (8\text{cm} \times h)$$


$$\frac{8h}{8} = \frac{64\text{cm}^2}{8\text{cm}}$$

$$h = \underline{8\text{cm}}$$

**Example 5:** based on the triangle below answer the following questions. If AD 12cm, BC=8cm and AC =6cm then calculate

A. Area of  $\triangle ABC$

B. the length BE



**Solution:**

A.  $A(\triangle ABC) = \frac{1}{2}bh$

Using base BC and height AD

$$= \frac{1}{2} \times BC \times AD \dots$$

$$= \frac{1}{2} \times 8\text{cm} \times 12\text{cm}$$

$$= \underline{48\text{cm}^2}$$

B. Using the other base and height, area of  $\triangle ABC$  is

$$A(\triangle ABC) = \frac{1}{2}bh$$

Using base AC and height BE

$$A(\triangle ABC) = \frac{1}{2} \times AC \times BE \dots$$



$$48\text{cm}^2 = \frac{1}{2} \times 6\text{cm} \times BE$$

$$\frac{48\text{cm}^2}{3\text{cm}} = \frac{3\text{cm} \times BE}{3\text{cm}}$$


$$BE = \underline{16\text{cm}}$$

**Exercise 5.3.1:**

1. Calculate the area and perimeter of the following triangles.




2. The area of a triangle is 63 . If its height is 9cm long, then calculate the length of its base.
3. The figure below represents a wall of a certain building. Find the area and perimeter of the wall



4. Based on the figure below, answer the following questions

- A. Calculate the area of shaded region
- B. Calculate the area of unshaded part



## 5.4. PERIMETERS AND AREAS OF FOUR SIDED FIGURES

### Learning Outcomes:

At the end of this unit, learners will be able to:

- Solve routine and non-routine problems involving Perimetre and area of four sided figures.
- Solves routine and non-routine problems involving area of composite figures formed by any two or more figures such as triangle, square, and rectangles.

### Introduction


In grade 6 you have learnt about perimeter and area of different closed figures. In this section you will also learn how to calculate, the area and perimeter of rectangle, square, parallelogram, trapezium, rhombus and kite. In order to have an obvious understanding about the topics you need to follow the following activities.

#### Activity 5.4.1

#### Discuss with your friends/partners

1. Take a metre scale and measure the length of all corners of your classroom. Tell your classmates the value you get.
2. Multiply only the value of length and width. What do you think about the value you get?
3. Considering the following rectangle answer the following questions?

A. What is the area of the rectangle above in square units?



*B. What is the perimeter of the rectangle?*


*4. What is area of a closed region?*

*5. What is perimeter of a closed region?*

*6. How do you calculate area and perimeter of a rectangle?*

### 1. Area and perimeter of rectangle

- ✓ Area of rectangle is the product of its base and height.



***Area of rectangle = base  $\times$  height***

$$A = b \times h$$

- ✓ The perimeter of rectangle is calculated as:


$$\text{perimeter} = b + h + b + h$$

$$P = 2b + 2h$$

$$P = 2(b + h)$$

### 2. Area and perimeter of square

- ✓ Square is a rectangle whose base and height are equal.
- ✓ The area of square whose side length 'S' is




$$A = S^2$$

- ✓ The perimeter of Square is

$$P = 4S$$

**Example:**

calculate the area and perimeter of the rectangle given below



**Solution**

$$A = b \times h$$

$$A = 6\text{cm} \times 8\text{cm}$$

$$A = 48\text{cm}^2$$

$$P = 2(b + h)$$

$$P = 2(6 + 8) = 2 \times 14$$

$$P = 28\text{cm}$$

**Example:**

calculate the area and perimeter of square whose side length is 10cm

**Solution:**

$$S = 10\text{m}, \quad A = ?, \quad P = ?$$

$$A = S^2$$

$$A = (10\text{m})^2$$

$$A = 100\text{m}^2$$

$$P = 4S$$

$$P = 4 \times 10\text{m}$$

$$P = 40\text{m}$$



**3. Perimeter and area of parallelogram**



**Activity 5.4.2**


**Discuss with your friends/partners**

1. What is parallelogram?
2. list its properties of parallelogram
3. Consider the parallelogram below, where  $PQ$  is base( $b$ ) and the altitude( $h$ ) from  $S$  to  $PQ$



- i. Construct altitude from  $Q$  to  $SR$ , and diagonal  $SQ$  as in  $b$  above
- ii. What is the area of  $\triangle PQS$  in terms of  $b$  and  $h$
- iii. What is the area of  $\triangle QRS$  in terms of  $b$  and  $h$
- iv. How do you find the area parallelogram using areas of  $\triangle PQS$  and  $\triangle QRS$
- v. From  $a, b, c$ , the formula to calculate area of parallelogram is \_\_\_\_\_.

As you have done in the above activity the formula to calculate area of parallelogram is derived as follows:



$$A(PQRS) = A(\triangle PQS) + A(\triangle QRS)$$

$$A(PQRS) = \frac{1}{2} \times PQ \times h + \frac{1}{2} \times RS \times h$$

$$A(PQRS) = \frac{1}{2}bh + \frac{1}{2}bh$$

$$A(PQRS) = bh$$

**Note:**

1. The area of parallelogram with base  $b$  and altitude  $h$  is given by the formula:


$$A = bh$$

2. The perimeter of parallelogram is calculated by adding all its sides

$$P = PQ + QR + RS + PS$$

$$P = b + a + b + a$$

$$P = 2(a + b)$$



**Example 1:** In the figure given below,  $AP, AQ$  are altitude of the parallelogram  $ABCD$ .

If  $AQ = 4\text{cm}$ ,  $CD = 5\text{cm}$  and  $AP = 8\text{cm}$ , then calculate

- Area of the parallelogram
- Length of  $BC$
- Perimetre of the parallelogram

**Solution:**

A.

$$A = bh$$

using base  $DC$   
And height  $AQ$

$$A = DC \times AQ$$

$$A = 5\text{cm} \times 4\text{cm}$$

$$A = 20\text{cm}^2$$

B.

$$A = bh$$

using base  $BC$  And  
height  $AP$

$$A = BC \times AP$$

$$(20\text{m}^2) = (BC \times 8\text{cm})$$

$$8\text{cm} \quad 8\text{cm}$$

$$BC = 2.5\text{cm}$$

C.

perimeter of the  
parallelogram is:


$$AB = DC = 2.5\text{cm},$$

$$AD = BC = 2.5\text{cm}$$

$$P = AB + BC + CD + AD$$

$$P = 5\text{cm} + 2.5\text{cm} + 5\text{cm} + 2.5$$

$$P = 15\text{cm}$$



**Example 2:**

The area of parallelogram is  $42\text{cm}^2$ . If the base of the trapezium is  $6\text{cm}$ , then find the corresponding altitude.

**Solution:**


$$A = 42\text{cm}^2, \quad b = 6\text{cm}, \quad h = ?$$

$$A = bh$$


$$\frac{(42\text{cm}^2)}{6\text{cm}} = \frac{(6\text{cm} \times h)}{6\text{cm}} \quad h = 7\text{cm}$$

### Exercise 5.4.1


- Calculate the area of the following parallelogram
- ABCD is a parallelogram of area  $18\text{cm}^2$ . Find the length of the corresponding altitudes, if  $AB = 5\text{cm}$ .



3. Calculate the area of the region PTRS, if PQRS is parallelogram



4. In the figure given below, WQ, WP are altitude of the parallelogram WXYZ. If WP = 6 cm, XY = 5 cm and WQ = 8 cm, then calculate
- Area of the parallelogram
  - Length of ZY
  - Perimetre of the parallelogram





#### 4. Perimeter and area of trapezium



#### Activity 5.4.3


*Discuss with your friends/partners*

- What is trapezium?
- Consider the trapezium below



- Construct diagonal AC and altitude from D to base  $AB_2$  (its extension) as in b above.
- What is the area of  $\triangle ADC$  in terms of  $b_2$  and  $h$ .
- What is the area of  $\triangle ABC$  in terms of  $b_1$  and  $h$ .
- The area of trapezium ABCD is the sum of the area of  $\triangle ADC$  and  $\triangle ABC$
- From  $a, b, c$  the formula to calculate area of trapezium is \_\_\_\_\_.

As you have done in the above activity the formula to calculate area of trapezium is derived as follows:



The area of trapezium ABCD is the sum of the area of  $\triangle ADC$  and  $\triangle ABC$

$$(ABCD) = A(\triangle ADC) + A(\triangle ABC)$$

$$= \frac{1}{2} \times b_1 \times h + \frac{1}{2} \times b_2 \times h$$

$$= \frac{1}{2} h (b_1 + b_2)$$

$$= \frac{h}{2} (b_1 + b_2)$$

**Note:**


Area of trapezium with base  $b_1$  and  $b_2$  and altitude  $h$  is given by the

formula:  $= \frac{h}{2} (b_1 + b_2)$

The perimeter of trapezium is the sum of the two base and its legs:

$$\text{perimeter} = AB + BC + DC + AD$$

$$P = b_1 + b_2 + l_1 + l_2$$



**Example 1:** calculate the area and perimeter of the trapezium given below

**Solution:**


the area of trapezium

$$= \frac{h}{2} (b_1 + b_2)$$

$$A = \frac{3}{2} (8 + 16)$$

$$A = \frac{3}{2} \times 24$$

$$A = 36 \text{ cm}^2$$



the perimeter of trapezium is

$$P = b_1 + b_2 + l_1 + l_2$$

$$= 8 + 16 + 5 + 5 \quad P = 34 \text{ cm}$$

**Example 2:** The area of trapezium is  $48 \text{ cm}^2$  and its height is  $4 \text{ cm}$ . If one of the base is  $4 \text{ cm}$  more than the other base, then calculate each base of the trapezium.

**Solution:**

let the smaller base is ,  $x$  and the larger base is  $x+4$ ,  $A = 48 \text{ cm}^2$

$$= \frac{h}{2} (b_1 + b_2)$$


$$A = \frac{4}{2} (x + x + 4)$$

$$48 = 2 (2x+4)$$

$$48 = 4x+8$$

$$4x = 40$$


$$x = 10 \text{ cm}$$



Therefore, the bases of trapezium are  
Smaller base  $x = 10 \text{ cm}$  and larger  
base is  $x+4 = 10+4 = 14 \text{ cm}$ .

**Exercise 5.4.2**



1. Calculate the area and perimeter of the following trapezium



2. The area of trapezium is  $170 \text{ cm}^2$ . If its height and one of the bases are  $17 \text{ cm}$  and  $12 \text{ cm}$  respectively, then calculate the other base of the trapezium.
3. One of the bases of trapezium exceeds the other by  $2 \text{ cm}$ . If the altitude and area of trapezium are  $6 \text{ cm}$  and  $42 \text{ cm}^2$  respectively, then calculate the larger base of the trapezium.


**5. Perimeter and area of rhombus and kite****Activity 5.4.4****Discuss with your friends/partners**

1. Define rhombus and list its properties.
2. Define kite and list its properties
3. Consider the following rhombus and kite



- i. Name the diagonals of the rhombus
- ii. Name the diagonals of kite
- iii. Calculate the area of rhombus by adding area of  $\Delta ABC$  and  $\Delta ADC$
- iv. Calculate the area of kite by adding area of  $\Delta WXY$  and  $\Delta WZY$
- v. Using the above result generalize the formula to calculate area of rhombus and kite.

As you have done in the above activity the formula to calculate area of trapezium is derived as follows



### Area of rhombus

is the sum of area  $\Delta ABC$  and  $\Delta ADC$

$$A = A(\Delta ABC) + A(\Delta ADC)$$

$$A = \frac{1}{2} AC \times BE + \frac{1}{2} AC \times ED$$

$$A = \frac{1}{2} AC(BE + ED)$$


since  $BE + ED = BD$ , then

$$A = \frac{1}{2} AC \times BD$$

where  $AC$  is diagonal 1( $d_1$ )

and  $BD$  is diagonal 2( $d_2$ )

$$A = \frac{1}{2} d_1 d_2$$



### Area of kite

is the sum of area  $\Delta WXY$  and  $\Delta WZY$

$$A = \Delta WXY + \Delta WZY$$

$$A = \frac{1}{2} WY \times NX + \frac{1}{2} WY \times NZ$$

$$A = \frac{1}{2} WY(NX + NZ)$$

since  $NX + NZ = XZ$ , then

$$A = \frac{1}{2} WY \times XZ$$

where  $WY$  is diagonal 1( $d_1$ )

and  $XZ$  is diagonal 2( $d_2$ )

$$A = \frac{1}{2} d_1 d_2$$

**Note:**

*The area of rhombus and kite is given by the formula*


$$A = \frac{1}{2} d_1 d_2, \text{ where } d_1 \text{ and } d_2 \text{ are diagonals.}$$

*The perimeter of rhombus and kite is calculated by adding the length of all the four sides.*

**Example 1:** Calculate the perimeter and area

of the kite given below, if  $WX=6\text{cm}$ ,  $XY=10\text{cm}$ ,

$XZ=8\text{cm}$  and  $WY=16\text{cm}$



**Solution:**

$$A = \frac{1}{2} d_1 d_2$$

$$A = \frac{1}{2} 8\text{ cm} \times 16\text{ cm}$$

$$A = \frac{1}{2} WY \times XZ$$

$$A = 64$$

$$P = WX + WZ + XY + YZ$$


since  $WX = WZ = 6\text{cm}$  And  $XZ = YZ = 8\text{cm}$  then,

$$P = 6 + 6 + 8 + 8 = 18\text{ cm}$$

### Exercise 5.4.3

1. Calculate the area and perimeter of the following rhombus if

$AB=5\text{cm}$ ,  $AC=8\text{cm}$  and  $BD=6\text{cm}$ .



2. Calculate the area of kite, whose diagonals are  $12\text{cm}$  and  $16\text{cm}$ .

3. Calculate the perimeter of kite, whose adjacent sides are  $18\text{mm}$  and  $10\text{mm}$ .

4. The area of rhombus is  $144\text{cm}^2$ . If one of its diagonals is  $18\text{ cm}$  long, then calculate the length of the other diagonal.

## 5.5. CIRCUMFERENCE AND AREA OF A CIRCLE.

### *Learning Outcomes:*

At the end of this unit, learners will be able to:

- *Define circumference and area of a circle*
- *Calculate the circumference and area of a circle.*


### **Activity 5.5.1**

1. Have you ever observed circular object in your locality? If yes, tell your classmates loudly.
2. Can you tell your classmates how to draw a circle?
3. What are the materials you use to draw a circle?
4. Draw a circle and indicate different parts of a circle.

### 5.5.1. Revision on the parts of a circle

In grade 6 you had learnt about parts of circles. It is necessary to note the following parts of a circle to clearly remember what you learnt in your previous class.

1. A circle is the set of all points on a plane that are at equal distance from the center of a given point called *center of a circle*.
2. A chord of a circle is any line segment that joins two points on the circle ( $\overline{AB}$ ) and ( $\overline{CD}$ ).
3. A diameter ( $d$ ) is a line segment that passing through the center and joining any two points on the circle.
4. A radius ( $r$ ) of a circle is the distance from the center to any point on the circle.  $OA$ ,  $OB$  and  $OD$  are radii of circle.




### 5.5.2. Circumference of circle

As you have seen before, the distance all-round the plane figure is perimeter. The analogy of perimeter for a circle is named as circumference of a circle.



#### Activity 5.5.2

1. *Materials needed: ruler, string, scissor and circular objects of various sizes.*
  - A. *Bring a circular object.*
  - B. *Use a ruler to measure the diameter of your circular object. Record your finding.*
  - C. *Wrap a string around the circular object once. Mark the string where it meets itself.*
  - D. *Lay the string out straight, measure and then record the length of the string with your ruler. Can you call the circumference of the circle?*
  - E. *Divide the measure of the circumference obtained in (d) by the measure of the diameter obtained in (b). Record your answer.*
  - F. *Repeat this activity with circular objects of various sizes.*
2. *Compare the results you obtained in (e) and (f). What do you observe or find?*
3. *How is the circumference related to the diameter?*
4. *What is your conclusion?*



From the activity above, you can understand that the ratio of the circumference to the diameter of the circles you considered approaches to a number very closer 3.14 called  $\pi$ .

Therefore, the circumference divided by the diameter of any circle is given as  $\frac{C}{d} = \pi$  (where  $\pi$  is Greek letter (read as pi)). Most of the time the rational numbers we used for approximately  $\pi$  are 3.14 or  $\frac{22}{7}$ .

### Circumference of a Circle

*For any circle with diameter  $d$ , or radius  $r$ , its circumference is given by:  $C = \pi d$  or  $C = 2\pi r$*

**Example:** Find the circumference of a circle with diameter 12m.

**Solution:**

$$C = \pi d \text{ and } d = 12m$$

$$\text{Then, } C = \pi d = \pi \times 12m = 3.14 \times 12m \approx 53.38m,$$

Therefore, the circumference of the circle is  $12\pi m$  or approximately 53.38m.

**Example:** The circumference of the circular basement of the building is 220m. Find the radius of the basement. (Use  $\pi = \frac{22}{7}$ )

**Solution:**

$$\text{Given: } C = 220m \quad \text{and} \quad C = 2\pi r$$

$$220m = (2 \times \frac{22}{7} \times r) = \frac{44}{7}r$$

$$220m \times \frac{7}{44} = \frac{44}{7}r \times \frac{7}{44}$$

$$35m = r$$

Therefore, the radius of the building is 35m.

**Example:** The figure below is formed a semicircle and a straight line. What is its perimeter? (Use  $\pi = \frac{22}{7}$ )

**Solution:**

Diameter of circle,  $d=6m$

Perimeter of a semicircle = Halve of the circumference of a circle

$$= \frac{1}{2} \times \pi \times d$$

$$= \frac{1}{2} \times \frac{22}{7} \times 6m = 11m$$

Perimeter of a semicircle =  $\frac{1}{2}$  of the circumference of a circle + length diameter

$$= 11m + 6m = 18m$$


**5.5.3. Area of a Circle**

From your previous grade lesson you could remember what to mean area of a plane figure. It is the number of square units that cover the plane figure. We can approximate the area of a circle by using square grid. That is, dividing a circle in to small pieces of unit squares (a square of area  $1\text{cm}^2$ ).

Number of squares covered by the quadrant is 39.

$$\text{Area of a quadrant} = 39 \times 1\text{cm}^2 = 39\text{cm}^2$$

$$\text{Area of a circle} = 39\text{ cm}^2 \times 4 = 156\text{cm}^2$$



The following activity will help you find the way of calculating the accurate area of a circle.

**Activity 5.5.3**

**Materials needed:** paper, compass, ruler or straightedge, scissors, pencil.


1. Draw a circle and several radii that separate the circle in to **equal-sized** sections. (Let  $r$  units represent the length of the radius and  $C$  units



represent the circumference of the circle).

2. Cut out each section of the circle.

3. Reassemble the sections in the form of a parallelogram.



4. What is the base length of this “parallelogram”? How about length of its height?

5. Find the area of the parallelogram. Remember area of a parallelogram is:  $A = bh$ .

6. How could you use this formula to find the area of a circle?

- The base of the parallelogram shown on the activity above is half the circumference of the circle

$$\frac{1}{2} C = \frac{1}{2} \times 2\pi r$$

- The height of the parallelogram is the length of the radius. Substitute this information into the formula for the area of a parallelogram.


$$\text{Area of a circle} = \text{area of the parallelogram} = \frac{1}{2} C \times r = \pi r \times r = \pi r^2$$

### Exercise 5.5.1


1. Complete the following table (Use  $\pi = \frac{22}{7}$ )

Radius	Diameter	Circumference	Area
12cm			
		16cm	
			$\frac{22}{7} \text{ cm}^2$
	24cm		

2. If the diameter of a circle is three times as long as the diameter of another circle, explain how the circumference of two circles relates to each other.
3. Find the perimeter and area of the following figure. (Use  $\pi = \frac{22}{7}$ )



4. What happens to the area of a circle if the radius of the circle is doubled?
5. Calculate the perimeter and area of the figure given below.



## 5.6. APPLICATIONS OF PERIMETER AND AREA OF PLANE FIGURES.

### *Learning Outcomes:*

**At the end of this unit, learners will be able to:**

- Apply the concept of perimeter and area of plane figures in real life problems.


In this section you will learn some of the application of area and perimeter in our real world.

The concept of area and perimeter is applied:

- In decorating the wall of a house, so that to estimate the cost of painting painting
- To estimate the fencing material required in fencing a garden.

- To estimate the area of roofing material.
- Making various furniture.
- To estimate carpet size for floor.

**Example 1:** the diagram below shows the floor plan of a hotel. Carpet costs Birr 100 per square meter. How much will it cost to carpet the hotel.



**Solution:**

First calculate the area of the floor

*Area of the floor is the sum of the areas of the two trapeziums.*

$$A = \frac{h}{2}(b_1 + b_2) + \frac{h}{2}(b_1 + b_2)$$

$$A = \frac{3}{2}(4 + 6) + \frac{3}{2}(4 + 6)$$

$$A = 15 + 15 = 30 \text{ m}^2$$


*The cost of  $1 \text{ m}^2$  carpet is Birr 100, so the cost of  $30 \text{ m}^2$  is Birr*

$$30 \times 100 = 3000$$

*Hence, it costs Birr 3000 to carpet the floor.*

**Example 2:** A farmer wants today fence a plot of land shown below

- Calculate the length of fencing material required
- If the cost of 1m fencing material is Birr 250, then calculate the cost to fence the plot of land



**Solution:**

$$A. \quad P = AB + BC + CD + DE + EF + FA \quad BC = DE - AF = 50 - 20 = 30$$

$$P = 40 + 30 + 20 + 50 + 60 + 20$$

$$P = 220 \text{ m}$$


And

$$CD = EF - AB = 60 - 40 = 20$$

**Therefore, 220m of fencing material is required.**

B. The cost of 1m fencing material is Birr 250, so the cost of 220m fencing material is  $\text{Birr } 250 \times 220 = \text{Birr } 55,000$

**Example 3:** The following diagram shows the front part of a building  
If you paint the wall of the building, then



A. calculate the area of the building painted.

B. If it costs Birr200 to paint 1 m<sup>2</sup> wall, then how much it costs to paint the wall?

**Solution:**

A. Note that the area of window and door will not be painted, so the area of wall painted is equal to the area of bigger rectangle minus the area of door and window

$$\begin{aligned} \text{painted area} &= 8 \times 3 - 1 \times 1.5 - 2 \times 1.7 \\ &= 24 - 1.5 - 3.4 = 19.1 \text{ m}^2 \end{aligned}$$

B. it costs Birr200 to paint 1 m<sup>2</sup>. So, to paint 19.1m<sup>2</sup>, it costs  
 $\text{Birr } 200 \times 19.1 = \text{Birr } 3820$

**Example 4:** A class room has length of 9m and width of 6m. The flooring is to be replaced by terazo tiles of size 30cm by 30cm.  
how many terazo tiles are needed to cover the class room.

**Solution:**

$$\begin{aligned} \text{Area of class room, } A &= 9\text{m} \times 6\text{m} = 54\text{m}^2 \\ &= 540,000\text{cm}^2, 1 \text{ m}^2 = 10,000 \text{ cm}^2 \end{aligned}$$

$$\text{Area of terazo tile, } A = 30\text{cm} \times 30\text{cm} = 900\text{cm}^2$$


$$\begin{aligned} \text{Number of terazo tiles} &= \frac{\text{Area of class room}}{\text{Area of terazo tile}} = \frac{540,000 \text{ cm}^2}{900 \text{ cm}^2} \\ &= 600 \end{aligned}$$

Therefore, 600 terazo tiles are required for flooring

### Exercise 5.6.1

1. The length and width of rectangular football field are 100m and 80m respectively. 1 m<sup>2</sup> artificial grass costs Birr 500, then how much it costs to cover the field by artificial grass.
2. The following diagram shows one side of a building. If this part will be covered by HCB (hollow concrete block), then how many HCB is required?


(the size of HCB is 20cm by 40cm, and 1 m<sup>2</sup>=10,000cm<sup>2</sup>)



3. A farmer wants to fence the following plot of land. If no fencing material is required along river side,

Then calculate,

- A. The length of fencing material required
  - B. If the cost of 1 m fencing material is Birr 150, then calculate the cost to fence the land.
4. A rectangular flower bed measures 10m by 6m. It has a path 2m wide around it. Find the area of the path



## SUMMARY FOR UNIT 5

- Triangles based on their sides and angles classified as follows

### Based on sides

1. Scalen triangle
2. Isosceles triangle
3. Equilateral triangle

### Based on angles

1. Acute angled triangle
2. Right angled triangle
3. Obtuse angled triangle

**A quadrilateral** is a four-sided geometric figure bounded by line segments.

- **A trapezium** is a special type of quadrilateral in which exactly one pair of opposite sides are parallel.
- **A parallelogram** is a quadrilateral in which both pair of opposite sides are parallel.
- **rectangle** is a parallelogram with all its angles are right angle.
- **square** is a parallelogram with four congruent sides and four right angles.
- **A rhombus** is a parallelogram in which all its sides are congruent.
- **Kite** is a quadrilateral that has two pairs of consecutive congruent sides, but opposite sides are not congruent.
- Formula to calculate area and perimeter of different geometric figure angles

### 1. Triangle

$$A = \frac{1}{2} bh \text{ and } P = a + b + c$$

### 2. Rectangle

$$A = bh \text{ and } P = 2(b + h)$$

### 3. Squar

$$A = S^2 \text{ and } P = 4S$$

### 4. Rhombus

$$A = \frac{1}{2} d_1 d_2 \text{ and } P = 4S$$

### 5. Kite

$$A = \frac{1}{2} d_1 d_2 \text{ and } P = 2(a + b)$$

### 6. Trapezium

$$\text{and } P = a + b + b_1 + b_2$$

## REVIEW EXERCISE FOR UNIT 5

**Write True for the correct statements and False for the incorrect statements.**

1. If the adjacent sides of a parallelogram are congruent, then the parallelogram is rhombus
2. If the adjacent sides of a parallelogram are congruent, then the parallelogram is square.
3. The sides of rectangle make  $45^\circ$  with its sides.
4. The diagonals of kite bisect each other.
5. If the diagonals of a parallelogram are perpendicular bisector of each other, then the parallelogram is necessarily rhombus.
6. All squares are rhombus.
7. All rectangles are squares.

**Fill in the blank space with the correct answer**

1. A triangle with all its sides are equal is called \_\_\_\_\_
2. A triangle with one right angle is called \_\_\_\_\_
3. A quadrilateral with adjacent sides congruent, but opposite sides are not congruent is called \_\_\_\_\_
4. A parallelogram with all four sides congruent is necessarily \_\_\_\_\_
5. A rhombus with four right angles is \_\_\_\_\_

**Choose the correct answer**

1. The area of triangle with base 4cm and height 3cm is \_\_\_\_\_  
A.  $12\text{cm}^2$       B.  $6\text{ cm}^2$       C.  $7\text{ cm}^2$       D.  $14\text{ cm}^2$
2. The area of rhombus whose diagonals are 12cm and 18cm is \_\_\_\_\_  
A.  $108\text{ cm}^2$       B.  $216\text{ cm}^2$       C.  $80\text{cm}^2$       D.  $160\text{cm}^2$
3. A toilet room has length of 2m and width of 2 m. The flooring is to be tiled by ceramic tiles of size 20cm by 20cm. how many ceramic tiles are needed to cover the class room?

A. 300

B. 150

C. 200

D. 100

4. The area of rectangular field is  $480\text{m}^2$ . If the length of base of the field is  $12\text{m}$ , then the perimeter of the field is \_\_\_\_\_

A.  $40\text{m}$ B.  $52\text{m}$ C.  $104\text{m}$ D.  $80\text{m}$ 


5. The perimeter of kite with its opposite sides are  $5\text{cm}$  and  $12\text{cm}$  is \_\_\_\_\_

A.  $60\text{cm}$ B.  $34\text{cm}$ C.  $17\text{m}$ D.  $30\text{cm}$ 


### Work out

1. Calculate the area and perimeter of the following figures.


A.




B.




C.




D.



E.



F.



# UNIT 6

## CONGRUENCY OF PLANE FIGURES

### *Learning Outcomes:*

At the end of this unit, learners will be able to:

- ⇒ identify congruent triangles by using the tests for congruency (SSS, SAS, ASA).
- ⇒ apply real-life situations in solving geometric problems

### *Introduction*

In this unit you will learn about congruence of geometric figures. The unit has two sections. The first section deals with the congruence of geometric shapes. The main focus of this section is to enable you identifying a given triangle is congruent to the other using congruence test for triangles. In the second section you learn the application of congruence of figures in your daily life.

### **6.1. CONGRUENT OF PLANE FIGURES**

At the end of this unit, learners will be able to:

- ⇒ Explain the concept of congruency of triangles
- ⇒ Check the congruence of given triangles by tracing, cutting and overlapping.
- ⇒ Apply real-life applications of congruency of triangles


## 6.1.1 . Definition and Illustration of Congruent Figures



### Activity 6.1.1

*Discuss with your friends/partners*

1. Which pair of the following line segments are congruent?



2. Which pair of the following circles are congruent?





3. Are the two maps below congruent?



4. Which pair of the following figures are congruent?



5. Which pair of angles are congruent?



6. when do we say that two figures are congruent?

7. When do we say that two-line segments are congruent?

8. When do we say that two circles are congruent?

9. When do we say that two angles are congruent?

**Definition 6.1:**

**Congruent figures** are figures that have the same size and shape. Congruent figures are exact copies of one another.


**Note:**

1. Two figures are congruent, if one figure cover the other completely and exactly.
2. Two-line segments are congruent if they have the same length.
3. Two circles are congruent if they have the same radius.
4. Two angles are congruent if they have the same measure.

**Exercise 6.1.1**

1. Which pair of the following figures are congruent?

A.



B.



C.



D.



2. Write true if the statement is correct and write false if the statement is wrong.
- A. All line segments are congruent
  - B. All circles are congruent
  - C. All rectangles are congruent
  - D. Two circles having the same diameter are congruent
  - E. Two-line segments having the same length are congruent

- F. Two angles that have the same measure are congruent  
 G. If two figures have the same shape, then they are congruent.  
 H. If two triangles are congruent, then they have the same area.

3. Answer the following questions using the following figures



- A. If you put one of the figures above over the other, does it cover completely and exactly?  
 B. What can you say about the figures?

### 6.1.2. Congruency of Triangles



#### Activity 6.1.2

*Discuss with your friends/partners*

1. Consider the following two triangles and measure each side and each angle of the two triangles and fill the following blank space

A. Length of  $AB$  congruent to \_\_\_\_\_


B. Length of  $BC$  congruent to \_\_\_\_\_

C. Length of  $AC$  congruent to \_\_\_\_\_

D.  $\angle A$  is congruent to \_\_\_\_\_

E.  $\angle B$  is congruent to \_\_\_\_\_

F.  $\angle C$  is congruent to \_\_\_\_\_



2. If you put  $\triangle ABC$  on  $\triangle XYZ$ , in such a way that

A.  $AB$  on  $XY$ ,  $BC$  on  $YZ$  and  $AC$  on  $XZ$ , then does  $\triangle ABC$  covers  $\triangle XYZ$  completely?

B. What can you say about the two triangles?

3. If  $\triangle ABC$  is congruent to  $\triangle XYZ$ , then

A. The corresponding side to  $AB$  (the side that will fit or cover side  $AB$  completely) is \_\_\_\_\_

B. The corresponding side to  $AC$  (the side that will fit or cover side  $AC$  completely) is \_\_\_\_\_

C. The corresponding side to  $BC$  (the side that will fit or cover side  $BC$  completely) is \_\_\_\_\_

D. The corresponding angle to  $\angle A$  (the angle that fit  $\angle A$ ) is \_\_\_\_\_

E. The corresponding angle to  $\angle B$  (the angle that fit  $\angle B$ ) is \_\_\_\_\_

F. The corresponding angle to  $\angle C$  (the angle that fit  $\angle C$ ) is \_\_\_\_\_

#### 4. When do we say that two triangles are congruent?

##### Definition 6.2:

**Two triangles are congruent, if their corresponding parts (angles and sides) that match one another are equal.**

##### Note:

1. Two triangles are congruent if they are copies of each other and when you place one triangle on another, they cover each other completely.


2. If  $\triangle ABC$  is congruent to  $\triangle XYZ$ , then symbolically written as  $\triangle ABC \cong \triangle XYZ$

3. If  $\triangle ABC \cong \triangle XYZ$ , then when you place  $\triangle XYZ$  on  $\triangle ABC$  it should

satisfy the following six conditions.

i.  $X$  falls on  $A$ , and  $\angle X \cong \angle A$ ,  
(i.e.  $\angle X$  is congruent to  $\angle A$ )

ii.  $Y$  falls on  $B$ , and  $\angle Y \cong \angle B$ ,  
(i.e.  $\angle Y$  is congruent to  $\angle B$ )




- iii.  $Z$  falls on  $C$ , and  $\angle Z \cong \angle C$ , (i.e.  $\angle Z$  is congruent to  $\angle C$ )
- iv.  $\overline{XY}$  falls along  $\overline{AB}$ , and  $\overline{XY} \cong \overline{AB}$  (i.e.  $\overline{XY}$  is congruent to  $\overline{AB}$ )
- v.  $\overline{YZ}$  falls along  $\overline{BC}$ , and  $\overline{YZ} \cong \overline{BC}$  (i.e.  $\overline{YZ}$  is congruent to  $\overline{BC}$ )
- vi.  $\overline{XZ}$  falls along  $\overline{AC}$ , and  $\overline{XZ} \cong \overline{AC}$  (i.e.  $\overline{XZ}$  is congruent to  $\overline{AC}$ )

**Example 1:** If  $\triangle ABC \cong \triangle FED$ , then find the six congruent corresponding parts of the triangles.

**Solution:**

- |                               |   |   |
|-------------------------------|---|---|
| i. $\angle A \cong \angle F$  | iii. $\angle C \cong \angle D$          | v. $\overline{BC} \cong \overline{ED}$  |
| ii. $\angle B \cong \angle E$ | iv. $\overline{AB} \cong \overline{FE}$ | vi. $\overline{AC} \cong \overline{FD}$ |

**Example 2:** consider the following triangles, congruent angles are marked as indicated, based on the figure answer the following questions



**Solution:**

A. The corresponding part of  $\angle P$  is  $\angle Z$ , The corresponding part of  $\angle Q$  is  $\angle Y$  The corresponding part of  $\angle R$  is  $\angle X$ ,

**Therefore,  $\triangle PQR \cong \triangle ZYX$**

B.  $\triangle QRP \cong \triangle YXZ$

**Example 3:** let  $\triangle ABC \cong \triangle DEF$ , and  $m(\angle A) = 70^\circ$ ,  $DE = 6\text{ cm}$ , then find  $m(\angle D)$  and the length of  $AB$

**Solution:**

Since the triangles are congruent, their corresponding parts are

congruent. So,  $m(\angle A) = m(\angle D) = 70^\circ$  and,  $DE = AB = 6\text{ cm}$

### Exercise 5.1.2

1. Write true if the statement is correct and write false if the statement is wrong

A. If  $\triangle ABC \cong \triangle EDF$ , then  $m(\angle A) = m(\angle D)$

B. All equilateral triangles are congruent

C. If  $\triangle ABC \cong \triangle EDF$ , then  $AB = ED$

D. If  $\triangle ABC \cong \triangle EDF$ , then  $\angle B \cong \angle D$ , then

E. If  $\triangle ABC \cong \triangle EDF$ , then  $\overline{AC} \cong \overline{DF}$

F. Any triangle is congruent to itself.

**Note:**


1.  $AB$  means measure of length of  $(AB)$   
2.  $m(\angle D)$  means measure of  $\angle D$

2. If  $\triangle ZXY \cong \triangle EDF$ , then,

A.  $\triangle XYZ \cong$  \_\_\_\_\_ C.  $\overline{FD} \cong$  \_\_\_\_\_ E.  $\angle D \cong$  \_\_\_\_\_

B.  $m(\angle D) =$  \_\_\_\_\_ D.  $FD =$  \_\_\_\_\_ F.  $\triangle YXZ \cong$  \_\_\_\_\_

3. Considering the following triangles and answer the following questions, congruent angles are indicated with the same mark



A.  $\triangle XYZ \cong$  \_\_\_\_\_

A.  $\triangle SRT \cong$  \_\_\_\_\_

A.  $\triangle ZYX \cong$  \_\_\_\_\_

### 6.1.3 Tests for congruency of triangles (ASA, SAS, SSS)

#### Learning Outcomes:

At the end of this unit, learners will be able to:

- Describe each of the tests for congruence SAS, SSS and ASA
- Identify the congruence of two given triangles by using the tests for congruence SAS, SSS and ASA

In previous discussion you have seen that, in order to say two triangles


are congruent, their six corresponding parts must be congruent. But, in this section, you will learn that you do not need all six pieces of information to show the triangles are congruent. By using only, the three parts of a triangle and applying the three tests for congruence (SSS, SAS, ASA), you can show whether two triangles are congruent or not.

### A. side – side – side (SSS) congruence test

#### Activity 6.1.3

*Read the following and discuss with your friends/partners*


Given  $\triangle ABC$ ,



- If only one side of is given say 5.5cm, can you draw the exact copy of it?
  - The answer is **NO**, because you can draw a number of triangles using two another arbitrary side as follows




- If only two sides of is given (say 5.5cm and 3.4cm) can you draw the exact copy of it?
  - The answer is **NO**, because you can draw a number of triangles using the two sides and another arbitrary side as follows



3. If all the three sides of  $\triangle ABC$  is given (say  $BC=5.5\text{cm}$ ,  $AC= 3.4\text{cm}$  and  $AB=5\text{cm}$ ) can you draw the exact copy of it?

In this case you can draw, a single triangle which the exact copy of  $\triangle ABC$ . So, to draw an exact copy of, we need the length of three sides of the triangle. We call this side – side – side congruence criterion.




### Side – Side – Side (SSS) congruence test:

*If the three sides of one triangle is congruent to the three corresponding sides of another triangle, then the triangles are congruent.*

#### Example 1:

examine whether the two triangles are congruent or not. If yes write the congruence relation in symbolic form



#### Solution:

$$AB=PR=3.5\text{cm} , AC=RQ=5\text{cm} \text{ and } BC=PQ=7.1\text{cm}$$


- The three sides of one triangle are equal to the three sides of other triangle, so by SSS congruence test the two triangles are congruent.
- You can easily see that A corresponds to R, B corresponds to P and C corresponds to Q,

Hence  $\triangle ABC \cong \triangle RPQ$ , by SSS congruence test

**Example 2:** consider the following kite  
 A.State the three pairs of equal parts in  $\triangle ABD$   
 and  $\triangle CBD$

B.Is  $\triangle ABD \cong \triangle CBD$ ? Why or why not?

C. Does  $\overline{BD}$  bisect  $\angle ABC$ ? Give reasons



**Solution:**

A. three pairs of equal parts in  $\triangle ABD$  and  $\triangle CBD$  are  
 $AB=CB$ ..... (given)

$AD=CD$ .....(given)


$BD=BD$ .....(common side for both)

B. From a above  $\triangle ABD \cong \triangle CBD$  by SSS

C. From b above  $\triangle ABD \cong \triangle CBD$  .... Corresponding parts of congruent triangles So,  $\angle ABC$  is bisected by  $\overline{BD}$

### Exercise 6.1.3

1. In the given figure below, lengths of the sides of the triangles are indicated. By applying the SSS congruence rule, state which pairs of triangles are congruent, in case of congruent triangles, write the result in symbolic form




2. In the figure below  $AB=AC$  and  $D$  is the mid point of  $BC$

A. State the three pairs equal parts in  $\triangle ADB$  and  $\triangle ADC$ .

B. Is  $\triangle ADB \cong \triangle ADC$ ? Give your reasons.


C. Is  $\angle B = \angle C$ ? Why?



3. In the figure,  $PR=QT$  and  $PT=QR$ .

Which of the following statements is correct?

- A.  $\triangle PQR \cong \triangle PQT$       C.  $\triangle TPQ \cong \triangle RPQ$   
 B.  $\triangle PQR \cong \triangle QPT$       D.  $\triangle PRQ \cong \triangle QTP$



## B. side – Angle – side (SSA) congruence test

### Activity 6.1.4



Read the following and discuss with your friends/partners

Given  $\triangle ABC$ ,

1. If only one side of  $\triangle ABC$  is given say 5.5cm, you can't draw the exact copy of it as you have seen in activity 5.3. can

you draw exact copy of it, if one side and one angle is given (say one side is 5.5cm and one angle is  $65^\circ$ )?


The answer is NO, because you can draw a number of triangles as follows



2. If two sides and the angle between these two sides is given (say  $BC = 5.5\text{cm}$ ,  $AC = 3.4\text{cm}$  and the angle between  $BC$  and  $AC$  is  $65^\circ$ ) can

you draw the exact copy of it?

- In this case you can draw, a single triangle which the exact copy of as follows




So, to draw an exact copy of  $\triangle ABC$ , we need the length of two sides of the triangle and the angle between the two sides. We call this side – Angle – side congruence criterion.

### Side – Angle – Side (SAS) congruence test:

*If the two sides and the angle included between them of a triangle are equal to two corresponding sides and the angle included between them of another triangle, then the triangles are congruent.*

**Example 1:** In the figure  $AB = AC$ , and  $\overline{AD}$  is the bisector of  $\angle BAC$



- State the three pairs of equal parts in  $\triangle ADB$  and  $\triangle ADC$
- Is  $\triangle ADB \cong \triangle ADC$ ? Give reason
- is  $\angle B \cong \angle C$ ? Give reason
- is  $BD = DC$ ? Give reason

**Solution:**

- The three pairs of equal parts are

$AB = AC$ ..... (Given)






$AD = AD$ ..... (common sides for both)

$m(\angle BAD) = m(\angle CAD)$ .... (  $\overline{AD}$  ) bisects  $\angle BAC$  )

- ii. yes,  $\triangle ADB \cong \triangle ADC$  (by SAS)
- iii.  $\angle B \cong \angle C$ , because they are corresponding parts of congruent triangles.
- iv.  $BD = DC$ , because they are corresponding parts of congruent triangles.


### Exercise 6.1.4

- Which angle included between side  $\overline{DE}$  and  $\overline{EF}$  in  $\triangle DEF$ ?
- By applying SAS congruence rule, you want to establish that  $\triangle PQR \cong \triangle FED$ . It is given that  $PQ = FE$  and  $RP = DF$ . What additional information is needed to be  $\triangle PQR \cong \triangle FED$ ?
- In the given figure below, by applying SAS congruence rule, state the pairs of congruent triangles. In case of congruent triangles, write them in symbolic form.



- In the given figure,  $PN = RN$  and  $TN = QN$ . Which of the following statements is correct?

- A.  $\triangle PNT \cong \triangle RNQ$   
 B.  $\triangle PNT \cong \triangle QNR$   
 C.  $\triangle TPN \cong \triangle RQN$   
 D.  $\triangle NTP \cong \triangle NQR$



### C. Angle – Side – Angle (ASA) congruence test



#### Activity 6.1.5

*Discuss with your friends/partners*

1. Can you draw the exact copy of a given triangle, if you know
  - A. Only one of its angles?
  - B. Only two of its angles?
  - C. Two angles and any one side?
  - D. Two angles and the side included between them?

### Angle – Side – Angle (ASA) congruence test:

*If two angles and the included side of a triangle are equal to two corresponding angles and the included side of another triangle, then the triangles are congruent.*

**Example 1:**  $\triangle ABC \cong \triangle QRP$  by ASA, and  $BC = RP$  is given. what additional information is needed, to establish ASA congruence test?

**Solution:**

*For ASA congruence test, we need the two angles between which the two sides  $BC$  and  $RP$  are included. So, the additional information is  $\angle B \cong \angle R$  and  $\angle C \cong \angle P$*

**Example 2:** Using the figure show that  $\triangle AOC \cong \triangle BOD$

**Solution:**

in  $\triangle AOC$ ,

$$m(\angle A) + m(\angle C) + m(\angle O) = 180^\circ$$

$$m(\angle A) + 70^\circ + 30^\circ = 180^\circ$$

$$m(\angle A) = 80^\circ$$


$m(\angle AOC) = m(\angle DOB) = 30^\circ \dots \dots$  vertically opposite angles

$$\text{in } \triangle DOB, m(\angle D) + m(\angle O) + m(\angle B) = 180^\circ$$

$$70^\circ + 30^\circ + m(\angle B) = 180^\circ \Rightarrow m(\angle B) = 80^\circ$$


Now,  $AC = DB$  and the angles that includes the sides  $AC$  and  $DB$  are also congruent (i.e.  $\angle A \cong \angle B$ , and  $\angle C \cong \angle D$ )




So, by ASA  $\triangle AOC \cong \triangle BOD$ .



### Exercise 6.1.5

1. What is the side between the angle M and N of  $\triangle MNP$ .
2. You want to establish  $\triangle DEF \cong \triangle MNP$ , using the ASA congruence rule. You are given that  $\angle D = \angle M$  and  $\angle F = \angle P$ . What additional information is needed to establish the congruence  $\triangle DEF \cong \triangle MNP$ ?
3. In figure below, by applying ASA congruence rule, state which pairs of triangles are congruent and write the result in symbolic form.






4. Given below are measurements of some parts of  $\triangle DEF$  and  $\triangle PQR$ . Check whether the two triangles are congruent or not, by ASA congruence rule and write it in symbolic form.

 **$\triangle DEF$**  **$\triangle PQR$** 

- |  |   |
|--|---|
| A. $\angle D = 70^\circ$ , $\angle F = 85^\circ$ , $DF = 7$ cm | $\angle Q = 70^\circ$ , $\angle R = 85^\circ$ , $QR = 7$ cm |
| B. $\angle D = 60^\circ$ , $\angle F = 80^\circ$ , $DF = 6$ cm | $\angle P = 60^\circ$ , $\angle R = 80^\circ$ , $PR = 6$ cm |
| C. $\angle E = 80^\circ$ , $\angle F = 30^\circ$ , $EF = 5$ cm | $\angle P = 80^\circ$ , $PQ = 5$ cm, $\angle R = 30^\circ$  |
| E. $\angle D = 80^\circ$ , $\angle F = 30^\circ$ , $DF = 5$ cm | $\angle P = 80^\circ$ , $PR = 5$ cm, $\angle R = 30^\circ$  |

5. In the figure, diagonal  $AC$  bisects  $\angle DAB$  and  $\angle DCB$ . State the three pairs of equal parts in triangles  $BAC$  and  $DAC$ .

- Is  $\triangle BAC \cong \triangle DAC$ ? Give reasons.
- $AB = AD$ ? Justify your answer.
- $DC = CB$ ? Give reasons



## 6.3. Applications

### *Learning Outcomes:*

At the end of this unit, learners will be able to:


- Apply congruency of plane figures to real life problems

In this section you learn some the applications of congruence of triangles

**Example 1:** show that the diagonal of rhombus bisects the angles at the vertex

**Solution:**

let  $ABCD$  is rhombus, we need to show  $\overline{AC}$  bisects  $\angle BAD$  and  $\angle BCD$



**Considering**

$\triangle ABC$  and  $\triangle ADC$

$AB \cong AD$  ... All sides of rhombus are congruent

$BC \cong DC$  ... All sides of rhombus are congruent

$AC \cong AC$  ... common side for both

Therefore,  $\triangle ABC \cong \triangle ADC$  by SSS congruence test

$\angle BAC \cong \angle DAC$  ... corresponding parts of congruent triangles, hence  $\angle BAD$  is bisected by  $\overline{AC}$


In similar way you can show that  $\overline{BD}$  also bisects  $\angle B$  and  $\angle D$

Example 2: the following is a part of one side of suspension bridge on Abay river

The length of cable  $AB = 5\text{m}$ ,

$BC = 4\text{m}$ ,  $EF = 4\text{m}$ ,

$AC = 3\text{m}$ , what should be the length of cable  $DE$ ? (assume  $AC \perp BC$  and  $DF \perp EF$ )



**Solution:**

consider  $\triangle ABC$  and  $\triangle DEF$

$BC = EF = 4\text{m}$  .... Given

$AC = DF = 3\text{m}$  ..... common side

$\angle C \cong \angle F$  .... Since  $AC \perp BC$  and  $DF \perp FE$ )


Therefore, by SAS test  $\triangle ABC$  and  $\triangle DEF$  and

$DE = AB = 5m$ .... Corresponding part of congruent triangles

**Example 3:** consider the following geometric figure that shows a plot of land to plant flowers.

If  $BC = CE = 6m$ ,  $\angle ABC \cong \angle CED$ ,  $AE = 4m$  and  $AC \perp BD$ , then calculate the length of

- A. CD
- B. BD



**Solution:**

A. consider  $\triangle ABC$  and  $\triangle CDE$

$BC = CE$  .... Given

$\angle ABC \cong \angle CED$ .... Given

$\angle ACB \cong \angle ECD$  ..... since  $AC \perp BD$

The corresponding two angles and included side of the two triangles are

congruent. So,  $\triangle ABC \cong \triangle DEC$  by ASA.

$CD = AC$  ... corresponding side of congruent triangles

Therefore,  $CD = AC = AE + EC = 4m + 6m = 10m$

b.  $BD = BC + CD$

$= 6m + 10m = 16m$


**Example 4:** in the figure ABCD is rectangle and F, G, E, and H are the mid points on side AB, BC, CD, and AD respectively. Show that

A.  $\triangle FAH \cong \triangle FBG$

C.  $\triangle FAH \cong \triangle EDH$

B.  $\triangle ECG \cong \triangle EDH$

D.  $HF = FG = EG = EH$



**Solution:**

A. consider  $\triangle FAH$  and  $\triangle FBG$

$FA=FB$ ..... since  $F$  is midpoint of  $AB$

$AH=HD$ .... Since  $AD = BC$  and,  $H$  is

mid point of  $AD$ ,  $G$  is midpoint of  $BC$

$m(\angle A)=m(\angle B)=90^\circ$ .... Angles of rectangle

**Therefore,**  $\triangle FAH \cong \triangle FBG$  by SAS

B. consider  $\triangle ECG$  and  $\triangle EDH$

$EC=ED$ ..... since  $E$  is midpoint of  $CD$

$CG=HD$ .... Since  $AD = BC$  and,  $H$  is midpoint of  $AD$ ,  $G$  is midpoint

of  $BC$   $m(\angle C)=m(\angle D)=90^\circ$ .... Angles of rectangle

**Therefore,**  $\triangle ECG \cong \triangle EDH$  by SAS

C. consider  $\triangle FAH$  and  $\triangle EDH$

$FA=ED$ ..... since  $AB = DC$ , and  $F$  is midpoint of  $AB$ ,  $E$  is midpoint  $DC$

$AH=HD$ .... Since  $H$  is midpoint of  $AD$

$m(\angle A)=m(\angle D)=90^\circ$ .... Angles of rectangle

**Therefore,**  $\triangle FAH \cong \triangle EDH$  by SAS


D. From A, B, and C you can see that

$$\triangle FAH \cong \triangle FBG \cong \triangle ECG \cong \triangle EDH$$

Hence,  $HF=GF=GE=HE$  since the corresponding sides of congruent triangles

### Exercise 6.2.1

1. Show that the diagonal of rectangle divides the rectangle in to two congruent triangles
2. In the figure below  $\triangle ABC$  is isosceles triangle with  $AB=BC$  and  $BD$  bisects  $\angle ABC$ , then show that  $D$  is the mid-point  $AC$ .
3. A portion of roof truss has the following shape. If  $AB = BD$ , and  $\angle ABC \cong \angle DBC$ , then show that  $\triangle ABC \cong \triangle DBC$ .



### Summary for unit 6

- ✓ Congruent figures are figures that have the same size and shape. Congruent figures are exact copies of one another.
- ✓ Two triangles are congruent, if their corresponding parts (angles and sides) that match one another are equal.
- ✓ Two triangles are congruent if they are copies of each other and when you place one triangle on another, they cover each other completely.
- ✓ If  $\triangle ABC$  is congruent to  $\triangle XYZ$ , then symbolically written as  $\triangle ABC \cong \triangle XYZ$
- ✓ If  $\triangle ABC \cong \triangle XYZ$ ,
  - i.  $\angle X \cong \angle A$ ,  $\angle Y \cong \angle B$  and  $\angle Z \cong \angle C$
  - ii.  $\overline{XY} \cong \overline{AB}$ ,  $\overline{YZ} \cong \overline{BC}$  and  $\overline{XZ} \cong \overline{AC}$
- ✓ If the three sides of one triangle is congruent to the three corresponding sides of another triangle, then the triangles are congruent by **Side – Side – Side (SSS)** congruence rule.

- ✓ If the two sides and the angle included between them of a triangle are equal to two corresponding sides and the angle included between them of another triangle, then the triangles are congruent by **Side – Angle – Side (SAS) congruence rule.**
- ✓ If two angles and the included side of a triangle are equal to two corresponding angles and the included side of another triangle, then the triangles are congruent by **Angle – Side – Angle (ASA) congruence rule.**

## Review exercise for unit 6

### *I. Write True if the statement is correct and False if it is incorrect*

1. Congruent objects are exact copies of one another.
2. Two-line segments are congruent if they have equal lengths.
3. Two angles are congruent if their measures are equal.
4. Two triangles are congruent if the three sides of the one triangle are equal to the three corresponding sides of the other.
5. Two triangles are congruent if the three angles of the one triangle are equal to the three corresponding angles of the other.
6. If two triangles are congruent, then they have the same perimeter.
7. If two squares have equal length of side, then they are congruent.
8. If  $\triangle DEF \cong \triangle RTS$ , then  $\angle E = \angle R$ .
9. Every triangle is congruent to itself.
10.  $\angle Y$  is between the side  $YZ$  and  $XZ$  of  $\triangle XYZ$ .

### *II. Choose the correct answer from the given alternatives*

11. Which congruence rule do you use in the following?  
 $AC = DF$ ,  $AB = DE$ ,  $BC = EF$ , So,  $\triangle ABC \cong \triangle DEF$   
 A. SSS      B. ASA      C. SAS      D. AAS
12. Which congruence rule do you use in the following?  
 $ZX = RP$ ,  $RQ = ZY$ ,  $\angle PRQ = \angle XZY$ , so,  $\triangle PQR \cong \triangle XYZ$

A. SSS      B. ASA      C. SAS      D. AAS

13. Which congruence rule do you use in the following?

$\angle MLN = \angle FGH$ ,  $\angle NML = \angle GFH$ ,  $ML = FG$ , So,  $\triangle LMN \cong \triangle GFH$

A. SSS      B. ASA      C. SAS      D. AAS

14. If  $\triangle ABC \cong \triangle PQR$ , then which of the following is not true?

A.  $BC = QR$     B.  $\angle C = \angle R$     C.  $\angle ABC = \angle PQR$     D.  $BC = PQ$

15. Diagonal rectangle intersects in to congruent triangles, which congruent criteria show these congruency

A. SSS      B. ASA      C. SAS      D. AAS


16. Which one of the following is not true about congruent figures?

- A. Any two rhombuses are congruent.
- B. Any two congruent figures have the same area.
- C. Any two congruent figures have the same size and shape.
- D. Any two congruent circles have the same length of radius.


### ***III. Work out***

17. Show and explain using a different congruence rule for triangle why a diagonal of a square divides the square into two congruent triangles.
18. Recall that a kite is a quadrilateral with two pairs of adjacent and congruent sides. Will one of the diagonals of a kite divide the kite into two congruent triangles? Show and explain your answer using congruence rule.
19. Two triangles ABC and PQR are such that;  $AB = 3.5$  cm,  $BC = 7.1$  cm,  $AC = 5$  cm,  $PQ = 7.1$  cm,  $QR = 5$  cm and  $PR = 3.5$  cm. Check whether the two triangles are congruent or not and explain your answer using congruence rule.


20. Given that  $\triangle ABC \cong \triangle PQR$  such that  $B = (2x + 30)^\circ$ ,  $\angle Q = 55^\circ$  and find the value of  $x$ .
21. Describe the rule of congruence in two triangles given by; In  $\triangle ABC$ ,  $AB = 7$  cm,  $BC = 5$  cm,  $\angle B = 50^\circ$  and In  $\triangle DEF$ ,  $DE = 5$  cm,  $EF = 7$  cm,  $\angle E = 50^\circ$
22.  $\triangle FGH$  is an isosceles triangle with  $FG = FH$  and  $FM$  is one of its height of triangle
- State the three pairs of equal parts in  $\triangle FGM$  and  $\triangle FHM$ .
  - is  $\triangle FGM \cong \triangle FHM$ ? Why or why not?
  - is  $\angle G = \angle H$ ? Why or why not?
  - is  $\angle GFM = \angle FHM$ ? Why or why not?
23. In the figure below, by applying congruence rule, state which pairs of triangles are congruent. In case of congruent triangles, write the result in symbolic form





A.



C



B.





# UNIT 7

# DATA HANDLING

## ***Learning Outcomes:***

At the end of this unit, learners will be able to:

- **organize data using frequency tables for a given data**
- **construct and Interpret data from pie charts**
- **calculate Mean, Mode, Median and range of a given data**
- ***apply the concept of data handling to organize and interpret real life problems***

## ***Introduction***

Collection of data from a group of things helps us to understand more about these things in the group. To do this the collected data should be presented systematically pictorially so as to analysis them. From this unit you will learn how to collect simple data and present them pictorially and do some calculation on them to study their nature or property.

## **7.1. ORGANIZATION OF DATA USING FREQUENCY TABLE**

## ***Learning Outcomes:***

At the end of this unit, learners will be able to:

- **Collect simple data from their environment using tally mark.**
- **Organizes data in a frequency distribution table.**

**Activity 7.1.1**

1. Define the word 'Data'.
2. Where do you get data?
3. The following data shows the age of students in a certain class.

15 13 13 13 15 16 14 13 14 15 12  
 17 15 13 13 14 15 13 16 14 13 14  
 15 13 16 15 16 13 12 13 15 14 14

4. Copy and Complete the table given below using the above information

Age of students	Tally	Number of students (frequency)
12		2
13		
14		
15		
16		
17		
Total number of students		

**Definition:**

***Data is a collection of facts, such as numbers, words, measurements, observations or even just descriptions of things.***

You can collect data:

- ✓ by using a questionnaire.
- ✓ by making observations and recording the results.
- ✓ by carrying out an experiment.
- ✓ from records or data base
- ✓ from the internet

One ways of presenting the data is tally chart or frequency table.

**Note:**

*A tally chart is a simple way of recording and counting frequencies*

- *A tally chart or frequency table is a quick and easy way of recording data.*
- **Frequency** is the number of times a data value occurs.

**Example 1:** Draw tally chart or frequency table using the following data. In a school, 40 students were asked what size of shoe does they dressed on Monday. Here are the results:


33 32 34 37 37 35 38 36 37 36  
 36 38 35 33 36 37 38 38 31 37  
 36 35 37 36 39 37 36 35 38 33  
 34 33 37 37 38 35 34 37 39 36


**Solution:**

Number of friends	Tally	Frequency(Num-ber of students)
31		1
32		1
33		4
34		3
35		5
36		8
37		10
38		6
39		2

**Note:**

The above tally chart or frequency table in **example 1**, shows the frequencies of the different size of shoe (how often each size occurred).

Tally marks are grouped in five to make them easier to count:  is easier to count than |||||.








- represents 1 member of the group (sample).
-  Represents 5 members of the group (sample).

**Example 2:** Consider the following data collected from the scores of 45 students in a mathematics final exam. Show this information more clearly by drawing a tally chart

36	38	32	37	34	25	28	30	33
28	32	38	32	36	34	34	38	36
32	25	37	35	29	27	32	35	37
36	31	35	36	33	24	31	34	22
27	31	29	32	37	26	34	35	32

**Solution:**

Score	Tally	Frequency (Number of students)
22		1
24		1
25		2
26		1
27		2
28		2
29		2
30		1
31		3

32		7
33		2
34		5
35		4
36		5
37		4
38		3
Total number of students		45

### Exercise 7.1.1

1. Show the following information more clearly by drawing a tally chart or frequency table.

The following are weights in kg of 40 students in a class

40    45    45    46    44    43    45    47    46    49  
 44    51    47    45    44    46    46    43    44    50  
 48    43    45    46    44    44    47    43    44    45  
 45    43    45    46    44    47    45    46    44    47

2. The table below shows the favorite color of grade 7<sup>th</sup> students.

White	Red	White	Yellow	Green	black	Green	Blue
Green	White	Black	Red	Yellow	Blue	Blue	Red
Yellow	Blue	White	Blue	Green	White	White	White
Yellow	Blue	Green	Green	White	Blue	Black	Red
Red	Blue	Yellow	Red	Green	White	White	Green

Show the above information (table) by tally chart.

3. For each of the following recorded data shows monthly average temperature in degree Celsius for Addis Ababa city, then display the information in a tally chart or frequency table.

21	19	19	20	20	20	22	22	23	22
22	21	20	19	19	20	21	22	22	21
19	21	21	22	22	23	22	23	24	23

## 7.2. CONSTRUCTION AND INTERPRETATION OF LINE GRAPHS AND PIE CHARTS

### Learning Outcomes:

At the end of this unit, learners will be able to:

- ↪ Construct line graphs and pie chart to represent organized data by using the given data or by collecting data from their environment
- ↪ Interpret simple pie charts.

### 7.2.1 Line graphs



#### Activity 7.2.1

*Discuss with your friends*

1a. Copy and complete the following table.

X	0	1	2	3	4	5
Y	1	3	5	7	9	11
(x,y)						

- b. Plot the above points on a Cartesian coordinate plane and connect by straight line.
2. The following table represents score of students. Draw the graph of this information.

<b>score</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>
<b>Number of students</b>	<b>2</b>	<b>6</b>	<b>8</b>	<b>12</b>	<b>7</b>	<b>5</b>	<b>5</b>

3. Define line graph by your own words.

### Definition:

***Line graph is a graph that uses lines to connect individual data points on a Cartesian coordinate plane.***

### Note:

*The line graph is most commonly used to represent two related facts. It use data point “markers,” which are connected by straight lines or smooth curves.*

***The following points are important to making a line graph***

1. Construct a Cartesian coordinate plane and label an appropriate scale.
2. Make a table of data arranged in order pairs and mark the points on a Cartesian coordinate plane
3. Connect the points by a straight line or smooth curve.

**Example 1:** The following table shows selling income of a shop-keeper in a week. Draw a line graph of the information given below.

<b>Days</b>	<b>Mon-day</b>	<b>Tues-day</b>	<b>Wednes-day</b>	<b>Thurs-day</b>	<b>Friday</b>	<b>Satur-day</b>	<b>Sun-day</b>
<b>Selling income in Birr</b>	<b>2000</b>	<b>2300</b>	<b>2500</b>	<b>2700</b>	<b>2500</b>	<b>2600</b>	<b>2500</b>


Solution: To draw the line graph of the above information

1. Draw a Cartesian coordinate plane and label the days horizontally

and the selling income vertically in appropriate scale.


2. Mark the points (Monday, 2000), ( Tuesday, 2300), ( Wednesday, 2500), ( Thursday, 2700), ( Thursday, 2500), (Friday), (Saturday, 2600) and (Sunday, 2500 )
3. Connect all points by smooth curve

### *Selling income*



Example 2: The graph shows the number of student's age from 13 to 20 in a School

### *Ages of students in school*



- A. What is the title of the graph?  
 B. At what age do the number of students highest?  
 C. At what age do the number of students lowest?  
 D. Prepare a table that contains ages and the number of students.

**Solution:**

A. *Ages of students in school*      B. 16      C. 20

D.

Ages	13	14	15	16	17	18	19	20
Number of students	70	95	105	140	85	70	30	10

**Exercise 7.2.1**

1. Draw a line graphs to represent each of the following data.

A. The number of letters delivered to an office in one week

<b>Days</b>	<b>Mon</b>	<b>Tue</b>	<b>Wed</b>	<b>Thu</b>	<b>Fri</b>	<b>Sat</b>	<b>Sun</b>
<b>Number of Letters</b>	<b>10</b>	<b>0</b>	<b>5</b>	<b>8</b>	<b>12</b>	<b>15</b>	<b>10</b>

B. The temperature in Addis Ababa at midday during the last week in April

<b>Days</b>	<b>Mon</b>	<b>Tue</b>	<b>Wed</b>	<b>Th</b>	<b>Fri-day</b>	<b>Sat</b>	<b>Sun</b>
<b>Temperature in °C</b>	<b>20</b>	<b>22</b>	<b>21</b>	<b>20</b>	<b>23</b>	<b>24</b>	<b>25</b>

- 2.. Copy and complete the table below gives some values between inches and centimeters. (Hint 1 inch = 2.54 centimeters).


<b>Inches</b>	<b>1</b>	<b>2</b>	<b>6</b>	<b>10</b>	<b>20</b>	<b>30</b>	<b>40</b>
<b>Centimeters</b>	<b>2.54</b>	<b>5.08</b>					

Draw the graph from of the relation between inches to centimeters.

3. The amount of petrol (in liters) in the storage tank at a garage was measured every hour between morning 1:00 and afternoon 11:30

in one day.

This is the shape of the line graph showing the results:



- When was the amount of petrol in the tank at its lowest?
- What happened to the amount of petrol between 3:00 and 4:00?
- What can you say about the sales like between 7:00 and 11:00?
- Give a reason for your answer to (question c).

## 7.2.2 Pie charts



### Activity 7.2.2

*Discuss with your friends*


- Draw a circle and divide the circle with angles at the center are  $90^\circ$ ,  $60^\circ$ ,  $30^\circ$ ,  $180^\circ$ ,
- Look at the table given below

Age of students	Below 13	13-14	15-16	17-18	Above 18
Number of students covered degree of sector	$20^\circ$	$244^\circ$	$54^\circ$	$32^\circ$	$10^\circ$

*Draw a pie chart to display the above data.*

**Note:**

The portion of a circular region enclosed between two radii and part of the circumference is called a **sector of the circle**. The size of the sector is determined by the size of the angle formed by the two radii.



- Radii means plural form of radius
- O is center of circle
- $\widehat{AB}$  is arc of the sector

**Note:**

Measure angle of circle is  $360^\circ$ .

$360^\circ$  is covered 100% of circle

$$\text{i.e } 100\% = 360^\circ \quad 1\% = 3.6^\circ \text{ and } 1^\circ = \frac{10}{36} \%$$

$$\text{Percentage} = \frac{\text{measure of angle} \times \text{total value}}{360^\circ}$$

$$\text{Measure of angle} = \frac{\text{percentage} \times 360^\circ}{\text{total value}}$$

**Example 1:** Divide the circle into 4 sectors with  $36^\circ$ ,  $54^\circ$ ,  $108^\circ$ ,  $162^\circ$

- The first sector measures  $36^\circ$   
 $= 36^\circ \times \frac{10\%}{36^\circ} = 10\%$  of a circle.
- The 2<sup>nd</sup> sector measures  $54^\circ$   
 $= 54^\circ \times \frac{10\%}{36^\circ} = 15\%$  of a circle.
- The 3<sup>rd</sup> sector measure  $108^\circ$   
 $= 108^\circ \times \frac{10\%}{36^\circ} = 30\%$  of a circle.



- The 4th sector measure  $162^\circ$   
 $= 162^\circ \times \frac{10\%}{36^\circ} = 45\%$  of a circle.

**Definition:**

**Pie chart is a type of graph that represents the data in the circular graph. It is a very common and accurate way of representing data especially useful for showing the relations of one item with another and one item with the whole items.**

**Example 2:** Show the following data by using pie chart.


The expenditure on different budget title of a family in a month is given below in table

Budget in Birr	Food	Education	Clothing	House rent	others	Saving	Total Birr
Expenditure in Birr	2400	1080	800	1800	720	400	7200

**Solution:**

Measure of angle =  $\frac{\text{Expenditure on the given budget in Birr} \times 360^\circ}{\text{Total expenditure}}$

Budget in Birr	Expenditure in Birr	Measure of angle of sectors
Food	2400	$\frac{2400 \times 360^\circ}{7200} = 120^\circ$
Education	1080	$\frac{1080 \times 360^\circ}{7200} = 54^\circ$
Clothing	800	$\frac{800 \times 360^\circ}{7200} = 40^\circ$
House rent	1800	$\frac{1800 \times 360^\circ}{7200} = 90^\circ$
others	720	$\frac{720 \times 360^\circ}{7200} = 36^\circ$
Saving	400	$\frac{400 \times 360^\circ}{7200} = 20^\circ$
Total Birr	7200	$= 360^\circ$



**Example 3:** Show the following data by using pie chart


There are 1200 students in a certain school from grade 4 up to grade 8 and the percent of students are given below

Grade	Grade 4	Grade 5	Grade 6	Grade 7	Grade 8
Number of students in %	20%	25%	15%	15%	10%

**Solution:**

$$\text{Measure of angle of sector} = \frac{\text{percentage of students} \times 360^\circ}{100}$$


Grade	Number of students in %	Measure of angle of sectors
Grade 4	20%	$\frac{20 \times 360^\circ}{100} = 72^\circ$
Grade 5	30%	$\frac{30 \times 360^\circ}{100} = 108^\circ$
Grade 6	25%	$\frac{25 \times 360^\circ}{100} = 90^\circ$
Grade 7	15%	$\frac{15 \times 360^\circ}{100} = 54^\circ$
Grade 8	10%	$\frac{10 \times 360^\circ}{100} = 36^\circ$
Total	100%	360°



**Example 3:** The pie chart given below shows W/ro Tseday's expenses and saving for last month. If the monthly income was Birr 10800, then find

- |                          |                           |
|--------------------------|---------------------------|
| A. Her food expenses     | D. Her saving             |
| B. Her house rent        | E. Her education expenses |
| C. Her clothing expenses | F. For other expenses     |

**Solution:**



$$\text{measure of angle of sector} = \frac{\text{each expenses} \times 360^\circ}{\text{Total amount of budget}}$$

$$\text{each Expenses} = \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ}$$

$$\begin{aligned} \text{A. Food expense} &= \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ} \\ &= \frac{80^\circ \times 10800}{360^\circ} = \text{Birr } 2580 \end{aligned}$$

$$\begin{aligned} \text{B. House rent} &= \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ} \\ &= \frac{70^\circ \times 10800}{360^\circ} = \text{Birr } 2100 \end{aligned}$$

$$\begin{aligned} \text{C. Clothing expense} &= \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ} \\ &= \frac{50^\circ \times 10800}{360^\circ} = \text{Birr } 1500 \end{aligned}$$

$$\begin{aligned} \text{D. Saving} &= \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ} \\ &= \frac{40^\circ \times 10800}{360^\circ} = \text{Birr } 1200 \end{aligned}$$

$$\begin{aligned} \text{E. Education} &= \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ} \\ &= \frac{60^\circ \times 10800}{360^\circ} = \text{Birr } 1800 \end{aligned}$$

$$\begin{aligned} \text{F. For other} &= \frac{\text{measure of angle of sector} \times \text{Total amount of budget}}{360^\circ} \\ &= \frac{54^\circ \times 10800}{360^\circ} = \text{Birr } 1620 \end{aligned}$$

### Exercise 7.2.2


1. 60 students were asked to name their favorite chewing gum. The results are shown in the given figure below

A. What does the whole Circle represent?

B.. Which chewing gum does the largest sector represent?

C. What does the smallest sector represent?

D. Use the given angles of sectors to calculate the number of students who liked



i. Strawberry chewing gum

ii. Chocolate

ii. Orange


2. The following pie chart shows a family budget based on a net income of Birr 8400 per month.

A.. Determine the amount spent on rent.

B. Determine the amount spent on food.

C. Determine the amount saving money.

D. How much more money is spent?



3. The budget for social development programme of wereda district is given as follows in table.

Item	Amount in Birr
Education	1,800,000
Public health	1,200,000
Community development	200,000
life skill Training for children	400,000
Total budget	3,600,000

Draw a pie chart to show the above information using percent and degree of sector.

### 7.3. THE MEAN, MODE, MEDIAN AND RANGE OF DATA

#### *Learning Outcomes:*

At the end of this unit, learners will be able to:

➤ *Calculate the mean, mode, median and range of the data.*

#### Mean

#### Activity 7.3.1

1. Find the sum of the following numbers

13, 18, 13, 14, 12

2. Divide the sum of the above numbers by five  
3. What do you say about your result

#### **Definition:**

The **mean** of a given data is the sum of all values divided by the number of values:  $\text{mean} = \frac{\text{sum of all values}}{\text{number of values}}$

**That mean of for the given data  $x_1, x_2, x_3, \dots, x_n$  is**

$$\text{mean } (\bar{X}) = \frac{(x_1 + x_2 + x_3 + \dots + x_n)}{n}, \text{ where } \bar{X} \text{ mean of data}$$

**Example1:** Find the mean of the following data

A. 22, 20, 14, 12, 27

C. 4500, 2560, 5000

B. 100, 200, 120, 320, 150, 160

**Solution:**

A.

$$\begin{aligned} \text{mean } (\bar{X}) &= \frac{(x_1 + x_2 + x_3 + \dots + x_n)}{n} \\ &= \frac{22 + 20 + 14 + 12 + 27}{5} \\ &= \frac{95}{5} = 19 \end{aligned}$$

$$\text{B. mean } (\bar{X}) = \frac{(x_1 + x_2 + x_3 + \dots + x_n)}{n}$$

$$\begin{aligned} &= \frac{100 + 200 + 120 + 320 + 150 + 160}{6} \\ &= \frac{1050}{6} = 175 \end{aligned}$$

$$\text{C. mean } (\bar{X}) = \frac{(x_1 + x_2 + x_3 + \dots + x_n)}{n}$$

$$\begin{aligned} &= \frac{4500 + 2560 + 5000}{3} \\ &= \frac{12060}{3} = 4020 \end{aligned}$$

2. The mean of five numbers is 30. Four of the numbers are 32, 28, 40 and 27, then find the value of the other numbers.

**Solution:** Let  $x$  be the missing number

$$\text{Mean } \bar{X} = \frac{\text{sum of the all data}}{\text{measure of the values}}$$

$$30 = \frac{32 + 28 + 40 + 27 + X}{5}$$

$$X + 127 = 150$$

$$X = 150 - 27$$

$$X = 23$$

3. The mean of 4 numbers is 15 and the mean of 6 other numbers is 20, then find the mean of all 10 numbers.

**Solution:**

$$\text{mean} = \frac{\text{sum of all values}}{\text{number of values}}$$

The sum of 4 numbers = mean  $\times$  number of value =  $15 \times 4 = 60$

The sum of 6 numbers = mean  $\times$  number of value =  $20 \times 6 = 120$

Thus, the total sum =  $60 + 120 = 180$

Total number of data =  $4 + 6 = 10$

$$\frac{180}{10} = 18$$

Therefore, the mean of all ten numbers is 18

### Exercise 7.3.1

- Find the mean of the following given data
  - 10 14 16 19 25 12
  - 28 35 70 140 160
  - 58 40 56 55 78 80 70 45
  - 600 750 950 800 2000
- If the age of 8 students are 11, 12, 14, 17, 15, 13, 14 and 16, then find the mean of age of students.
- The mean of four numbers is 70. Three of the numbers are 38, 85 and 60, then find the value of the other numbers
- The mean of three numbers is 18 and the mean of five other numbers is 24, then find the mean of all 8 numbers.
- Find the value of x, so that the mean of the given data 16, 2x, 6, 10 and 4 is

## Mode



### Activity 7.3.2

1. Which number occurs most frequently?

A. 6 12 10 6 23 12 6 25

B. 23 22 21 24 23 0 23 24 25 24

C. 100 500 600 700 800 900

1. Define Mode of data by your own words.

### Definition:

The **mode** of list of data is the value which occurs **most frequently**.

**Example 1:** find the mode of given data below

A. 20 30 40 20 20 30 50

B. 12 14 12 14 13 14 12

C. 25 28 50 60 70 50 60 70 90

D. 21 78 90 40 20 10 15 25 35

### Solution:

A. 20 occurs more frequently than any other values of data, then mode=20

B. 12 and 14 occurs three times, hence there are two modes **12** and **14**.

C. 50, 60 and 70 occurs two times equally, then there are two modes 50, 60 and 70.

D. Each value occurs only once, so there is no mode for the given data.

### Note:



A data that has one mode is called unimodal.



A data that has two mode is called bimodal

- ✓ A data that has three mode is called trimodal
- ✓ If each value occurs only once, so there is no mode.

### Exercise 7.3.2

1. Find the mode of the following given data below and identify it is unimodal, bimodal, trimodal and no mode.

- A. 8 11 9 14 9 15 18 6 9 10
- B. 24 15 18 20 18 22 24 26 18 26 24
- C. 35 43 39 46 51 47 38 51 43 38 40 45
- D. 123 121 119 116 130 121 131
- E. 117 119 135 121 129 139 134
- F. 7.2 6.2 7.7 8.1 6.4 7.2 5.4

### Median



#### Activity 7.3.3

Arrange the following data in increasing order

A. 14 10 12 18 17

B. 12 18 6 8 2 4 14 4 18

Circle the middle value of the above arrangement of numbers?

### Definition:

The **median** is the middle value when data is arranged in order of **size**.

*To find the median of list of data*

**I. Arrange data in increasing or decreasing order**

**II. Median= the middle value of arranged data**

**III. If the middle value is two data, then median is the mean of two middle data.**

**Example 1:** Find the median of the following data

- A. 4   5   6   10   14  
 B. 12 18   14   28   26   14   22  
 C. 8 12   18   13   16   15  
 D. 20 30   65   70   15   60   90   45

**Solution:**

A. *The increasing order of given data is*

4   5   6   10   14

*Then the middle value is 6*

*Therefore, the median is 6*

B. *The increasing order of given data is*

12   14   14   18   22   26   28

*Then the middle value is 18*

*Therefore, the median = 18*

C. *The increasing order of given data is*

8   12   13   15   16   18

*Then the middle values are 13 and 15*

The mean of 13 and 15 is  $= \frac{13 + 15}{2} = 14$

*Therefore, the median = 14*

D. *The increasing order of given data is*

15   20   30   45   60   65   70   90

*Then the middle value is 45 and 60*

The mean of 45 and 60 is  $= \frac{45 + 60}{2} = 52.5$

*Therefore, the median 52.5*

### Exercise 7.3.3

- Find the median of the following list of data
  - 14 12 24 36 23
  - 104 112 100 150 102 160
  - 250 300 100 150 220 230 90
  - 450 400 670 260 450 300 420 350 150 430
- The salaries of 8 employees in Birr who work for a small company are listed below. What is the median salary?  
8000, 9000, 5500, 10,000, 14,000, 30,000, 20,000

### Range



#### Activity 7.3.4

- identify the lowest and the highest value from the given data below
  - 26 29 60 70 80 100 20 15
  - 108 150 155 183 213
- Find the difference between the highest and the lowest of above data?
- Define range by your own words?

#### Definition:

*The **range** of the listed data is the difference between the highest value and the lowest value*

$$\text{Range} = \text{highest value} - \text{lowest value}$$

**Example 1:** find the range of the following data

- 70 55 74 63 80 40
- 800 -200 -600 0 -300
- 12 43 -55 70 10 -200 68 90

**Solution:**

A. the lowest value is 40 and the highest value is 80

$\text{Range} = \text{highest value} - \text{lowest value}$

$$\text{Range} = 80 - 40 = 40$$

B. the lowest value is -800 and the highest value is 0

$\text{Range} = \text{highest value} - \text{lowest value}$

$$\text{Range} = 0 - (-800) = 800$$

C. the lowest value is -200 and the highest value is 90

$\text{Range} = \text{highest value} - \text{lowest value}$

$$\text{Range} = 90 - (-200) = 290$$

### Exercise 7.3.4

1. Find the range of the following given data

A. 61 23 13 90 72 30 50

B. 46 70 85 84 83 92 94 58

C. -900 -300 -600 -460 0 -500 -250

D. 8.5 7.2 9.7 8.3 7.4 7.8 8.4

2. In a class of 20 students the highest score in mathematics exam was 94 and the lowest was 41. What was the range?

## 7.4. APPLICATION OF DATA HANDLING

### Learning Outcomes:

At the end of this unit, learners will be able to:

- ➔ **Apply the concept of data handling to organize and interpret real life problems.**

**Example 1:** The exam scores of 9 grade seven students are listed below.

Find the mean, mode, median and range of given data.

82, 92, 78.5, 91, 92, 89, 95, 100, 86

**Solution:**

$$\begin{aligned}
 I. \quad \text{mean } (\bar{X}) &= \frac{x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 + x_9}{9} \\
 &= \frac{82 + 92 + 78.5 + 91 + 92 + 89 + 95 + 100 + 86}{9} \\
 &= \frac{805.5}{9} = 89.5
 \end{aligned}$$

II. 92 occurs more frequently than any other values of data, then  
mode=92

III. The increasing order of data is

78.5   82   86   89   **91**   92   92   95   100

The middle value is 91, therefore the median is 91

IV. The lowest value is 78.5 and the highest value is 100

$$\text{Range} = \text{highest} - \text{lowest} = 100 - 78.5$$

$$\text{Range} = 21.5$$

**Example 2:** The ages of 10 college students are listed below. Find the mean, mode, median and range for age of students.

18, 24, 20, 35, 19, 23, 26, 23, 19, 20

**Solution:**

$$\begin{aligned}
 I. \quad \text{mean } (\bar{X}) &= \frac{x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 + x_9 + x_{10}}{n} \\
 &= \frac{18 + 24 + 20 + 35 + 19 + 23 + 26 + 23 + 19 + 20}{10} \\
 &= \frac{227}{10} = 22.7
 \end{aligned}$$

II. 19, 20, and 23 occurs twice than any other values of data, then modes are 19, 20 and 23

III. The increasing order of data is

18 19 86 89 20 20 23 23 24 26 35


The middle values are 20 and 23  $\frac{(20+23)}{2} = \frac{43}{2} = 21.5$

Therefore, the median is 21.5

IV. The lowest value is 18 and the highest value is 35

Range = highest – lowest = 35 – 18 = 17

**Example 3:** The pie chart shown below is the number of students in a certain school.



There are 1200 students in the school, then what is the number of students in grade 6, 7 and 8?

**Solution:**

$$\text{Grade 8} = 100\% - 30\% - 15\% - 20\% = 35\%$$

Grade 5	Grade 6	Grade 7	Grade 8
$= \frac{1200 \times 15}{100} = 180$	$= \frac{1200 \times 30}{100} = 360$	$= \frac{1200 \times 20}{100} = 240$	$= \frac{1200 \times 35}{100} = 420$

**Example 4:** The table shows the daily sales in Birr grocery of

different categories of items for five days.

Day	Mon	Tues	Wed	Thurs	Fri
Drinks	3000	4500	1500	4000	6500
Food	4000	5000	3500	3000	5000

- Construct a line graph for the frequency table.
- On what days were the sales for drinks better than the sales for food?
- What is the total earnings for food and drinks on Wednesday?

**Solution:**

A.



B. Thursday and Friday

C. Sale in Wednesday = Birr 1500 + Birr 3500 = Birr 5000

### Exercise 7.4.1


- The height (in centimeters) of the members of a school football team has been listed below. 142, 140, 130, 150, 160, 135, 158, 132. Find the mean, mode, median and range of the above given data.
- The table shows the daily earnings of a store for five days in Birr.
  - Construct a line graph for the frequency table.
  - On which days were the earnings above Birr 400

Day	Mon	Tues	Wed	Thurs	Fri
Earnings	300	450	200	400	650

3. The number of points scored in Ethiopian premier league of football games is listed below. Find the mean, mode, median and range of scores?

13 18 23 9 3 18

4. 3000 students appeared for an examination from five different centers C1, C2, C3, C4 and C5 of a city. From the given pie chart, find the number of students appearing for the examination from each center



## SUMMARY FOR UNIT 7

- ✓ A tally chart or frequency table is a quick and easy way of recording data.
- ✓ Line graph is a graph that uses lines to connect individual data points on a Cartesian coordinate plane
- ✓ The line graph is most commonly used to represent two related facts. It uses data point “markers,” which are connected by straight lines or smooth curves
- ✓ Pie chart is a type of graph that represents the data in the circular Graph.
- ✓ The portion of a circular region enclosed between two radii and part of the circumference (an arc) is called a **sector of the circle**

- ✓ The **mean** of given data is the sum of all values divided by the

$$\text{mean } (\bar{X}) = \frac{\text{sum of all values}}{\text{number of values}}$$

- ✓ The **mode** of list of data is the value which occurs **most**
- ✓ A data that has one mode is called unimodal.
- ✓ A data that has two mode is called bimodal
- ✓ A data that has three mode is called trimodal
- ✓ If each value occurs only once, so there is no mode
- ✓ The **median** is the middle value when data is arranged in order of **size**.
- ✓ That is range=highest value –lowest value

## REVIEW EXERCISE FOR UNIT 7

*I. Write True if the statement is correct and False if the statement is incorrect*


- 1 Mean is the difference between maximum and minimum value.
2. The mean of 1,3,5 and 7 is 5.
3. A data which has two modes is called bimodal
4. Pie chart is called circular graph.
5.  $360^\circ$  is 100% of a circle.

### II. Work out

6. 1440 students were asked how they travelled to school. The pie chart shows the results of this survey;

Find


- A. How many of the students travelled to school by bus.
- B. How many students travelled on foot?



7. The total expenditure of a sub city is Birr 720,000,000. The pie chart as shown below then how much

money was spent on

- A. Education
- B. Public health
- C. Social service
- D. Others



8. If the mean of 7 numbers is 20 and the mean of 5 other numbers is 44, then find the mean of all 12 numbers?

9. Find the mean median, mode and range of each set of numbers below.

- A. 3, 4, 7, 3, 5, 2, 6, 10
- B. 8, 10, 12, 14, 7, 16, 5, 7, 9, 11
- C. 17, 18, 16, 17, 17, 14, 22, 15, 16, 17, 14, 12

10. The weights, in kilograms, of the 8 sport club members at a school are: 45, 53, 47, 46, 44, 46, 47, 48

- A. What is the modal weight?
- B. Calculate the mean and range of weight.

11. Students are arranged in eleven classes. The class sizes are 23, 24, 24, 26, 27, 28, 30, 24, 29, 24, 27

- A. What is the modal class size?
- B. Calculate the mean class size.

12. Find mean, mode, median of the following number

- A. 49 70 24 76 97 32 32 31
- B. 73 93 72 44 16 43 22 29


13. If the mean of  $x$ ,  $x+3$ ,  $x-5$ ,  $2x$  and  $3x$  is 8, then find the value  $x$ ?

14. 1200 students were interviewed to find out which of five clubs they Preferred. The results as follows

Club	Girls club	HIV club	Maths club	Sport club	Mini media club
Number of students	180	150	240	270	360

Draw a pie chart to represent the above information.

15. Use the information given to find the value of  $n$  in each of the following of numbers.
- A. 5, 7, 4, 1,  $n$ , 5: the mean is 6    D. 2.6, 3.5,  $n$ , 6.2: the mean is 4
- B. 3, 1, 4, 5, 4,  $n$ : the mode is 4
- C. 1, 7, 2, 1,  $n$ , 4, 3: the modes are 1 and 2
16. The line graph given below shows weekly recorded Addis Ababa's temperature in degree Celsius



- A. What is the range of temperature?
- B. What is mean of temperature?
- C. When was the temperature highest?